



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:21 AM UTC

PDB ID : 1S38 / pdb_00001s38
Title : CRYSTAL STRUCTURE OF TGT IN COMPLEX WITH 2-AMINO-8-MET
HYLQUINAZOLIN-4(3H)-ONE
Authors : Meyer, E.A.; Furler, M.; Diederich, F.; Brenk, R.; Klebe, G.
Deposited on : 2004-01-12
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	5% 75% 17% 6%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3223 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA guanine transglycosylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	364	2865	1795	520	528	22	27	3	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P28720

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 2-AMINO-8-METHYLQUINAZOLIN-4(3H)-ONE (CCD ID: MAQ) (formula: C₉H₉N₃O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			13	9	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	344	Total	O	0	0
			344	344		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: tRNA guanine transglycosylase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.60Å 65.48Å 70.43Å 90.00° 96.53° 90.00°	Depositor
Resolution (Å)	20.00 – 1.81 20.00 – 1.81	Depositor EDS
% Data completeness (in resolution range)	93.6 (20.00-1.81) 93.6 (20.00-1.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 1.81Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.197 , 0.222 0.199 , 0.223	Depositor DCC
R_{free} test set	3563 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3223	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MAQ, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	1/2924 (0.0%)	0.88	9/3934 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	131	SER	C-N	-5.75	1.25	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131	SER	O-C-N	-7.95	110.28	123.00
1	A	51	VAL	N-CA-C	-7.29	96.97	107.98
1	A	131	SER	CA-C-N	6.94	134.79	121.54
1	A	131	SER	C-N-CA	6.94	134.79	121.54
1	A	47	THR	N-CA-C	5.95	118.28	111.02
1	A	325	LYS	N-CA-C	5.76	120.68	113.43
1	A	190	LYS	N-CA-C	5.39	117.58	111.11
1	A	205	SER	CB-CA-C	-5.36	110.37	116.54
1	A	52	LYS	N-CA-C	5.15	117.94	110.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2865	0	2815	58	0
2	A	1	0	0	0	0
3	A	13	0	9	0	0
4	A	344	0	0	9	0
All	All	3223	0	2824	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:TYR:HA	1:A:75:MET:HE3	1.39	1.01
1:A:263[A]:GLY:HA3	1:A:281[A]:CYS:SG	2.15	0.86
1:A:71:THR:HG22	1:A:75:MET:HE2	1.59	0.85
1:A:131:SER:OG	4:A:1272:HOH:O	2.03	0.76
1:A:297:ASP:OD1	1:A:380:ARG:HD3	1.92	0.68
1:A:190:LYS:N	1:A:190:LYS:HD2	2.10	0.65
1:A:72:TYR:HA	1:A:75:MET:CE	2.21	0.64
1:A:188:SER:HB2	4:A:1135:HOH:O	1.96	0.63
1:A:38:ARG:O	1:A:41:ALA:HB2	1.99	0.62
1:A:190:LYS:HD2	1:A:190:LYS:H	1.65	0.61
1:A:263[B]:GLY:O	1:A:264[B]:LYS:HB3	1.98	0.61
1:A:39:THR:HA	1:A:40:PRO:C	2.27	0.60
1:A:43:MET:HG2	1:A:66:ILE:HG23	1.84	0.60
1:A:145:HIS:CD2	1:A:189:ARG:HD3	2.37	0.59
1:A:124:PHE:O	1:A:132:ARG:HA	2.04	0.57
1:A:47:THR:HB	4:A:1362:HOH:O	2.03	0.57
1:A:75:MET:HE1	1:A:135:LEU:HD22	1.86	0.57
1:A:359:GLN:NE2	1:A:362:ARG:HH21	2.03	0.56
1:A:190:LYS:HD3	4:A:1076:HOH:O	2.06	0.55
1:A:75:MET:HE1	1:A:135:LEU:CD2	2.36	0.55
1:A:71:THR:HG23	1:A:143:ILE:HG21	1.90	0.53
1:A:190:LYS:H	1:A:190:LYS:CD	2.20	0.53
1:A:72:TYR:O	1:A:75:MET:HG2	2.08	0.53
1:A:252:PRO:O	1:A:257:HIS:HE1	1.92	0.52
1:A:320:CYS:O	1:A:324:GLN:HG3	2.10	0.52
1:A:303:ARG:HD2	4:A:1162:HOH:O	2.10	0.52
1:A:47:THR:HG22	1:A:48:ALA:N	2.25	0.51
1:A:124:PHE:CD1	1:A:124:PHE:N	2.80	0.50
1:A:281[A]:CYS:SG	1:A:284:PRO:HD3	2.52	0.49
1:A:52:LYS:HE2	4:A:1362:HOH:O	2.14	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:THR:O	1:A:62:THR:HG23	2.14	0.47
1:A:306:ARG:NH2	4:A:1127:HOH:O	2.46	0.47
1:A:319:HIS:H	1:A:356:GLN:HE22	1.61	0.47
1:A:322:VAL:HG13	1:A:345:LEU:HG	1.97	0.47
1:A:97:ARG:HB3	1:A:98:PRO:CD	2.45	0.47
1:A:319:HIS:HA	4:A:1215:HOH:O	2.15	0.47
1:A:176:MET:SD	1:A:218:LEU:HD13	2.56	0.46
1:A:186:PHE:CD2	1:A:198:LEU:HG	2.51	0.46
1:A:257:HIS:HD2	1:A:275:GLY:O	1.98	0.45
1:A:304:ASN:OD1	1:A:306:ARG:HG2	2.15	0.45
1:A:47:THR:HG23	1:A:73:HIS:CG	2.51	0.45
1:A:190:LYS:N	1:A:190:LYS:CD	2.78	0.45
1:A:241:PHE:CZ	1:A:270:GLY:HA3	2.51	0.45
1:A:263[A]:GLY:O	1:A:354:TYR:CE2	2.70	0.44
1:A:68:LEU:C	1:A:68:LEU:HD23	2.42	0.44
1:A:161:TYR:CD1	1:A:162:PRO:HA	2.52	0.44
1:A:330:TYR:CE2	1:A:334:LEU:HD11	2.53	0.44
1:A:47:THR:HG23	1:A:73:HIS:CD2	2.53	0.43
1:A:97:ARG:HB3	1:A:98:PRO:HD2	2.00	0.42
1:A:293:ALA:HB2	1:A:346:MET:HE1	2.02	0.42
1:A:21:ARG:HD3	4:A:1159:HOH:O	2.19	0.42
1:A:263[B]:GLY:O	1:A:267:ASP:HB2	2.20	0.42
1:A:259:LEU:HG	1:A:262:VAL:CG2	2.50	0.41
1:A:283:LEU:HB3	1:A:284:PRO:HD3	2.02	0.41
1:A:284:PRO:HG3	1:A:354:TYR:CZ	2.55	0.41
1:A:314:LEU:HD23	1:A:349:HIS:CG	2.56	0.41
1:A:280:ASP:O	1:A:281[A]:CYS:HB3	2.21	0.41
1:A:285:THR:HG22	1:A:347:THR:HG23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/386 (94%)	347 (96%)	12 (3%)	2 (1%)	21	11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	264[A]	LYS
1	A	264[B]	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/315 (95%)	290 (97%)	9 (3%)	36	17

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	THR
1	A	86	LEU
1	A	107	GLN
1	A	132	ARG
1	A	134	MET
1	A	198	LEU
1	A	218	LEU
1	A	314	LEU
1	A	345	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	145	HIS
1	A	202	GLN
1	A	212	GLN
1	A	213	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	257	HIS
1	A	324	GLN
1	A	356	GLN
1	A	359	GLN
1	A	372	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MAQ	A	900	-	14,14,14	2.47	9 (64%)	20,20,20	1.67	4 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAQ	A	900	-	-	-	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	900	MAQ	C4-C7	4.15	1.55	1.48
3	A	900	MAQ	C3-C2	3.96	1.45	1.40
3	A	900	MAQ	C4-C3	3.04	1.45	1.41
3	A	900	MAQ	C6-C1	3.01	1.44	1.38
3	A	900	MAQ	C6-C5	2.92	1.43	1.38
3	A	900	MAQ	C5-C4	2.40	1.43	1.39
3	A	900	MAQ	C9-C2	2.30	1.55	1.51
3	A	900	MAQ	C8-N3	-2.21	1.32	1.37
3	A	900	MAQ	C1-C2	2.02	1.43	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	900	MAQ	C3-C4-C7	-4.66	114.39	119.99
3	A	900	MAQ	C9-C2-C3	-2.47	118.42	121.42
3	A	900	MAQ	O1-C7-N1	2.34	123.78	120.23
3	A	900	MAQ	O1-C7-C4	-2.32	117.59	121.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	364/386 (94%)	0.12	21 (5%)	29 32	10, 20, 34, 60	11 (3%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	263[A]	GLY	8.1
1	A	108	VAL	4.7
1	A	131	SER	4.6
1	A	106	TYR	4.4
1	A	125	LYS	3.9
1	A	319	HIS	3.6
1	A	123	THR	3.4
1	A	124	PHE	3.2
1	A	111	LEU	2.9
1	A	47	THR	2.8
1	A	134	MET	2.6
1	A	133	HIS	2.5
1	A	105	GLY	2.3
1	A	135	LEU	2.3
1	A	109	MET	2.3
1	A	11	ARG	2.3
1	A	382	PHE	2.2
1	A	71	THR	2.2
1	A	264[A]	LYS	2.1
1	A	324	GLN	2.1
1	A	72	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MAQ	A	900	13/13	0.92	0.09	21,25,28,29	0
2	ZN	A	400	1/1	1.00	0.03	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.