



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 11:12 AM UTC

PDB ID : 6S3D / pdb_00006s3d
Title : Structure of D25 Fab in complex with scaffold S0_2.126
Authors : Cramer, J.T.; Krey, T.
Deposited on : 2019-06-25
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	241	74% 14% 12%
1	C	241	79% 10% 11%
1	E	241	2% 73% 16% 11%
1	H	241	4% 72% 16% 12%
2	B	225	2% 80% 13% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	225	% 74% 17% 8%
2	F	225	3% 76% 15% 8%
2	L	225	2% 78% 16% 6%
3	M	71	% 63% 20% 17%
3	N	71	61% 21% 18%
3	O	71	62% 18% •• 17%
3	P	71	% 61% 21% 18%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14439 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	0	0	0
			1572	1002	259	304	7			
1	C	215	Total	C	N	O	S	0	0	0
			1601	1020	265	309	7			
1	E	215	Total	C	N	O	S	0	0	0
			1590	1012	264	307	7			
1	H	212	Total	C	N	O	S	0	0	0
			1581	1006	261	307	7			




- Molecule 2 is a protein called Light Chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	210	Total	C	N	O	S	0	0	0
			1580	991	260	324	5			
2	D	206	Total	C	N	O	S	0	0	0
			1570	986	260	319	5			
2	F	206	Total	C	N	O	S	0	0	0
			1553	972	256	320	5			
2	L	211	Total	C	N	O	S	0	0	0
			1585	992	261	327	5			

- Molecule 3 is a protein called S0_2.126.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	59	Total	C	N	O	S	0	0	0
			444	277	77	85	5			
3	N	58	Total	C	N	O	S	0	0	0
			456	284	80	87	5			
3	O	59	Total	C	N	O	S	0	0	0
			452	280	79	88	5			
3	P	58	Total	C	N	O	S	0	0	0
			455	282	82	87	4			

- Molecule 1: Heavy Chain









• Molecule 2: Light Chain





• Molecule 2: Light Chain



• Molecule 2: Light Chain




• Molecule 2: Light Chain



• Molecule 3: S0_2.126



• Molecule 3: S0_2.126



• Molecule 3: S0_2.126



• Molecule 3: S0_2.126



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	126.27Å 126.98Å 156.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.09 – 3.00 49.09 – 3.00	Depositor EDS
% Data completeness (in resolution range)	98.8 (49.09-3.00) 99.7 (49.09-3.00)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.14 _3260	Depositor
R, R_{free}	0.269 , 0.292 0.271 , 0.294	Depositor DCC
R_{free} test set	2542 reflections (2.91%)	wwPDB-VP
Wilson B-factor (Å ²)	73.4	Xtriage
Anisotropy	0.646	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 87.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14439	wwPDB-VP
Average B, all atoms (Å ²)	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 88.81 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.2895e-08. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.18	0/1612	0.45	0/2211
1	C	0.19	0/1642	0.44	0/2250
1	E	0.29	1/1629 (0.1%)	0.50	0/2232
1	H	0.16	0/1621	0.43	0/2221
2	B	0.18	0/1612	0.43	0/2199
2	D	0.19	0/1600	0.46	0/2175
2	F	0.17	0/1583	0.44	0/2158
2	L	0.17	0/1617	0.42	0/2206
3	M	0.24	0/447	0.51	0/596
3	N	0.25	0/460	0.57	0/612
3	O	0.43	0/456	0.95	1/609 (0.2%)
3	P	0.24	0/459	0.56	0/612
All	All	0.21	1/14738 (0.0%)	0.48	1/20081 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	172	GLY	C-O	5.81	1.29	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	O	29	GLU	N-CA-C	-5.20	105.30	110.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1572	0	1525	21	0
1	C	1601	0	1562	15	0
1	E	1590	0	1539	24	0
1	H	1581	0	1546	22	0
2	B	1580	0	1503	19	0
2	D	1570	0	1502	29	0
2	F	1553	0	1454	24	0
2	L	1585	0	1506	24	0
3	M	444	0	440	11	0
3	N	456	0	470	8	0
3	O	452	0	451	11	0
3	P	455	0	463	9	0
All	All	14439	0	13961	204	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 204 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:O:26:ARG:O	3:O:29:GLU:HB2	1.54	1.08
1:E:159:ASP:HB3	1:E:190:LEU:HD22	1.55	0.87
3:O:28:GLU:HA	3:O:31:GLU:HG3	1.62	0.79
1:E:14:LYS:HG3	1:E:20:VAL:HG12	1.68	0.74
3:O:26:ARG:CZ	3:O:28:GLU:HB2	2.18	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/241 (86%)	199 (96%)	8 (4%)	1 (0%)	24	60
1	C	211/241 (88%)	200 (95%)	11 (5%)	0	100	100
1	E	209/241 (87%)	201 (96%)	8 (4%)	0	100	100
1	H	208/241 (86%)	201 (97%)	6 (3%)	1 (0%)	24	60
2	B	206/225 (92%)	197 (96%)	8 (4%)	1 (0%)	24	60
2	D	198/225 (88%)	192 (97%)	5 (2%)	1 (0%)	24	60
2	F	198/225 (88%)	192 (97%)	5 (2%)	1 (0%)	24	60
2	L	209/225 (93%)	203 (97%)	5 (2%)	1 (0%)	24	60
3	M	57/71 (80%)	54 (95%)	2 (4%)	1 (2%)	6	31
3	N	56/71 (79%)	51 (91%)	5 (9%)	0	100	100
3	O	57/71 (80%)	53 (93%)	3 (5%)	1 (2%)	6	31
3	P	56/71 (79%)	51 (91%)	5 (9%)	0	100	100
All	All	1873/2148 (87%)	1794 (96%)	71 (4%)	8 (0%)	30	65

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	32	VAL
3	O	23	GLY
2	D	32	VAL
2	B	32	VAL
2	L	32	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	173/205 (84%)	173 (100%)	0	100	100
1	C	178/205 (87%)	178 (100%)	0	100	100
1	E	174/205 (85%)	173 (99%)	1 (1%)	78	88
1	H	177/205 (86%)	177 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	177/198 (89%)	177 (100%)	0	100	100
2	D	177/198 (89%)	176 (99%)	1 (1%)	78	88
2	F	172/198 (87%)	172 (100%)	0	100	100
2	L	177/198 (89%)	177 (100%)	0	100	100
3	M	47/63 (75%)	47 (100%)	0	100	100
3	N	52/63 (82%)	52 (100%)	0	100	100
3	O	50/63 (79%)	46 (92%)	4 (8%)	11	38
3	P	51/63 (81%)	51 (100%)	0	100	100
All	All	1605/1864 (86%)	1599 (100%)	6 (0%)	84	90

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	O	29	GLU
3	O	32	GLU
3	O	33	ARG
1	E	158	LYS
2	D	156	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	140	ASN
3	O	14	GLN
2	F	29	GLN
2	F	39	GLN
2	F	162	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	212/241 (87%)	0.18	2 (0%) 81 61	46, 89, 135, 169	0
1	C	215/241 (89%)	0.32	3 (1%) 73 51	51, 100, 165, 200	0
1	E	215/241 (89%)	0.31	6 (2%) 55 32	46, 101, 160, 194	0
1	H	212/241 (87%)	0.55	9 (4%) 40 22	50, 103, 190, 249	0
2	B	210/225 (93%)	0.14	4 (1%) 66 43	52, 84, 159, 176	0
2	D	206/225 (91%)	0.04	3 (1%) 72 49	36, 80, 149, 166	0
2	F	206/225 (91%)	0.18	6 (2%) 53 31	50, 85, 189, 230	0
2	L	211/225 (93%)	0.17	4 (1%) 66 43	46, 86, 172, 190	0
3	M	59/71 (83%)	0.47	1 (1%) 69 45	89, 130, 160, 163	0
3	N	58/71 (81%)	0.21	0 100 100	76, 113, 148, 156	0
3	O	59/71 (83%)	0.25	0 100 100	82, 131, 152, 168	0
3	P	58/71 (81%)	0.20	1 (1%) 69 45	74, 114, 141, 146	0
All	All	1921/2148 (89%)	0.24	39 (2%) 65 41	36, 95, 165, 249	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	138	PRO	4.8
1	H	140	ALA	4.7
1	H	215	PRO	4.0
1	E	140	ALA	3.8
2	D	212	ASN	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.