



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:31 PM UTC

PDB ID : 3S4A / pdb_00003s4a
Title : Cellobiose phosphorylase from *Cellulomonas uda* in complex with cellobiose
Authors : Van Hoorebeke, A.; Stout, J.; Soetaert, W.; Van Beeumen, J.; Desmet, T.; Savvides, S.
Deposited on : 2011-05-19
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	822	5% 89% 11%
1	B	822	6% 88% 11%
2	C	2	100%
2	D	2	100%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 26625 atoms, of which 12150 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellobiose phosphorylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	822	Total	C	H	N	O	S	0	3	0
			12471	4072	6026	1111	1245	17			
1	B	822	Total	C	H	N	O	S	9	5	0
			12559	4091	6082	1112	1257	17			

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	H	O	0	0	0
			44	12	21	11			
2	D	2	Total	C	H	O	0	0	0
			44	12	21	11			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	745	Total	O	0	0
			745	745		
3	B	762	Total	O	0	0
			762	762		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cellobiose phosphorylase



- Molecule 1: Cellobiose phosphorylase



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain C:  100%

BGC1
BGC2

- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain D:  100%

BGC1
BGC2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.69Å 195.56Å 103.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.79 – 1.99 45.79 – 1.99	Depositor EDS
% Data completeness (in resolution range)	88.2 (45.79-1.99) 99.0 (45.79-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 1.98Å)	Xtriage
Refinement program	PHENIX 1.6.4_486	Depositor
R, R_{free}	0.200 , 0.255 0.209 , 0.254	Depositor DCC
R_{free} test set	5954 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	10.0	Xtriage
Anisotropy	0.596	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.57$, $\langle L^2 \rangle = 0.42$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	26625	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 61.30 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3373e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	4/6622 (0.1%)	0.95	5/9029 (0.1%)
1	B	0.99	3/6657 (0.0%)	0.95	4/9075 (0.0%)
All	All	0.99	7/13279 (0.1%)	0.95	9/18104 (0.0%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	191	TYR	C-O	7.07	1.32	1.24
1	A	650	VAL	CA-CB	6.69	1.61	1.54
1	A	192	ARG	C-N	-6.42	1.24	1.33
1	A	741	VAL	CA-CB	5.63	1.61	1.54
1	B	561	THR	CA-C	5.44	1.55	1.52
1	B	706	GLU	C-N	5.41	1.38	1.33
1	B	677	VAL	CA-CB	5.23	1.60	1.54

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	765	VAL	CA-C-N	6.61	126.31	119.56
1	A	765	VAL	C-N-CA	6.61	126.31	119.56
1	A	793	GLY	N-CA-C	6.21	118.63	111.36
1	B	762	GLY	CA-C-N	5.88	125.50	119.56
1	B	762	GLY	C-N-CA	5.88	125.50	119.56
1	B	191	TYR	CA-C-O	5.78	126.61	119.97
1	A	222	ASN	N-CA-C	5.51	119.04	110.17
1	A	668	ASN	CA-C-O	5.35	123.53	118.34
1	B	654	PRO	O-C-N	5.30	123.75	121.31

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6445	6026	6050	59	0
1	B	6477	6082	6087	69	0
2	C	23	21	21	0	0
2	D	23	21	21	0	0
3	A	745	0	0	4	0
3	B	762	0	0	6	0
All	All	14475	12150	12179	126	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (126) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:504:SER:O	1:B:508[B]:THR:HG22	1.80	0.82
1:A:51:LYS:HG2	1:A:52:MET:HG2	1.63	0.80
1:A:51:LYS:HE3	1:A:158:GLN:OE1	1.88	0.73
1:B:633:MET:H	1:B:668:ASN:HD21	1.37	0.70
1:B:503:GLU:HB3	1:B:508[A]:THR:HG21	1.74	0.69
1:A:31:GLU:O	1:A:34:PHE:CZ	2.46	0.69
1:B:715:ALA:O	1:B:722:HIS:HD2	1.76	0.68
1:B:198:TYR:CE2	1:B:286[B]:VAL:HG11	2.30	0.66
1:A:51:LYS:HD2	1:A:360:ILE:O	1.98	0.64
1:A:633:MET:H	1:A:668:ASN:HD21	1.47	0.63
1:B:28:LEU:HB2	1:B:35:SER:HB2	1.79	0.63
1:B:794:ARG:NH1	3:B:1217:HOH:O	2.33	0.62
1:A:347:PHE:HE2	1:A:383:ARG:NH2	1.97	0.62
1:B:198:TYR:CE2	1:B:286[B]:VAL:CG1	2.82	0.62
1:B:22:TYR:CG	1:B:23:PRO:HD2	2.34	0.61
1:A:322:TYR:CD2	1:A:772:ARG:HD3	2.35	0.61
1:A:668:ASN:N	1:A:669:PRO:CD	2.66	0.58
1:A:190:GLU:HG2	1:A:193:GLU:CD	2.31	0.56
1:A:795:LEU:HD22	1:A:820:VAL:HG22	1.86	0.55
3:A:981:HOH:O	1:B:192:ARG:HD3	2.04	0.55
1:B:496:CYS:SG	1:B:508[B]:THR:HG23	2.45	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:191:TYR:CE2	1:B:286[B]:VAL:HG22	2.42	0.55
1:A:406:GLN:HE21	1:A:413:ASN:HB2	1.71	0.55
1:B:278:TRP:CD1	1:B:283:LYS:HG2	2.41	0.55
1:B:22:TYR:CD2	1:B:23:PRO:HD2	2.42	0.55
1:B:402:TYR:CG	1:B:412:GLY:HA3	2.41	0.54
1:A:795:LEU:HD23	1:A:820:VAL:HG13	1.88	0.54
1:B:716:GLY:C	1:B:722:HIS:CD2	2.85	0.54
1:A:429:ALA:HB2	1:A:528:TYR:CZ	2.44	0.52
1:B:539:ARG:HA	3:B:1497:HOH:O	2.08	0.52
1:A:46:PHE:HZ	1:A:349:MET:HE1	1.75	0.52
1:B:317:ASP:HB2	3:B:972:HOH:O	2.10	0.52
1:B:633:MET:H	1:B:668:ASN:ND2	2.08	0.51
1:B:329:GLU:HG3	3:B:1329:HOH:O	2.09	0.51
1:A:402:TYR:CG	1:A:412:GLY:HA3	2.46	0.51
1:B:794:ARG:HH21	1:B:821:THR:HG22	1.75	0.51
1:A:743:GLN:HB3	1:A:749:ARG:HB3	1.93	0.51
1:A:485:ARG:HD3	1:A:506:GLN:O	2.11	0.51
1:A:571:ARG:HG2	1:A:591:TRP:CD2	2.47	0.50
1:A:593:GLU:N	1:A:594:PRO:CD	2.75	0.50
1:B:382:GLU:HG2	1:B:383:ARG:N	2.26	0.50
1:B:768:PHE:CE1	1:B:783:VAL:HG21	2.47	0.50
1:A:666:HIS:HB2	3:A:924:HOH:O	2.13	0.48
1:B:705:LEU:HD11	1:B:726:LYS:HA	1.95	0.48
1:A:771:THR:HG23	1:A:780:GLU:OE2	2.13	0.48
1:A:46:PHE:CZ	1:A:349:MET:HE1	2.48	0.48
1:B:504:SER:O	1:B:508[A]:THR:HG23	2.13	0.48
1:A:52:MET:HA	1:A:156:ASN:HA	1.95	0.48
1:A:657:TYR:HB3	1:B:169:SER:HB3	1.96	0.48
1:A:101:LEU:HD22	1:A:341:TYR:CG	2.48	0.48
1:A:683:ARG:HD2	1:A:687:TYR:CZ	2.49	0.47
1:B:487:ASP:O	1:B:488:TRP:C	2.56	0.47
1:B:668:ASN:N	1:B:669:PRO:CD	2.77	0.47
1:B:716:GLY:O	1:B:722:HIS:HB2	2.14	0.47
1:B:1:MET:CE	3:B:1282:HOH:O	2.62	0.47
1:B:136:ASN:ND2	1:B:142:LYS:HG2	2.30	0.47
1:B:686:ASP:O	1:B:690:ARG:HG3	2.15	0.47
1:A:101:LEU:HD23	1:A:341:TYR:CD2	2.50	0.47
1:B:504:SER:O	1:B:508[B]:THR:CG2	2.59	0.46
1:B:53:ARG:HA	1:B:53:ARG:HD3	1.70	0.46
1:B:191:TYR:CD2	1:B:286[B]:VAL:HG22	2.51	0.46
1:B:405:TYR:O	1:B:407:PRO:HD3	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:PHE:O	1:A:237:ASN:HA	2.16	0.46
1:B:758:ASP:N	1:B:759:PRO:HD3	2.31	0.46
1:A:711:ALA:HA	1:A:728:SER:HA	1.98	0.45
1:B:693:PRO:HA	3:B:1357:HOH:O	2.16	0.45
3:A:1026:HOH:O	1:B:713:MET:HE2	2.16	0.45
1:B:483:ILE:HG21	1:B:494:LEU:HD12	1.99	0.45
1:A:703:HIS:O	1:A:704:ARG:HB2	2.16	0.45
1:B:795:LEU:HD22	1:B:820:VAL:HG22	1.99	0.44
1:A:483:ILE:HG21	1:A:494:LEU:HD12	1.98	0.44
1:B:656:GLY:HA3	1:B:719:ALA:HB2	1.99	0.44
1:B:74:VAL:O	1:B:80:VAL:HA	2.17	0.44
1:B:210:PHE:O	1:B:237:ASN:HA	2.17	0.44
1:B:224:LEU:HD23	1:B:224:LEU:HA	1.77	0.44
1:B:715:ALA:O	1:B:722:HIS:CD2	2.63	0.44
1:A:527:LEU:O	1:A:531:GLN:HG3	2.17	0.44
1:A:761:ILE:HG21	1:A:765:VAL:HB	1.99	0.44
1:B:448:VAL:HG12	1:B:463:HIS:CE1	2.53	0.44
1:B:22:TYR:CG	1:B:23:PRO:CD	3.01	0.43
1:A:705:LEU:HD11	1:A:726:LYS:HA	1.99	0.43
1:B:269:TYR:CE2	1:B:271:GLU:HG3	2.54	0.43
1:A:568:TRP:CZ2	1:A:583:ASP:HB2	2.53	0.43
1:A:191:TYR:CD2	1:A:286:VAL:HG12	2.54	0.43
1:B:728:SER:O	1:B:729:TRP:HB2	2.18	0.43
1:A:347:PHE:HE2	1:A:383:ARG:CZ	2.31	0.42
1:A:656:GLY:HA3	1:A:719:ALA:HB2	2.01	0.42
1:B:113:VAL:HG23	1:B:142:LYS:HE3	2.00	0.42
1:B:347:PHE:HB2	1:B:376:PHE:CD2	2.54	0.42
1:A:794:ARG:NH2	1:A:821:THR:HG22	2.35	0.42
1:B:496:CYS:SG	1:B:508[B]:THR:CG2	3.08	0.42
1:A:166:ARG:HD2	1:B:649:GLU:OE2	2.19	0.42
1:A:146:LEU:HD12	1:A:253:VAL:HG21	2.02	0.42
1:B:70:ARG:N	1:B:70:ARG:HD2	2.34	0.42
1:B:190:GLU:O	1:B:193:GLU:HG3	2.19	0.42
1:B:593:GLU:HB2	1:B:594:PRO:HD3	2.01	0.42
1:A:60:TYR:CZ	1:A:358:THR:HA	2.55	0.42
1:A:337:ILE:HG13	3:A:1260:HOH:O	2.20	0.42
1:A:458:VAL:HB	1:A:459:PRO:HD2	2.02	0.42
1:B:649:GLU:O	1:B:650:VAL:C	2.63	0.41
1:B:711:ALA:HA	1:B:728:SER:HA	2.02	0.41
1:A:55:LEU:HD12	1:A:55:LEU:N	2.35	0.41
1:A:229:VAL:HB	1:A:230:PRO:HD3	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:749:ARG:HA	1:A:750:PRO:HD3	1.73	0.41
1:B:113:VAL:CG2	1:B:142:LYS:HE3	2.50	0.41
1:A:52:MET:SD	1:A:156:ASN:ND2	2.94	0.41
1:A:535:LEU:HD23	1:A:535:LEU:HA	1.93	0.41
1:A:666:HIS:CE1	1:A:732:GLY:HA3	2.56	0.41
1:A:429:ALA:HB2	1:A:528:TYR:CE2	2.55	0.41
1:A:716:GLY:C	1:A:722:HIS:ND1	2.78	0.41
1:B:49:ASP:OD1	1:B:362:ARG:HD3	2.21	0.41
1:B:446:GLU:HA	1:B:447:PRO:HD3	1.95	0.41
1:B:665:CYS:O	1:B:669:PRO:HD3	2.21	0.41
1:A:101:LEU:CD2	1:A:341:TYR:CG	3.04	0.41
1:A:795:LEU:HD11	1:A:805:ASN:HA	2.03	0.41
1:B:757:VAL:HG11	1:B:820:VAL:HG21	2.02	0.41
1:A:380:ILE:HB	1:A:383:ARG:HD3	2.03	0.41
1:A:382:GLU:HG2	1:A:383:ARG:N	2.36	0.41
1:A:574:ASP:HB3	1:A:580[A]:ILE:HD11	2.03	0.40
1:A:672:ILE:HG12	1:A:687:TYR:HB2	2.03	0.40
1:B:175:VAL:HG21	1:B:210:PHE:CE2	2.56	0.40
1:B:193:GLU:CG	1:B:194:ARG:H	2.34	0.40
1:B:574:ASP:OD2	1:B:578:ASN:HB2	2.21	0.40
1:A:322:TYR:HD2	1:A:772:ARG:HD3	1.86	0.40
1:B:593:GLU:N	1:B:594:PRO:CD	2.84	0.40
1:B:6:PHE:CD1	1:B:333:ARG:HD3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	823/822 (100%)	786 (96%)	37 (4%)	0	100 100
1	B	825/822 (100%)	786 (95%)	38 (5%)	1 (0%)	48 46

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1648/1644 (100%)	1572 (95%)	75 (5%)	1 (0%)	48 46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	85	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	667/675 (99%)	664 (100%)	3 (0%)	84 89
1	B	674/675 (100%)	673 (100%)	1 (0%)	88 92
All	All	1341/1350 (99%)	1337 (100%)	4 (0%)	86 91

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	193	GLU
1	A	312	LYS
1	A	543	ASP
1	B	636	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	ASN
1	A	126	ASN
1	A	181	HIS
1	A	404	GLN
1	A	406	GLN
1	A	668	ASN
1	B	5	HIS
1	B	158	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	404	GLN
1	B	511	GLN
1	B	531	GLN
1	B	668	ASN
1	B	722	HIS
1	B	760	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BGC	C	1	2	12,12,12	1.11	2 (16%)	17,17,17	1.26	1 (5%)
2	BGC	C	2	2	11,11,12	0.97	1 (9%)	15,15,17	1.49	2 (13%)
2	BGC	D	1	2	12,12,12	0.95	0	17,17,17	1.04	1 (5%)
2	BGC	D	2	2	11,11,12	1.06	1 (9%)	15,15,17	1.81	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	1/2/22/22	0/1/1/1
2	BGC	C	2	2	-	2/2/19/22	0/1/1/1
2	BGC	D	1	2	-	0/2/22/22	0/1/1/1
2	BGC	D	2	2	-	1/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	BGC	O5-C5	-2.28	1.38	1.44
2	D	2	BGC	O5-C5	-2.18	1.39	1.43
2	C	1	BGC	O5-C1	-2.13	1.37	1.42
2	C	2	BGC	O5-C5	-2.00	1.39	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	BGC	O4-C4-C5	4.14	119.51	109.32
2	D	2	BGC	C1-C2-C3	3.33	114.50	109.64
2	C	2	BGC	C1-C2-C3	3.29	114.43	109.64
2	C	2	BGC	O4-C4-C5	2.91	116.49	109.32
2	C	1	BGC	O5-C1-C2	-2.86	105.26	110.30
2	D	1	BGC	O5-C5-C4	2.43	114.08	109.70
2	D	2	BGC	C3-C4-C5	-2.12	106.39	110.23

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	BGC	O5-C5-C6-O6
2	C	2	BGC	C4-C5-C6-O6
2	C	1	BGC	C4-C5-C6-O6
2	D	2	BGC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	822/822 (100%)	0.12	43 (5%)	33 31	3, 11, 33, 170	3 (0%)
1	B	822/822 (100%)	0.10	47 (5%)	29 28	3, 10, 29, 152	6 (0%)
All	All	1644/1644 (100%)	0.11	90 (5%)	30 29	3, 10, 32, 170	9 (0%)

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	512	ALA	13.8
1	B	612	THR	12.4
1	A	512	ALA	11.6
1	A	612	THR	10.9
1	B	610	GLN	10.7
1	B	274	ASP	9.2
1	B	613	ASP	8.7
1	B	608	GLY	8.2
1	A	607	GLU	8.2
1	B	607	GLU	8.0
1	A	557[A]	ASP	7.9
1	A	613	ASP	7.8
1	A	382	GLU	7.4
1	B	803	GLU	7.1
1	B	382	GLU	7.0
1	A	610	GLN	7.0
1	A	139	ASP	6.8
1	B	281	ASP	6.8
1	B	1	MET	6.4
1	A	52	MET	6.3
1	B	139	ASP	6.1
1	A	511	GLN	6.1
1	A	234	GLU	6.0
1	A	611	ASP	6.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	5.9
1	A	543	ASP	5.8
1	A	274	ASP	5.7
1	B	530	GLU	5.6
1	B	611	ASP	5.5
1	B	383	ARG	5.4
1	A	281	ASP	5.4
1	A	11	ARG	5.3
1	B	554	GLU	5.2
1	A	415	ASP	5.2
1	A	32	GLN	5.0
1	A	547	ARG	5.0
1	A	114	ARG	5.0
1	B	234	GLU	4.9
1	B	764	ASP	4.7
1	B	609	PRO	4.6
1	A	513	GLY	4.3
1	B	178	GLU	4.3
1	B	543	ASP	4.2
1	B	386	GLU	4.0
1	B	698	ASP	4.0
1	B	362	ARG	4.0
1	B	32	GLN	4.0
1	B	116	GLU	4.0
1	A	178	GLU	3.9
1	A	116	GLU	3.8
1	B	317	ASP	3.8
1	B	275	GLU	3.8
1	B	453	GLU	3.7
1	B	513	GLY	3.7
1	B	316	THR	3.7
1	B	751	GLU	3.7
1	B	286[A]	VAL	3.5
1	A	549	ARG	3.5
1	A	283	LYS	3.5
1	B	415	ASP	3.5
1	A	554	GLU	3.4
1	A	317	ASP	3.4
1	A	158	GLN	3.4
1	B	547	ARG	3.3
1	A	386	GLU	3.2
1	A	453	GLU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	327	ASN	3.2
1	A	362	ARG	3.2
1	A	67	ALA	3.1
1	B	11	ARG	3.0
1	A	803	GLU	2.9
1	A	383	ARG	2.9
1	B	509	GLU	2.8
1	A	48	ARG	2.7
1	B	289	GLU	2.7
1	B	511	GLN	2.7
1	A	580[A]	ILE	2.7
1	B	48	ARG	2.6
1	A	597	PHE	2.6
1	A	608	GLY	2.6
1	A	143	SER	2.6
1	B	360	ILE	2.4
1	B	93	ASP	2.4
1	A	360	ILE	2.3
1	B	260	SER	2.2
1	A	259	GLU	2.2
1	B	2	ARG	2.2
1	A	140	SER	2.2
1	B	753	ASP	2.1
1	B	597	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	C	1	12/12	0.92	0.08	10,19,23,26	0
2	BGC	C	2	11/12	0.94	0.08	5,12,19,25	0
2	BGC	D	1	12/12	0.95	0.07	6,13,21,26	0
2	BGC	D	2	11/12	0.95	0.07	5,9,15,18	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.