



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 07:21 AM UTC

PDB ID : 6S64 / pdb_00006s64
Title : Crystal structure of hTEAD2 in complex with a trisubstituted pyrazole inhibitor
Authors : Sturbaut, M.; Allemand, F.; Guichou, J.F.
Deposited on : 2019-07-02
Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	
1	B	240	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3427 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Transcriptional enhancer factor TEF-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	195	Total	C	N	O	S	0	1	0
			1613	1037	281	287	8			
1	B	210	Total	C	N	O	S	0	2	0
			1730	1106	302	313	9			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	216	MET	-	initiating methionine	UNP Q15562
A	448	VAL	-	expression tag	UNP Q15562
A	449	GLU	-	expression tag	UNP Q15562
A	450	HIS	-	expression tag	UNP Q15562
A	451	HIS	-	expression tag	UNP Q15562
A	452	HIS	-	expression tag	UNP Q15562
A	453	HIS	-	expression tag	UNP Q15562
A	454	HIS	-	expression tag	UNP Q15562
A	455	HIS	-	expression tag	UNP Q15562
B	216	MET	-	initiating methionine	UNP Q15562
B	448	VAL	-	expression tag	UNP Q15562
B	449	GLU	-	expression tag	UNP Q15562
B	450	HIS	-	expression tag	UNP Q15562
B	451	HIS	-	expression tag	UNP Q15562
B	452	HIS	-	expression tag	UNP Q15562
B	453	HIS	-	expression tag	UNP Q15562
B	454	HIS	-	expression tag	UNP Q15562
B	455	HIS	-	expression tag	UNP Q15562

- Molecule 2 is MYRISTIC ACID (CCD ID: MYR) (formula: C₁₄H₂₈O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			16	14	2		

- Molecule 3 is PALMITIC ACID (CCD ID: PLM) (formula: $C_{16}H_{32}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			17	16	1		

- Molecule 4 is 3-[3-(3,4-dichlorophenyl)-4-[(phenylmethyl)carbamoyl]pyrazol-1-yl]propanoic acid (CCD ID: KXE) (formula: $C_{20}H_{17}Cl_2N_3O_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	Cl	N	O	0	0
			28	20	2	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	11	Total	O	0	0
			11	11		
5	B	12	Total	O	0	0
			12	12		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional enhancer factor TEF-4



- Molecule 1: Transcriptional enhancer factor TEF-4



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.69Å 61.66Å 80.24Å 90.00° 117.36° 90.00°	Depositor
Resolution (Å)	71.26 – 2.22 71.26 – 2.22	Depositor EDS
% Data completeness (in resolution range)	98.4 (71.26-2.22) 98.4 (71.26-2.22)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.203 , 0.259 0.208 , 0.260	Depositor DCC
R_{free} test set	1296 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	55.0	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3427	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLM, MYR, KXE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	0/1654	1.06	5/2230 (0.2%)
1	B	0.94	0/1777	1.01	2/2401 (0.1%)
All	All	0.95	0/3431	1.03	7/4631 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	264	PRO	CA-C-N	10.99	133.57	119.84
1	A	264	PRO	C-N-CA	10.99	133.57	119.84
1	A	293	PRO	CA-C-N	5.99	126.15	119.32
1	A	293	PRO	C-N-CA	5.99	126.15	119.32
1	A	236	PHE	N-CA-C	5.24	116.24	108.60
1	B	411	VAL	CB-CA-C	5.17	117.88	110.33
1	B	400	ASN	N-CA-C	5.03	117.50	111.71

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	264	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1613	0	1593	14	0
1	B	1730	0	1693	7	0
2	A	16	0	27	0	0
3	B	17	0	31	0	0
4	B	28	0	0	1	0
5	A	11	0	0	0	0
5	B	12	0	0	0	0
All	All	3427	0	3344	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:382:TYR:HB3	4:B:502:KXE:CL1	2.17	0.81
1:A:266:LEU:HD23	1:A:443:ARG:HG3	1.85	0.58
1:A:237:VAL:CG1	1:A:327:TYR:CE2	2.92	0.53
1:A:237:VAL:HG11	1:A:327:TYR:CE2	2.44	0.52
1:A:266:LEU:HD23	1:A:443:ARG:CG	2.40	0.51
1:A:264:PRO:HB2	1:A:265:PRO:CD	2.41	0.50
1:A:396:ARG:HG2	1:A:430:VAL:CG2	2.41	0.50
1:A:389:LYS:O	1:A:392:GLN:HG2	2.11	0.50
1:A:238:GLU:OE1	1:A:376:ARG:NH2	2.45	0.50
1:B:234:SER:HB3	1:B:253:HIS:HD2	1.79	0.48
1:A:237:VAL:HG11	1:A:327:TYR:CZ	2.49	0.48
1:B:366:GLU:OE1	1:B:373:ARG:NH2	2.49	0.45
1:A:237:VAL:HG22	1:A:251:PHE:CZ	2.52	0.44
1:B:239:PRO:HG3	1:B:247:GLN:HE21	1.83	0.44
1:B:220:ALA:HB3	1:B:443:ARG:HD3	2.00	0.43
1:B:332:GLN:HG2	1:B:373:ARG:HG2	2.01	0.43
1:A:374:LEU:N	1:A:374:LEU:HD12	2.34	0.42
1:A:390:LEU:HD21	1:A:403:LEU:HD21	2.01	0.42
1:A:237:VAL:CG1	1:A:327:TYR:CD2	3.03	0.42
1:B:278:PHE:CE1	1:B:407:THR:HB	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:VAL:HG13	1:A:444:LEU:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/240 (78%)	180 (96%)	6 (3%)	2 (1%)	11	9
1	B	206/240 (86%)	193 (94%)	13 (6%)	0	100	100
All	All	394/480 (82%)	373 (95%)	19 (5%)	2 (0%)	24	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	265	PRO
1	A	271	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/210 (85%)	169 (95%)	9 (5%)	21	26
1	B	190/210 (90%)	174 (92%)	16 (8%)	10	10
All	All	368/420 (88%)	343 (93%)	25 (7%)	15	16

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	237	VAL
1	A	255	SER
1	A	271	VAL
1	A	288	LEU
1	A	410	GLN
1	A	418	GLN
1	A	420	LEU
1	A	430	VAL
1	A	433	SER
1	B	219	GLN
1	B	221	ARG
1	B	232	GLU
1	B	248	ARG
1	B	291	ARG
1	B	329	VAL
1	B	344	SER
1	B	354	VAL
1	B	355	VAL
1	B	380	CYS
1	B	392[A]	GLN
1	B	392[B]	GLN
1	B	393	LEU
1	B	411	VAL
1	B	434	GLU
1	B	446	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	307	ASN
1	A	392	GLN
1	A	410	GLN
1	B	247	GLN
1	B	388	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	MYR	A	501	-	15,15,15	1.01	2 (13%)	15,15,15	1.07	2 (13%)
3	PLM	B	501	1	15,16,17	0.33	0	14,15,17	0.56	0
4	KXE	B	502	-	30,30,30	0.83	2 (6%)	40,41,41	0.83	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MYR	A	501	-	-	7/13/13/13	-
3	PLM	B	501	1	-	8/14/14/15	-
4	KXE	B	502	-	-	9/18/18/18	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	502	KXE	O1-C12	3.44	1.33	1.22
2	A	501	MYR	O2-C1	-2.76	1.21	1.30
2	A	501	MYR	O1-C1	2.55	1.30	1.22
4	B	502	KXE	O2-C12	-2.47	1.22	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	502	KXE	O1-C12-C11	-3.14	113.15	123.09
2	A	501	MYR	O1-C1-C2	-2.85	114.05	123.09
2	A	501	MYR	O2-C1-C2	2.74	122.64	114.00
4	B	502	KXE	O2-C12-C11	2.70	122.54	114.00

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	501	PLM	O2-C1-C2-C3
4	B	502	KXE	N1-C10-C11-C12
4	B	502	KXE	O-C-N-C1
3	B	501	PLM	CA-CB-CC-CD
2	A	501	MYR	C4-C5-C6-C7
3	B	501	PLM	C3-C4-C5-C6
2	A	501	MYR	C2-C3-C4-C5
2	A	501	MYR	C10-C11-C12-C13
3	B	501	PLM	C2-C3-C4-C5
2	A	501	MYR	C5-C6-C7-C8
2	A	501	MYR	C11-C10-C9-C8
3	B	501	PLM	C9-CA-CB-CC
4	B	502	KXE	O-C-C8-C13
4	B	502	KXE	N-C-C8-C13
4	B	502	KXE	C8-C-N-C1
3	B	501	PLM	C5-C6-C7-C8
3	B	501	PLM	CC-CD-CE-CF
4	B	502	KXE	C11-C10-N1-N2
2	A	501	MYR	C7-C8-C9-C10
4	B	502	KXE	C10-C11-C12-O1
4	B	502	KXE	C10-C11-C12-O2
4	B	502	KXE	N-C-C8-C9
2	A	501	MYR	C6-C7-C8-C9
3	B	501	PLM	CB-CC-CD-CE


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	502	KXE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	195/240 (81%)	0.42	8 (4%) 41 38	38, 64, 106, 128	1 (0%)
1	B	210/240 (87%)	0.24	7 (3%) 49 46	35, 61, 98, 120	2 (0%)
All	All	405/480 (84%)	0.33	15 (3%) 45 42	35, 63, 101, 128	3 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	323	SER	5.6
1	A	239	PRO	4.9
1	A	264	PRO	3.7
1	A	265	PRO	3.1
1	B	263	ALA	3.0
1	A	222	GLY	2.9
1	A	324	GLY	2.7
1	B	447	ASP	2.7
1	B	221	ARG	2.3
1	B	365	LEU	2.3
1	A	226	ALA	2.3
1	A	326	PHE	2.2
1	B	282	LYS	2.1
1	B	283	GLY	2.1
1	A	283	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	KXE	B	502	28/28	0.86	0.21	80,87,94,101	28
3	PLM	B	501	17/18	0.91	0.19	54,67,82,109	0
2	MYR	A	501	16/16	0.96	0.14	59,69,94,100	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.