



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:42 AM UTC

PDB ID : 6S6F / pdb_00006s6f
Title : Crystal structure of 2-methylcitrate synthase (PrpC) from *Pseudomonas aeruginosa* in apo form.
Authors : Wijaya, A.J.; Brear, P.; Dolan, S.K.; Welch, M.
Deposited on : 2019-07-03
Resolution : 1.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1003 (1.54-1.54)
Clashscore	190562	1025 (1.54-1.54)
Ramachandran outliers	187476	1007 (1.54-1.54)
Sidechain outliers	187428	1007 (1.54-1.54)
RSRZ outliers	180081	1002 (1.54-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	5% 86% 10% .
1	B	375	7% 90% 7% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	401	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6103 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Citrate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	1	0
			2844	1807	494	530	13			
1	B	370	Total	C	N	O	S	0	4	0
			2931	1860	510	548	13			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	1
			12	6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	124	Total	O	0	1
			125	125		
4	B	166	Total	O	0	0
			166	166		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Citrate synthase



• Molecule 1: Citrate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.01Å 83.19Å 82.71Å 90.00° 101.93° 90.00°	Depositor
Resolution (Å)	57.99 – 1.53 57.99 – 1.53	Depositor EDS
% Data completeness (in resolution range)	98.9 (57.99-1.53) 99.1 (57.99-1.53)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 1.53Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.196 , 0.213 0.198 , 0.199	Depositor DCC
R_{free} test set	5085 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.364	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6103	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.07	10/2908 (0.3%)	0.94	2/3935 (0.1%)
1	B	0.95	14/2995 (0.5%)	0.90	1/4052 (0.0%)
All	All	1.01	24/5903 (0.4%)	0.92	3/7987 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	371	PRO	N-CD	-8.54	1.35	1.47
1	A	331	PRO	C-O	-6.59	1.15	1.24
1	A	132	ALA	C-O	-6.35	1.16	1.24
1	B	71	LYS	C-O	-6.31	1.16	1.24
1	B	331	PRO	C-O	-6.16	1.16	1.24
1	A	202	THR	C-O	-5.99	1.16	1.24
1	A	61	PRO	C-O	-5.88	1.16	1.23
1	B	70	LEU	C-O	-5.86	1.17	1.24
1	B	333	PHE	C-O	-5.53	1.17	1.24
1	A	344	HIS	CE1-NE2	-5.40	1.27	1.32
1	B	309	PRO	C-O	-5.38	1.18	1.23
1	A	122	VAL	C-O	-5.37	1.17	1.24
1	B	220	PRO	C-O	-5.34	1.17	1.24
1	A	131	PRO	C-O	-5.31	1.17	1.24
1	B	357	PRO	C-O	-5.28	1.17	1.23
1	B	344	HIS	CE1-NE2	-5.24	1.27	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	185	ALA	C-O	-5.17	1.18	1.24
1	B	226	ASN	C-O	-5.17	1.17	1.24
1	B	218	ARG	C-O	-5.16	1.17	1.24
1	A	130	PHE	C-O	-5.11	1.19	1.24
1	B	194	PHE	C-O	-5.10	1.18	1.24
1	B	311	ALA	C-O	-5.07	1.17	1.24
1	B	188	GLU	C-O	-5.06	1.19	1.24
1	A	187	HIS	C-O	-5.01	1.18	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	190	ASN	CA-CB-CG	-5.64	106.96	112.60
1	A	329	PHE	CA-CB-CG	5.52	119.32	113.80
1	A	352	ASN	N-CA-C	5.32	118.16	109.59

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	24[B]	THR	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2844	0	2808	16	0
1	B	2931	0	2899	18	0
2	A	12	0	16	4	0
2	B	24	0	32	3	0
3	B	1	0	0	1	0
4	A	125	0	0	1	0
4	B	166	0	0	0	0
All	All	6103	0	5755	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:13:ARG:HG2	1:B:13:ARG:NH1	2.09	0.67
1:A:333:PHE:CD2	2:A:401:GOL:H2	2.32	0.64
1:B:371:PRO:HG2	1:B:374:GLN:HB3	1.80	0.62
1:A:16:VAL:HG23	1:A:16:VAL:O	2.03	0.59
1:B:13:ARG:HG2	1:B:13:ARG:HH11	1.66	0.58
1:A:226:ASN:ND2	2:A:401:GOL:H12	2.20	0.56
1:A:372:LEU:HD21	1:B:63[A]:LYS:HD2	1.90	0.54
1:B:371:PRO:HG2	1:B:374:GLN:CB	2.38	0.54
1:A:194:PHE:CZ	1:B:357:PRO:HG2	2.44	0.53
1:A:228:ALA:HB1	1:A:257:MET:HG3	1.92	0.51
1:B:224:GLY:H	2:B:402[A]:GOL:H31	1.77	0.50
1:B:247:LEU:O	1:B:251:GLU:HG2	2.12	0.49
1:A:139[A]:ARG:NH2	1:A:145:GLN:OE1	2.45	0.49
4:A:573:HOH:O	1:B:24[B]:THR:HG21	2.13	0.48
1:A:252:ARG:NH2	1:A:254:ASP:OD2	2.46	0.48
1:B:56:LEU:HD22	1:B:346:PHE:CE2	2.51	0.46
1:B:63[B]:LYS:HE2	1:B:67:ASP:OD2	2.17	0.45
1:B:56:LEU:HD21	1:B:342:THR:OG1	2.17	0.45
1:A:312:ASP:OD1	2:A:401:GOL:H11	2.17	0.45
1:A:176:VAL:HG11	1:A:281:LEU:HD11	1.99	0.44
1:B:246:LEU:HD23	1:B:300:THR:HB	1.98	0.44
1:A:86:VAL:HG11	1:A:106:VAL:HG21	2.00	0.43
1:A:217:LEU:O	1:A:223:GLY:HA3	2.19	0.42
1:A:17:ALA:HB2	1:B:357:PRO:HB3	2.02	0.41
1:B:320:HIS:CE1	2:B:403:GOL:H11	2.55	0.41
1:B:320:HIS:NE2	2:B:403:GOL:H11	2.34	0.41
1:A:139[B]:ARG:NH2	1:A:164:LEU:O	2.47	0.41
1:B:121:ASP:O	3:B:404:CL:CL	2.76	0.41
1:A:226:ASN:CG	2:A:401:GOL:H12	2.45	0.41
1:A:139[B]:ARG:CZ	1:A:143:GLU:OE1	2.68	0.41
1:B:86:VAL:HG11	1:B:106:VAL:HG21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/375 (96%)	354 (99%)	5 (1%)	0	100	100
1	B	372/375 (99%)	365 (98%)	7 (2%)	0	100	100
All	All	731/750 (98%)	719 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/306 (97%)	295 (99%)	3 (1%)	68	42
1	B	307/306 (100%)	306 (100%)	1 (0%)	86	74
All	All	605/612 (99%)	601 (99%)	4 (1%)	76	55

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	188	GLU
1	A	315	HIS
1	A	356	ARG
1	B	188	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN
1	A	76	GLN
1	A	150	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	403	-	5,5,5	0.11	0	5,5,5	0.66	0
2	GOL	B	401	-	5,5,5	0.31	0	5,5,5	0.70	0
2	GOL	A	401	-	5,5,5	0.31	0	5,5,5	0.48	0
2	GOL	B	402[B]	-	5,5,5	0.16	0	5,5,5	0.47	0
2	GOL	B	402[A]	-	5,5,5	0.17	0	5,5,5	0.61	0
2	GOL	A	402	-	5,5,5	0.19	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	403	-	-	2/4/4/4	-
2	GOL	B	401	-	-	2/4/4/4	-
2	GOL	A	401	-	-	2/4/4/4	-
2	GOL	B	402[B]	-	-	1/4/4/4	-
2	GOL	B	402[A]	-	-	4/4/4/4	-
2	GOL	A	402	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	GOL	O1-C1-C2-C3
2	B	402[A]	GOL	C1-C2-C3-O3
2	B	403	GOL	C1-C2-C3-O3
2	A	401	GOL	O1-C1-C2-C3
2	A	402	GOL	C1-C2-C3-O3
2	B	402[A]	GOL	O1-C1-C2-C3
2	B	402[A]	GOL	O2-C2-C3-O3
2	B	403	GOL	O2-C2-C3-O3
2	B	402[B]	GOL	O2-C2-C3-O3
2	B	401	GOL	O1-C1-C2-O2
2	A	402	GOL	O2-C2-C3-O3
2	B	402[A]	GOL	O1-C1-C2-O2
2	A	401	GOL	C1-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	403	GOL	2	0
2	A	401	GOL	4	0
2	B	402[A]	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	360/375 (96%)	0.62	19 (5%)	32 40	14, 25, 46, 82	1 (0%)
1	B	370/375 (98%)	0.38	25 (6%)	23 29	8, 20, 46, 86	4 (1%)
All	All	730/750 (97%)	0.50	44 (6%)	27 35	8, 23, 46, 86	5 (0%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	16	VAL	8.2
1	A	17	ALA	5.9
1	B	371	PRO	4.3
1	B	374	GLN	3.9
1	B	372	LEU	3.9
1	B	30	ALA	3.8
1	B	27	GLN	3.5
1	B	8	SER	3.4
1	B	26	GLY	3.2
1	B	6	VAL	3.2
1	B	373	GLU	3.2
1	B	7	LEU	3.0
1	A	265	LYS	3.0
1	B	251	GLU	2.9
1	B	14	GLY	2.8
1	B	25	VAL	2.7
1	B	302	TRP	2.7
1	B	369	PHE	2.7
1	A	263	ILE	2.6
1	B	285	VAL	2.6
1	A	29	GLY	2.5
1	B	370	VAL	2.5
1	B	375	ARG	2.5
1	B	114	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	253	LYS	2.5
1	A	30	ALA	2.4
1	A	18	GLY	2.4
1	B	357	PRO	2.4
1	A	252	ARG	2.4
1	B	364	VAL	2.4
1	B	356	ARG	2.3
1	A	168	LYS	2.3
1	B	29	GLY	2.2
1	B	250	LEU	2.2
1	A	288	LYS	2.2
1	A	254	ASP	2.2
1	A	251	GLU	2.2
1	A	262	ALA	2.1
1	A	37	TYR	2.1
1	A	283	ASP	2.1
1	A	289	VAL	2.1
1	A	45	ALA	2.1
1	A	286	GLY	2.0
1	A	257	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	403	6/6	0.40	0.26	44,46,50,50	0
2	GOL	B	402[B]	6/6	0.65	0.17	25,29,32,32	6
2	GOL	B	402[A]	6/6	0.65	0.17	20,28,31,33	6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	B	401	6/6	0.74	0.20	42,44,49,51	0
2	GOL	A	401	6/6	0.81	0.13	30,32,33,35	0
2	GOL	A	402	6/6	0.86	0.14	39,49,51,55	0
3	CL	B	404	1/1	0.88	0.20	59,59,59,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.