



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 07:38 PM UTC

PDB ID : 1S7Y / pdb_00001s7y
Title : Crystal structure of the GluR6 ligand binding core in complex with glutamate at 1.75 Å resolution orthorhombic form
Authors : Mayer, M.L.
Deposited on : 2004-01-30
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	
1	B	259	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutamate receptor, ionotropic kainate 2 precursor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	28	0	0
			2012	1286	334	381	11			
1	B	251	Total	C	N	O	S	11	0	0
			2012	1286	334	381	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	SEE REMARK 999	UNP P42260
A	118	GLY	-	SEE REMARK 999	UNP P42260
A	119	THR	-	SEE REMARK 999	UNP P42260
B	1	GLY	-	SEE REMARK 999	UNP P42260
B	118	GLY	-	SEE REMARK 999	UNP P42260
B	119	THR	-	SEE REMARK 999	UNP P42260

- Molecule 2 is GLUTAMIC ACID (CCD ID: GLU) (formula: C₅H₉NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			10	5	1	4		
2	A	1	Total	C	N	O	0	0
			10	5	1	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	211	Total	O	0	0
			211	211		
3	B	216	Total	O	0	0
			216	216		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate receptor, ionotropic kainate 2 precursor



- Molecule 1: Glutamate receptor, ionotropic kainate 2 precursor



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.10Å 91.19Å 105.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.51 – 1.75 24.51 – 1.75	Depositor EDS
% Data completeness (in resolution range)	90.4 (24.51-1.75) 99.0 (24.51-1.75)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.78 (at 1.75Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.220 , 0.243 0.227 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	14.7	Xtriage
Anisotropy	0.551	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4471	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.44 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.9261e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/2050	0.86	4/2763 (0.1%)
1	B	0.47	0/2050	0.87	2/2763 (0.1%)
All	All	0.46	0/4100	0.87	6/5526 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	57	GLU	N-CA-C	7.72	119.70	111.28
1	B	57	GLU	N-CA-C	7.08	119.85	111.71
1	A	104	LYS	N-CA-C	-5.49	102.29	110.20
1	B	198	VAL	N-CA-C	5.48	115.68	110.42
1	A	99	VAL	N-CA-C	5.46	118.88	113.53
1	A	103	SER	N-CA-C	-5.32	103.00	110.35

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2012	0	2026	23	0
1	B	2012	0	2026	28	0
2	A	20	0	10	0	0
3	A	211	0	0	1	0
3	B	216	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4471	0	4062	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:95:VAL:HG11	1:B:149:LYS:HE3	1.59	0.83
1:B:88:ALA:HB1	1:B:89:PRO:HD2	1.68	0.75
1:A:172:LYS:HA	1:A:172:LYS:HE2	1.69	0.73
1:A:88:ALA:HB1	1:A:89:PRO:HD2	1.73	0.71
1:B:136:GLY:HA2	1:B:162:MET:HE2	1.71	0.70
1:B:54:ARG:NH2	1:B:57:GLU:HG3	2.08	0.67
1:A:133:ILE:HG12	1:A:185:ASP:OD1	1.97	0.65
1:A:54:ARG:NH2	1:A:57:GLU:HG3	2.13	0.63
1:B:201:ARG:HG3	1:B:201:ARG:HH11	1.65	0.61
1:B:202:ASN:HB2	3:B:417:HOH:O	2.01	0.61
1:A:236:LEU:HD12	3:B:405:HOH:O	2.02	0.59
1:B:188:PHE:HD2	1:B:190:MET:HE3	1.67	0.59
1:A:236:LEU:HD23	1:A:236:LEU:O	2.02	0.58
1:A:122:ASP:OD1	1:B:239:GLN:NE2	2.35	0.57
1:B:133:ILE:HG12	1:B:185:ASP:OD1	2.04	0.57
1:B:188:PHE:CD2	1:B:190:MET:HE3	2.41	0.56
1:B:98:LYS:HA	1:B:98:LYS:HE3	1.88	0.55
1:A:122:ASP:CG	1:B:239:GLN:HE22	2.15	0.54
1:B:136:GLY:CA	1:B:162:MET:HE2	2.38	0.54
1:A:188:PHE:HD2	1:A:190:MET:HE3	1.73	0.54
1:A:93:THR:O	1:A:97:GLU:HG3	2.07	0.54
1:A:228:ARG:HD2	1:A:228:ARG:C	2.32	0.54
1:A:201:ARG:HH11	1:A:201:ARG:HG3	1.71	0.54
1:B:180:ARG:HG2	1:B:186:TYR:CD2	2.42	0.54
1:B:93:THR:O	1:B:97:GLU:HG3	2.09	0.51
1:B:201:ARG:HG3	1:B:201:ARG:NH1	2.24	0.51
1:A:188:PHE:CD2	1:A:190:MET:HE3	2.48	0.48
1:B:228:ARG:HD2	1:B:228:ARG:C	2.38	0.48
1:A:201:ARG:HG3	1:A:201:ARG:NH1	2.31	0.46
1:A:202:ASN:HB2	3:A:718:HOH:O	2.15	0.46
1:A:211:LEU:HD22	1:B:211:LEU:HD22	1.98	0.46
1:B:113:ILE:HB	1:B:208:ILE:HB	1.98	0.46
1:B:109:LEU:CD1	1:B:191:GLU:HB3	2.48	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:154:THR:O	1:B:157:LYS:HG2	2.18	0.44
1:A:240:GLU:HG3	1:B:119:THR:O	2.17	0.44
1:A:236:LEU:HD23	1:A:236:LEU:C	2.42	0.44
1:B:88:ALA:HB1	1:B:89:PRO:CD	2.44	0.43
1:B:232:THR:O	1:B:236:LEU:HG	2.18	0.43
1:A:135:TYR:HA	1:A:187:ALA:O	2.19	0.42
1:A:166:ARG:HG3	1:A:170:LEU:HD12	2.01	0.42
1:A:180:ARG:HG2	1:A:186:TYR:CD2	2.55	0.42
1:B:66:ASP:OD1	1:B:148:LYS:HE2	2.19	0.42
1:B:117:LYS:HD3	1:B:203:CYS:O	2.20	0.42
1:A:133:ILE:HG23	1:A:185:ASP:OD1	2.20	0.41
1:B:135:TYR:HA	1:B:187:ALA:O	2.20	0.41
1:A:119:THR:O	1:B:240:GLU:HG3	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/259 (96%)	241 (97%)	7 (3%)	1 (0%)	30	15
1	B	249/259 (96%)	240 (96%)	9 (4%)	0	100	100
All	All	498/518 (96%)	481 (97%)	16 (3%)	1 (0%)	43	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/225 (98%)	217 (99%)	3 (1%)	59	44
1	B	220/225 (98%)	217 (99%)	3 (1%)	59	44
All	All	440/450 (98%)	434 (99%)	6 (1%)	59	44

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	PRO
1	A	18	LEU
1	A	236	LEU
1	B	15	PRO
1	B	98	LYS
1	B	253	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	ASN
1	A	207	GLN
1	B	29	ASN
1	B	239	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLU	A	552	-	8,9,9	1.04	0	8,11,11	0.82	0
2	GLU	A	551	-	8,9,9	0.99	0	8,11,11	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLU	A	552	-	-	2/9/9/9	-
2	GLU	A	551	-	-	2/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	551	GLU	O-C-CA-CB
2	A	552	GLU	O-C-CA-CB
2	A	552	GLU	OXT-C-CA-CB
2	A	551	GLU	OXT-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/259 (96%)	0.63	32 (12%) 8 8	7, 15, 33, 45	8 (3%)
1	B	251/259 (96%)	0.51	27 (10%) 11 12	7, 15, 33, 46	2 (0%)
All	All	502/518 (96%)	0.57	59 (11%) 9 10	7, 15, 33, 46	10 (1%)

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	253	ARG	7.9
1	A	67	VAL	5.2
1	A	203	CYS	4.8
1	A	3	ASN	4.8
1	A	66	ASP	4.5
1	A	23	ASP	4.5
1	B	183	THR	4.3
1	B	24	LYS	4.1
1	A	122	ASP	4.0
1	A	183	THR	3.8
1	B	3	ASN	3.6
1	A	133	ILE	3.4
1	A	185	ASP	3.4
1	B	203	CYS	3.4
1	A	202	ASN	3.3
1	B	23	ASP	3.3
1	A	152	ILE	3.2
1	B	253	ARG	3.2
1	A	69	GLY	3.2
1	B	202	ASN	3.2
1	A	182	LEU	3.2
1	A	24	LYS	3.1
1	A	129	LYS	3.1
1	A	200	GLN	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	157	LYS	3.1
1	B	185	ASP	3.1
1	B	204	ASN	3.0
1	A	204	ASN	3.0
1	B	4	ARG	2.9
1	A	25	PRO	2.9
1	A	236	LEU	2.7
1	A	181	VAL	2.7
1	A	68	ASN	2.6
1	B	125	ASP	2.5
1	A	70	GLN	2.5
1	B	184	SER	2.5
1	A	156	ASP	2.5
1	B	182	LEU	2.4
1	B	153	SER	2.4
1	A	175	GLU	2.4
1	B	167	GLN	2.3
1	A	65	ASP	2.3
1	A	116	ARG	2.3
1	B	25	PRO	2.3
1	A	127	LEU	2.3
1	A	117	LYS	2.3
1	B	181	VAL	2.3
1	B	152	ILE	2.2
1	A	153	SER	2.2
1	B	30	ASP	2.2
1	B	236	LEU	2.2
1	B	66	ASP	2.1
1	A	5	SER	2.1
1	B	200	GLN	2.1
1	B	239	GLN	2.1
1	B	129	LYS	2.1
1	B	122	ASP	2.0
1	B	128	ALA	2.0
1	A	151	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLU	A	551	10/10	0.96	0.05	6,7,9,10	0
2	GLU	A	552	10/10	0.96	0.06	7,8,8,8	0

6.5 Other polymers [i](#)

There are no such residues in this entry.