



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 03:45 AM UTC

PDB ID : 3S71 / pdb_00003s71
Title : The origin of the hydrophobic effect in the molecular recognition of arylsulfonamides by carbonic anhydrase
Authors : Snyder, P.W.; Heroux, A.; Whitesides, G.W.
Deposited on : 2011-05-26
Resolution : 1.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1693 (1.28-1.24)
Clashscore	190562	1730 (1.28-1.24)
Ramachandran outliers	187476	1695 (1.28-1.24)
Sidechain outliers	187428	1694 (1.28-1.24)
RSRZ outliers	180081	1693 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	258	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2549 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Carbonic anhydrase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	258	Total	C	N	O	S	0	15	0
			2155	1381	368	404	2			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 1-benzofuran-2-sulfonamide (CCD ID: EVD) (formula: C₈H₇NO₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			13	8	1	3	1		
3	B	1	Total	C	N	O	S	0	0
			13	8	1	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	367	Total 367	O 367	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carbonic anhydrase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.37Å 41.40Å 72.47Å 90.00° 104.69° 90.00°	Depositor
Resolution (Å)	35.64 – 1.25 35.64 – 1.25	Depositor EDS
% Data completeness (in resolution range)	90.9 (35.64-1.25) 90.8 (35.64-1.25)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 1.25Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.157 , 0.181 0.154 , 0.180	Depositor DCC
R_{free} test set	3088 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	10.1	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2549	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EVD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	1.64	20/2241 (0.9%)	1.37	12/3042 (0.4%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	161	VAL	CA-CB	11.78	1.69	1.54
1	B	176	PHE	CA-CB	8.47	1.59	1.53
1	B	253	ASN	C-O	6.64	1.36	1.23
1	B	34	ASP	C-N	-6.47	1.25	1.33
1	B	37	THR	CA-C	6.34	1.61	1.52
1	B	47	LEU	CA-C	6.04	1.60	1.52
1	B	173	SER	C-O	5.96	1.31	1.23
1	B	34	ASP	N-CA	5.87	1.53	1.46
1	B	90	LEU	CA-C	-5.83	1.45	1.52
1	B	214	GLU	CD-OE1	-5.66	1.14	1.25
1	B	96	HIS	ND1-CE1	5.56	1.38	1.32
1	B	73	SER	N-CA	5.37	1.53	1.46
1	B	47	LEU	CB-CG	-5.33	1.42	1.53
1	B	171	GLY	C-O	5.32	1.31	1.24
1	B	89	ARG	CA-C	-5.22	1.46	1.52
1	B	8	GLY	CA-C	-5.19	1.45	1.52
1	B	254	ARG	CA-C	-5.12	1.46	1.52
1	B	132	GLY	N-CA	-5.11	1.38	1.45
1	B	22[A]	ILE	CA-CB	5.09	1.63	1.54
1	B	22[B]	ILE	CA-CB	5.09	1.63	1.54

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	25	GLY	O-C-N	9.19	129.88	122.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	253	ASN	CA-C-O	-6.63	111.79	122.20
1	B	47	LEU	CB-CG-CD2	-6.10	92.41	110.70
1	B	125	THR	O-C-N	5.90	130.28	122.43
1	B	14	GLU	CB-CG-CD	5.67	122.24	112.60
1	B	46	PRO	O-C-N	5.57	129.91	123.01
1	B	46	PRO	CA-C-O	-5.42	115.27	121.56
1	B	17	HIS	O-C-N	5.25	128.36	122.22
1	B	127	LYS	N-CA-C	-5.24	106.72	113.01
1	B	46	PRO	CA-C-N	-5.08	112.97	120.94
1	B	46	PRO	C-N-CA	-5.08	112.97	120.94
1	B	74	GLN	CB-CA-C	-5.03	99.67	111.09

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2155	0	2093	37	0
2	B	1	0	0	0	0
3	B	26	0	13	0	0
4	B	367	0	0	18	0
All	All	2549	0	2106	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:HIS:HE1	4:B:344:HOH:O	1.40	1.04
1:B:3:HIS:CE1	4:B:344:HOH:O	2.18	0.91
1:B:26[A]:GLU:OE2	4:B:276:HOH:O	1.94	0.83
1:B:214:GLU:OE2	4:B:538:HOH:O	1.95	0.82
1:B:165[A]:ASP:OD1	4:B:291:HOH:O	1.96	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64[B]:HIS:HE1	4:B:323:HOH:O	1.66	0.79
1:B:3:HIS:N	4:B:460:HOH:O	2.17	0.77
1:B:257:LYS:HE2	4:B:285:HOH:O	1.87	0.75
1:B:80:LYS:CE	4:B:329:HOH:O	2.41	0.68
1:B:61:ASN:HD22	1:B:63:GLY:H	1.42	0.67
1:B:255:GLN:HG2	4:B:285:HOH:O	1.94	0.67
1:B:61:ASN:ND2	1:B:63:GLY:H	1.94	0.65
1:B:61:ASN:HD22	1:B:61:ASN:C	2.06	0.63
1:B:27[B]:ARG:NH1	4:B:264:HOH:O	2.33	0.60
1:B:163[B]:VAL:HG23	4:B:533:HOH:O	2.01	0.60
1:B:26[A]:GLU:HA	1:B:252:LYS:HD2	1.85	0.58
1:B:26[B]:GLU:OE1	1:B:252:LYS:HD3	2.05	0.56
1:B:39:LYS:HG3	4:B:610:HOH:O	2.07	0.54
1:B:17:HIS:HD2	4:B:486:HOH:O	1.90	0.54
1:B:89:ARG:HG3	1:B:125:THR:CG2	2.38	0.54
1:B:27[A]:ARG:HG3	1:B:205:GLU:HB3	1.94	0.49
1:B:27[B]:ARG:CZ	4:B:264:HOH:O	2.60	0.47
1:B:64[B]:HIS:CE1	4:B:323:HOH:O	2.50	0.47
1:B:134:ALA:O	1:B:140:GLY:HA3	2.16	0.46
1:B:47:LEU:HD21	1:B:210:ILE:HD13	1.98	0.45
1:B:163[B]:VAL:O	1:B:163[B]:VAL:CG1	2.65	0.45
1:B:163[B]:VAL:O	1:B:163[B]:VAL:HG12	2.15	0.45
1:B:61:ASN:ND2	1:B:61:ASN:C	2.75	0.44
1:B:154:LYS:HB2	1:B:217[C]:SER:O	2.19	0.42
1:B:89:ARG:NH1	4:B:606:HOH:O	2.35	0.41
1:B:213:LYS:HD2	1:B:260:PHE:CZ	2.55	0.41
1:B:80:LYS:HE3	4:B:329:HOH:O	2.15	0.41
1:B:89:ARG:HG3	1:B:125:THR:HG22	2.03	0.41
1:B:161:VAL:CG1	1:B:225:LYS:HD2	2.51	0.41
1:B:39:LYS:HE2	1:B:39:LYS:HB3	1.47	0.41
1:B:27[A]:ARG:CG	1:B:205:GLU:HB3	2.52	0.40
1:B:22[A]:ILE:HD12	1:B:22[A]:ILE:HA	1.97	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	272/258 (105%)	262 (96%)	10 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	238/223 (107%)	234 (98%)	4 (2%)	53	17

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	61	ASN
1	B	92	GLN
1	B	228	LYS
1	B	255	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	17	HIS
1	B	61	ASN
1	B	249	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EVD	B	1	2	11,14,14	0.96	0	17,21,21	1.71	4 (23%)
3	EVD	B	263	-	11,14,14	2.10	4 (36%)	17,21,21	2.09	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EVD	B	1	2	-	1/4/6/6	0/2/2/2
3	EVD	B	263	-	-	0/4/6/6	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	263	EVD	C6-C7	3.64	1.45	1.38
3	B	263	EVD	O1-S	-3.19	1.38	1.43
3	B	263	EVD	C7-C8	3.14	1.46	1.39
3	B	263	EVD	S-N	2.73	1.64	1.58

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	263	EVD	C4-C9-C8	4.06	123.83	118.59
3	B	263	EVD	C6-C5-C4	3.59	124.67	120.24
3	B	263	EVD	C5-C4-C9	-3.36	115.87	120.95
3	B	263	EVD	C8-C9-C3	3.22	108.95	105.64
3	B	263	EVD	C6-C7-C8	-3.18	113.82	119.60
3	B	1	EVD	C8-C9-C3	3.15	108.89	105.64
3	B	1	EVD	C6-C5-C4	2.62	123.47	120.24
3	B	1	EVD	O2-S-N	2.47	111.83	108.39
3	B	1	EVD	C5-C4-C9	-2.24	117.57	120.95

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1	EVD	C3-C2-S-O1

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	258/258 (100%)	-0.17	6 (2%) 61 62	3, 10, 21, 30	15 (5%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	253	ASN	4.3
1	B	3	HIS	2.6
1	B	36	HIS	2.6
1	B	42	PRO	2.5
1	B	37	THR	2.0
1	B	40	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EVD	B	263	13/13	0.97	0.08	10,13,16,16	13

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EVD	B	1	13/13	0.99	0.03	5,6,7,8	0
2	ZN	B	262	1/1	1.00	0.01	5,5,5,5	0

6.5 Other polymers [i](#)

There are no such residues in this entry.