



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 04:46 PM EDT

PDB ID : 6S71 / pdb_00006s71
Title : Crystal structure of CARM1 in complex with inhibitor WH5C
Authors : Gunnell, E.A.; Muhsen, U.; Dowden, J.; Dreveny, I.
Deposited on : 2019-07-04
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	92% 5%
1	B	351	92% 5%
1	C	351	3% 92%
1	D	351	4% 89% 7%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-arginine methyltransferase CARM1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	2	0
			2757	1781	454	506	16			
1	B	342	Total	C	N	O	S	0	1	0
			2754	1778	456	506	14			
1	C	342	Total	C	N	O	S	0	5	0
			2781	1795	463	507	16			
1	D	342	Total	C	N	O	S	0	1	0
			2747	1774	453	505	15			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	480	HIS	-	expression tag	UNP Q86X55
A	481	HIS	-	expression tag	UNP Q86X55
A	482	HIS	-	expression tag	UNP Q86X55
A	483	HIS	-	expression tag	UNP Q86X55
A	484	HIS	-	expression tag	UNP Q86X55
A	485	HIS	-	expression tag	UNP Q86X55
B	480	HIS	-	expression tag	UNP Q86X55
B	481	HIS	-	expression tag	UNP Q86X55
B	482	HIS	-	expression tag	UNP Q86X55
B	483	HIS	-	expression tag	UNP Q86X55
B	484	HIS	-	expression tag	UNP Q86X55
B	485	HIS	-	expression tag	UNP Q86X55
C	480	HIS	-	expression tag	UNP Q86X55
C	481	HIS	-	expression tag	UNP Q86X55
C	482	HIS	-	expression tag	UNP Q86X55
C	483	HIS	-	expression tag	UNP Q86X55
C	484	HIS	-	expression tag	UNP Q86X55
C	485	HIS	-	expression tag	UNP Q86X55
D	480	HIS	-	expression tag	UNP Q86X55
D	481	HIS	-	expression tag	UNP Q86X55
D	482	HIS	-	expression tag	UNP Q86X55

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	483	HIS	-	expression tag	UNP Q86X55
D	484	HIS	-	expression tag	UNP Q86X55
D	485	HIS	-	expression tag	UNP Q86X55

- Molecule 2 is (2 {S})-4-[[[(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyloxy)-2-yl]methyl-(5-carbamimidamidopentyl)amino]-2-azanyl-butanoic acid (CCD ID: KYE) (formula: C₂₀H₃₄N₁₀O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			35	20	10	5		
2	B	1	Total	C	N	O	0	0
			35	20	10	5		
2	C	1	Total	C	N	O	0	0
			35	20	10	5		
2	D	1	Total	C	N	O	0	0
			35	20	10	5		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	129	Total	O	0	0
			129	129		
4	B	164	Total	O	0	0
			164	164		
4	C	103	Total	O	0	0
			103	103		
4	D	138	Total	O	0	0
			138	138		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone-arginine methyltransferase CARM1





- Molecule 1: Histone-arginine methyltransferase CARM1



- Molecule 1: Histone-arginine methyltransferase CARM1



- Molecule 1: Histone-arginine methyltransferase CARM1



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	74.72 Å 98.63 Å 206.36 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	103.18 – 2.06 103.18 – 2.06	Depositor EDS
% Data completeness (in resolution range)	96.6 (103.18-2.06) 91.6 (103.18-2.06)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.07 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.186 , 0.210 0.188 , 0.214	Depositor DCC
R_{free} test set	2000 reflections (2.18%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11755	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.43 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.8456e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KYE, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	2/2830 (0.1%)	0.83	0/3833
1	B	0.82	1/2825 (0.0%)	0.84	2/3828 (0.1%)
1	C	0.75	2/2861 (0.1%)	0.80	1/3874 (0.0%)
1	D	0.82	2/2820 (0.1%)	0.83	1/3821 (0.0%)
All	All	0.80	7/11336 (0.1%)	0.83	4/15356 (0.0%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	159	GLN	CD-OE1	6.26	1.35	1.23
1	D	465	ASN	CG-OD1	5.99	1.34	1.23
1	B	159	GLN	CD-NE2	-5.88	1.21	1.33
1	C	446	GLN	CD-OE1	5.70	1.34	1.23
1	A	159	GLN	CD-NE2	-5.63	1.21	1.33
1	D	465	ASN	CG-ND2	-5.29	1.22	1.33
1	C	234	ARG	CD-NE	-5.12	1.39	1.46

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	168	ARG	NE-CZ-NH1	-9.40	112.10	121.50
1	B	168	ARG	NE-CZ-NH2	9.10	127.39	119.20
1	C	234	ARG	NE-CZ-NH2	-5.86	113.93	119.20
1	D	168	ARG	NE-CZ-NH2	-5.05	114.65	119.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2757	0	2709	13	0
1	B	2754	0	2698	14	0
1	C	2781	0	2737	8	0
1	D	2747	0	2697	22	0
2	A	35	0	0	3	0
2	B	35	0	0	2	0
2	C	35	0	0	0	0
2	D	35	0	0	1	0
3	A	12	0	16	3	0
3	B	12	0	16	1	0
3	C	6	0	8	0	0
3	D	12	0	16	1	0
4	A	129	0	0	2	0
4	B	164	0	0	1	0
4	C	103	0	0	1	0
4	D	138	0	0	1	0
All	All	11755	0	10897	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:313:TRP:CH2	1:D:163:MET:HE1	1.95	1.00
1:D:148:GLN:OE1	3:D:502:GOL:O3	1.96	0.84
1:A:352:SER:HB3	3:A:503:GOL:H12	1.64	0.80
1:C:284:ASN:HB2	4:C:687:HOH:O	1.83	0.79
1:A:197:ILE:HG22	2:A:501:KYE:O25	1.88	0.74
1:D:420:CYS:SG	1:D:467:LEU:HD22	2.30	0.70
1:B:313:TRP:CZ2	1:D:163:MET:HE1	2.31	0.65
1:C:336:ARG:HG2	1:C:466:LEU:O	1.96	0.64
1:D:420:CYS:SG	1:D:467:LEU:CD2	2.86	0.64
1:B:312:PHE:CE2	1:D:163:MET:HE3	2.34	0.62
1:D:336:ARG:HG2	1:D:466:LEU:O	2.00	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:SER:CB	3:A:503:GOL:H12	2.31	0.61
1:D:159:GLN:C	1:D:163:MET:HE2	2.26	0.60
1:C:168:ARG:HD2	1:C:257:GLU:OE1	2.04	0.57
1:D:168:ARG:HD2	1:D:257:GLU:OE1	2.04	0.57
1:B:168:ARG:NH1	2:B:501:KYE:O24	2.36	0.57
1:A:247:LEU:HB3	4:A:609:HOH:O	2.04	0.56
1:A:268[B]:MET:HE1	2:A:501:KYE:C29	2.37	0.55
1:B:159:GLN:NE2	2:B:501:KYE:O25	2.38	0.54
1:D:337:GLN:NE2	4:D:606:HOH:O	2.36	0.50
1:D:195:SER:OG	1:D:197:ILE:HG12	2.12	0.48
1:D:188:VAL:HG22	1:D:210:ILE:HG23	1.95	0.48
1:B:312:PHE:HE2	1:D:163:MET:HE3	1.76	0.47
1:C:440:LEU:HD22	1:C:450:ILE:HG12	1.97	0.47
1:A:352:SER:H	3:A:503:GOL:H12	1.80	0.47
1:A:455:GLN:OE1	1:A:462:LYS:NZ	2.48	0.46
1:B:188:VAL:HG22	1:B:210:ILE:HG23	1.96	0.46
1:B:440:LEU:HD22	1:B:450:ILE:HG12	1.98	0.46
1:A:188:VAL:HG22	1:A:210:ILE:HG23	1.96	0.46
1:B:300:GLU:OE1	1:B:304:MET:HE3	2.16	0.45
1:A:440:LEU:HD22	1:A:450:ILE:HG12	1.99	0.45
1:D:160:GLN:HA	1:D:163:MET:CE	2.48	0.44
1:A:315:GLN:HG3	4:A:679:HOH:O	2.17	0.44
1:A:264:PHE:CE2	1:A:292:VAL:HG21	2.53	0.44
1:D:369:ARG:HG3	1:D:441:ILE:CD1	2.48	0.44
1:C:294:LEU:N	1:C:294:LEU:HD23	2.32	0.44
1:D:420:CYS:SG	1:D:467:LEU:HD21	2.57	0.44
1:D:338:PRO:HB3	1:D:467:LEU:HD21	1.99	0.43
1:D:455:GLN:HB3	1:D:462:LYS:HG3	2.00	0.43
1:D:294:LEU:N	1:D:294:LEU:HD12	2.33	0.43
1:A:197:ILE:CG2	2:A:501:KYE:O25	2.63	0.43
1:D:351:LYS:HD3	1:D:351:LYS:HA	1.89	0.43
1:C:313:TRP:O	1:C:324:SER:HA	2.19	0.42
1:C:369:ARG:HG3	1:C:441:ILE:CD1	2.50	0.41
1:D:313:TRP:O	1:D:324:SER:HA	2.20	0.41
1:B:264:PHE:CE2	1:B:292:VAL:HG21	2.56	0.41
1:B:148:GLN:OE1	3:B:503:GOL:O1	2.30	0.41
1:C:464:SER:C	1:C:465:ASN:CG	2.89	0.41
1:D:464:SER:C	1:D:465:ASN:CG	2.89	0.41
1:B:445:ARG:HH22	1:B:468:ASP:CG	2.29	0.40
1:B:294:LEU:HD12	1:B:294:LEU:N	2.36	0.40
1:B:269:LEU:HB2	4:B:723:HOH:O	2.20	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:349[A]:MET:SD	1:A:384:VAL:HG22	2.62	0.40
2:D:501:KYE:O24	2:D:501:KYE:C20	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/351 (97%)	334 (98%)	8 (2%)	0	100	100
1	B	341/351 (97%)	333 (98%)	8 (2%)	0	100	100
1	C	345/351 (98%)	336 (97%)	9 (3%)	0	100	100
1	D	341/351 (97%)	333 (98%)	8 (2%)	0	100	100
All	All	1369/1404 (98%)	1336 (98%)	33 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/306 (98%)	296 (99%)	4 (1%)	61	62
1	B	299/306 (98%)	294 (98%)	5 (2%)	53	53
1	C	303/306 (99%)	298 (98%)	5 (2%)	53	53

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	299/306 (98%)	291 (97%)	8 (3%)	39	35
All	All	1201/1224 (98%)	1179 (98%)	22 (2%)	51	51

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	142	GLU
1	A	148	GLN
1	A	188	VAL
1	A	294	LEU
1	B	188	VAL
1	B	208	ARG
1	B	209	LYS
1	B	294	LEU
1	B	424	SER
1	C	168	ARG
1	C	294	LEU
1	C	429	LYS
1	C	463	SER
1	C	465	ASN
1	D	168	ARG
1	D	188	VAL
1	D	208	ARG
1	D	220	GLN
1	D	294	LEU
1	D	429	LYS
1	D	444	LYS
1	D	467	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	159	GLN
1	A	164	GLN
1	A	315	GLN
1	A	368	HIS
1	A	471	ASN
1	B	164	GLN
1	B	380	HIS
1	B	471	ASN
1	C	164	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	380	HIS
1	C	455	GLN
1	D	164	GLN
1	D	220	GLN
1	D	337	GLN
1	D	471	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	502	-	5,5,5	0.11	0	5,5,5	0.63	0
3	GOL	B	503	-	5,5,5	0.21	0	5,5,5	0.52	0
3	GOL	C	502	-	5,5,5	0.19	0	5,5,5	0.26	0
2	KYE	A	501	-	36,37,37	2.92	11 (30%)	45,51,51	1.92	11 (24%)
3	GOL	D	502	-	5,5,5	0.14	0	5,5,5	0.48	0
3	GOL	A	503	-	5,5,5	0.39	0	5,5,5	0.27	0
2	KYE	C	501	-	36,37,37	2.38	9 (25%)	45,51,51	1.99	12 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	502	-	5,5,5	0.19	0	5,5,5	0.37	0
2	KYE	B	501	-	36,37,37	2.77	12 (33%)	45,51,51	2.00	14 (31%)
2	KYE	D	501	-	36,37,37	2.77	13 (36%)	45,51,51	2.04	12 (26%)
3	GOL	D	503	-	5,5,5	0.28	0	5,5,5	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	502	-	-	2/4/4/4	-
3	GOL	B	503	-	-	0/4/4/4	-
3	GOL	C	502	-	-	1/4/4/4	-
2	KYE	A	501	-	-	6/26/42/42	0/3/3/3
3	GOL	D	502	-	-	2/4/4/4	-
3	GOL	A	503	-	-	2/4/4/4	-
2	KYE	C	501	-	-	6/26/42/42	0/3/3/3
3	GOL	A	502	-	-	2/4/4/4	-
2	KYE	B	501	-	-	11/26/42/42	0/3/3/3
2	KYE	D	501	-	-	8/26/42/42	0/3/3/3
3	GOL	D	503	-	-	0/4/4/4	-

All (45) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	KYE	C03-C02	-10.62	1.24	1.53
2	A	501	KYE	C03-C02	-10.52	1.24	1.53
2	D	501	KYE	C03-C02	-10.52	1.24	1.53
2	C	501	KYE	C03-C02	-9.03	1.28	1.53
2	A	501	KYE	O16-C05	-7.75	1.24	1.42
2	D	501	KYE	O16-C05	-7.03	1.25	1.42
2	B	501	KYE	O16-C05	-7.00	1.25	1.42
2	C	501	KYE	O16-C05	-5.64	1.29	1.42
2	A	501	KYE	C11-N12	4.44	1.45	1.34
2	A	501	KYE	C15-N06	-4.33	1.30	1.37
2	B	501	KYE	C11-N12	4.29	1.45	1.34
2	C	501	KYE	C11-N12	4.26	1.45	1.34
2	D	501	KYE	C11-N12	4.23	1.45	1.34
2	A	501	KYE	C18-C17	-4.14	1.38	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	KYE	O24-C23	4.02	1.34	1.22
2	B	501	KYE	O24-C23	3.86	1.33	1.22
2	A	501	KYE	O24-C23	3.77	1.33	1.22
2	A	501	KYE	C13-N14	-3.63	1.32	1.39
2	B	501	KYE	C13-N14	-3.46	1.32	1.39
2	D	501	KYE	C03-C05	3.43	1.64	1.53
2	D	501	KYE	C15-N06	-3.42	1.31	1.37
2	C	501	KYE	C03-C05	3.33	1.64	1.53
2	B	501	KYE	C18-C17	-3.31	1.41	1.51
2	A	501	KYE	O01-C02	3.26	1.51	1.43
2	D	501	KYE	C18-C17	-3.21	1.41	1.51
2	B	501	KYE	C15-N06	-3.15	1.32	1.37
2	D	501	KYE	C13-C07	-3.12	1.33	1.39
2	D	501	KYE	C13-N14	-3.09	1.33	1.39
2	A	501	KYE	C03-C05	3.07	1.63	1.53
2	D	501	KYE	O24-C23	3.03	1.31	1.22
2	B	501	KYE	C03-C05	2.99	1.62	1.53
2	A	501	KYE	C13-C07	-2.94	1.33	1.39
2	B	501	KYE	O01-C02	2.81	1.49	1.43
2	C	501	KYE	C18-C17	-2.79	1.43	1.51
2	C	501	KYE	O01-C02	2.68	1.49	1.43
2	D	501	KYE	O01-C02	2.51	1.49	1.43
2	B	501	KYE	C13-C07	-2.46	1.34	1.39
2	D	501	KYE	O16-C17	2.36	1.50	1.45
2	A	501	KYE	C09-N08	-2.22	1.30	1.33
2	B	501	KYE	C09-N08	-2.18	1.30	1.33
2	C	501	KYE	C07-N08	2.12	1.38	1.34
2	D	501	KYE	C09-N08	-2.09	1.30	1.33
2	C	501	KYE	O16-C17	2.08	1.49	1.45
2	D	501	KYE	C07-N06	-2.07	1.33	1.37
2	B	501	KYE	C11-N10	-2.00	1.30	1.35

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	KYE	C13-C07-N08	-5.15	119.62	126.72
2	D	501	KYE	C13-C07-N08	-4.94	119.92	126.72
2	D	501	KYE	N10-C09-N08	-4.77	121.36	128.58
2	C	501	KYE	C13-C07-N08	-4.66	120.30	126.72
2	A	501	KYE	C13-C07-N08	-4.65	120.31	126.72
2	A	501	KYE	N10-C09-N08	-4.49	121.78	128.58
2	C	501	KYE	O01-C02-C17	-4.41	98.42	111.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	KYE	N10-C09-N08	-4.13	122.33	128.58
2	A	501	KYE	N06-C15-N14	-4.07	108.16	113.94
2	C	501	KYE	N06-C15-N14	-4.06	108.17	113.94
2	B	501	KYE	N10-C09-N08	-4.03	122.48	128.58
2	D	501	KYE	O16-C17-C18	3.97	115.37	108.88
2	A	501	KYE	C13-N14-C15	3.91	109.59	103.45
2	D	501	KYE	C09-N08-C07	3.78	121.07	111.83
2	D	501	KYE	N06-C15-N14	-3.75	108.61	113.94
2	B	501	KYE	O25-C23-O24	-3.73	115.62	124.08
2	B	501	KYE	C13-N14-C15	3.67	109.22	103.45
2	C	501	KYE	C13-N14-C15	3.66	109.21	103.45
2	A	501	KYE	C07-C13-N14	-3.60	106.47	110.58
2	B	501	KYE	N06-C15-N14	-3.55	108.90	113.94
2	D	501	KYE	N08-C07-N06	3.52	133.15	127.17
2	C	501	KYE	N08-C07-N06	3.44	133.01	127.17
2	A	501	KYE	C09-N08-C07	3.43	120.20	111.83
2	D	501	KYE	C27-N19-C20	-3.33	103.51	111.44
2	B	501	KYE	N08-C07-N06	3.31	132.80	127.17
2	B	501	KYE	C09-N08-C07	3.21	119.66	111.83
2	B	501	KYE	C07-C13-N14	-3.12	107.02	110.58
2	D	501	KYE	C13-N14-C15	3.08	108.29	103.45
2	C	501	KYE	C09-N08-C07	3.07	119.34	111.83
2	A	501	KYE	N08-C07-N06	3.01	132.28	127.17
2	C	501	KYE	C07-C13-N14	-2.98	107.17	110.58
2	C	501	KYE	C07-N06-C15	2.91	108.79	105.74
2	D	501	KYE	O25-C23-O24	-2.82	117.68	124.08
2	A	501	KYE	O25-C23-O24	-2.65	118.07	124.08
2	B	501	KYE	O16-C05-C03	-2.61	101.03	106.62
2	D	501	KYE	C07-N06-C15	2.57	108.44	105.74
2	A	501	KYE	C07-N06-C15	2.52	108.39	105.74
2	D	501	KYE	C07-C13-N14	-2.50	107.72	110.58
2	B	501	KYE	N32-C33-N35	2.40	124.80	120.67
2	B	501	KYE	C27-N19-C20	-2.36	105.81	111.44
2	C	501	KYE	C03-C02-C17	2.30	107.05	102.61
2	A	501	KYE	O01-C02-C17	-2.29	104.51	111.08
2	A	501	KYE	C17-O16-C05	-2.25	104.49	109.47
2	B	501	KYE	O01-C02-C17	-2.22	104.70	111.08
2	B	501	KYE	O16-C17-C18	2.22	112.51	108.88
2	C	501	KYE	N34-C33-N32	2.12	124.08	119.27
2	D	501	KYE	O01-C02-C17	-2.12	105.00	111.08
2	B	501	KYE	C03-C02-C17	2.09	106.64	102.61
2	C	501	KYE	O01-C02-C03	2.05	118.40	111.82

There are no chirality outliers.

All (40) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	KYE	C02-C17-C18-N19
2	A	501	KYE	O16-C17-C18-N19
2	A	501	KYE	N19-C20-C21-C22
2	B	501	KYE	C02-C17-C18-N19
2	B	501	KYE	O16-C17-C18-N19
2	B	501	KYE	N19-C20-C21-C22
2	B	501	KYE	C20-C21-C22-C23
2	B	501	KYE	C20-C21-C22-N26
2	B	501	KYE	C28-C27-N19-C18
2	C	501	KYE	O16-C17-C18-N19
2	D	501	KYE	C02-C17-C18-N19
2	D	501	KYE	O16-C17-C18-N19
3	A	502	GOL	O1-C1-C2-C3
3	B	502	GOL	C1-C2-C3-O3
3	D	502	GOL	O1-C1-C2-C3
2	C	501	KYE	C29-C30-C31-N32
2	D	501	KYE	C17-C18-N19-C20
2	A	501	KYE	N19-C27-C28-C29
3	A	502	GOL	O1-C1-C2-O2
3	B	502	GOL	O2-C2-C3-O3
3	D	502	GOL	O1-C1-C2-O2
2	C	501	KYE	C28-C29-C30-C31
2	B	501	KYE	C28-C29-C30-C31
2	B	501	KYE	C27-C28-C29-C30
3	A	503	GOL	O1-C1-C2-O2
2	B	501	KYE	C29-C30-C31-N32
2	C	501	KYE	N19-C20-C21-C22
2	D	501	KYE	C28-C29-C30-C31
2	D	501	KYE	C21-C22-C23-O24
3	C	502	GOL	O2-C2-C3-O3
2	C	501	KYE	C27-C28-C29-C30
2	B	501	KYE	C30-C31-N32-C33
3	A	503	GOL	O1-C1-C2-C3
2	C	501	KYE	C02-C17-C18-N19
2	B	501	KYE	C17-C18-N19-C20
2	D	501	KYE	N19-C27-C28-C29
2	D	501	KYE	C30-C31-N32-C33
2	A	501	KYE	C27-C28-C29-C30
2	D	501	KYE	N19-C20-C21-C22
2	A	501	KYE	C28-C27-N19-C20


There are no ring outliers.


6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	503	GOL	1	0
2	A	501	KYE	3	0
3	D	502	GOL	1	0
3	A	503	GOL	3	0
2	B	501	KYE	2	0
2	D	501	KYE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	342/351 (97%)	0.28	2 (0%) 85 88	23, 46, 64, 108	2 (0%)
1	B	342/351 (97%)	-0.04	4 (1%) 76 78	19, 35, 54, 73	1 (0%)
1	C	342/351 (97%)	0.33	10 (2%) 53 55	21, 46, 67, 97	5 (1%)
1	D	342/351 (97%)	0.13	13 (3%) 44 44	25, 40, 61, 80	1 (0%)
All	All	1368/1404 (97%)	0.17	29 (2%) 63 66	19, 42, 64, 108	9 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	314	TYR	5.8
1	C	476	TYR	4.1
1	A	476	TYR	3.5
1	D	314	TYR	3.5
1	D	420	CYS	3.4
1	D	467	LEU	3.2
1	B	464	SER	3.1
1	D	476	TYR	3.0
1	B	476	TYR	3.0
1	C	474	PHE	2.9
1	D	307	PHE	2.8
1	D	466	LEU	2.8
1	C	307	PHE	2.8
1	D	469	LEU	2.7
1	D	470	LYS	2.6
1	C	137	PHE	2.6
1	D	327	ARG	2.6
1	C	414[A]	HIS	2.5
1	D	377	HIS	2.4
1	D	137	PHE	2.4
1	A	136	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	136	VAL	2.4
1	B	140	ARG	2.2
1	C	470	LYS	2.2
1	C	464	SER	2.1
1	C	261	TYR	2.1
1	D	136	VAL	2.0
1	D	246	SER	2.0
1	C	466	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	502	6/6	0.78	0.14	72,80,82,86	0
3	GOL	A	503	6/6	0.78	0.16	60,64,66,71	0
3	GOL	C	502	6/6	0.80	0.20	77,79,80,80	0
3	GOL	D	502	6/6	0.82	0.16	64,66,67,70	0
2	KYE	A	501	35/35	0.84	0.15	45,58,87,92	0
3	GOL	B	503	6/6	0.86	0.15	70,72,77,77	0
2	KYE	B	501	35/35	0.86	0.15	42,52,82,86	0
3	GOL	B	502	6/6	0.86	0.12	60,63,65,66	0
3	GOL	D	503	6/6	0.87	0.14	67,68,72,79	0
2	KYE	D	501	35/35	0.91	0.12	30,39,59,62	0
2	KYE	C	501	35/35	0.93	0.11	38,44,79,83	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around KYE A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around KYE B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around KYE D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around KYE C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.