



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:02 PM UTC

PDB ID : 6S7S / pdb_00006s7s
Title : Fumarate hydratase of Mycobacterium tuberculosis in complex with formate and allosteric modulator N-(2-Methoxy-5-(N-phenylsulfamoyl)phenyl)-2-(4-oxo-3,4-dihydrophthalazin-1-yl)acetamide
Authors : Whitehouse, A.J.; Libardo, M.D.; Kasbekar, M.; Brear, P.; Fischer, G.; Thomas, C.J.; Barry, C.E.; Boshoff, H.I.; Coyne, A.G.; Abell, C.
Deposited on : 2019-07-05
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	3% 90% 5%
1	B	474	8% 89% 8%
1	C	474	4% 88% 7% 5%
1	D	474	5% 88% 6% 5%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 15135 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fumarate hydratase class II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	450	Total	C	N	O	S	0	11	0
			3440	2144	624	660	12			
1	B	458	Total	C	N	O	S	0	15	0
			3532	2195	644	679	14			
1	C	451	Total	C	N	O	S	0	15	0
			3474	2162	633	668	11			
1	D	448	Total	C	N	O	S	0	9	0
			3413	2125	623	653	12			

- Molecule 2 is {N}-[2-methoxy-5-(phenylsulfamoyl)phenyl]-2-(4-oxidanylidene-3 {H}-phthalazin-1-yl)ethanamide (CCD ID: KZ8) (formula: C₂₃H₂₀N₄O₅S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			33	23	4	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	S	0	0
			33	23	4	5	1		

- Molecule 3 is FORMIC ACID (CCD ID: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			3	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	341	Total	O	0	0
			341	341		
4	B	307	Total	O	0	0
			307	307		
4	C	281	Total	O	0	0
			281	281		
4	D	278	Total	O	0	0
			278	278		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Fumarate hydratase class II




• Molecule 1: Fumarate hydratase class II




• Molecule 1: Fumarate hydratase class II



• Molecule 1: Fumarate hydratase class II





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	175.57Å 96.92Å 124.75Å 90.00° 102.86° 90.00°	Depositor
Resolution (Å)	121.62 – 1.70 121.62 – 1.70	Depositor EDS
% Data completeness (in resolution range)	85.3 (121.62-1.70) 85.2 (121.62-1.70)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.186 , 0.217 0.186 , 0.217	Depositor DCC
R_{free} test set	9648 reflections (4.28%)	wwPDB-VP
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.194	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15135	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, KZ8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.09	0/3498	1.34	2/4751 (0.0%)
1	B	1.10	1/3581 (0.0%)	1.36	4/4864 (0.1%)
1	C	1.10	1/3524 (0.0%)	1.38	4/4788 (0.1%)
1	D	1.07	0/3459	1.38	2/4698 (0.0%)
All	All	1.09	2/14062 (0.0%)	1.36	12/19101 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	139	SER	C-O	8.10	1.33	1.24
1	B	378	ASN	C-O	5.38	1.30	1.24

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	141	ASP	CA-CB-CG	6.22	118.82	112.60
1	A	111	ASN	CA-CB-CG	5.70	118.30	112.60
1	D	15	ASP	CA-CB-CG	5.64	118.24	112.60
1	A	141	ASP	CA-CB-CG	5.59	118.19	112.60
1	D	111	ASN	CA-CB-CG	5.58	118.18	112.60
1	C	15	ASP	CA-CB-CG	5.46	118.06	112.60
1	B	141	ASP	CA-CB-CG	5.45	118.05	112.60
1	C	435	ILE	N-CA-C	-5.33	105.30	110.42
1	B	362	ILE	N-CA-CB	5.17	114.48	110.45
1	B	111	ASN	CA-CB-CG	5.16	117.76	112.60
1	C	67	ASP	CA-CB-CG	5.06	117.66	112.60
1	B	15	ASP	CA-CB-CG	5.03	117.63	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3440	0	3467	20	0
1	B	3532	0	3564	22	0
1	C	3474	0	3508	23	0
1	D	3413	0	3451	18	0
2	A	33	0	0	0	0
2	C	33	0	0	1	0
3	B	3	0	1	0	0
4	A	341	0	0	5	0
4	B	307	0	0	4	0
4	C	281	0	0	4	0
4	D	278	0	0	5	0
All	All	15135	0	13991	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:826:HOH:O	1:C:102:THR:HG21	1.67	0.94
1:A:215[C]:ARG:CA	1:A:216:ALA:N	2.52	0.72
1:A:214:VAL:C	1:A:215[C]:ARG:CA	2.63	0.71
1:B:360:VAL:HG22	4:B:820:HOH:O	1.92	0.70
1:C:437[A]:GLN:NE2	1:C:441[A]:ASP:OD1	2.23	0.70
1:D:360:VAL:HG22	4:D:675:HOH:O	1.91	0.70
1:D:243[B]:VAL:HG23	4:D:692:HOH:O	1.95	0.65
1:D:10:TYR:HB2	1:D:21:ARG:HD3	1.83	0.61
1:A:360:VAL:HG22	4:A:841:HOH:O	2.01	0.60
1:D:310[A]:GLN:NE2	4:D:503:HOH:O	2.35	0.58
1:A:50[B]:GLN:HE21	1:A:50[B]:GLN:CA	2.17	0.58
1:D:308:GLU:HB3	1:D:398:LEU:HD21	1.87	0.56
1:A:20:VAL:HG22	1:A:33:GLN:HG2	1.88	0.56
1:A:10:TYR:CD1	1:A:10:TYR:O	2.59	0.56
1:C:165:GLN:OE1	1:C:381:ARG:NH1	2.40	0.55
1:C:360:VAL:HG22	4:C:798:HOH:O	2.05	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:406:PRO:HD2	4:C:657:HOH:O	2.08	0.53
1:A:323:GLY:HA2	4:A:869:HOH:O	2.08	0.52
1:B:337:ALA:O	1:B:341[B]:ILE:HD13	2.09	0.52
1:C:406:PRO:HB3	1:C:425:ALA:HB1	1.92	0.52
1:D:12:ILE:HG21	1:D:19:GLU:OE2	2.10	0.51
1:A:50[A]:GLN:HG2	1:A:370:LEU:CD1	2.41	0.50
1:A:50[B]:GLN:CA	1:A:50[B]:GLN:NE2	2.73	0.50
1:B:102[B]:THR:HG21	1:B:106:THR:CG2	2.41	0.50
1:A:215[C]:ARG:HD2	4:A:606:HOH:O	2.11	0.49
1:A:215[A]:ARG:C	1:A:215[A]:ARG:N	2.70	0.49
2:C:501:KZ8:O	2:C:501:KZ8:C22	2.61	0.49
1:B:341[A]:ILE:HD13	1:C:349:TRP:CH2	2.48	0.49
1:B:20:VAL:HG22	1:B:34:ARG:CZ	2.42	0.48
1:D:20:VAL:HG22	1:D:34:ARG:CZ	2.44	0.48
1:B:354:GLY:HA2	4:B:820:HOH:O	2.13	0.47
1:A:341[B]:ILE:HD13	1:D:349:TRP:CH2	2.49	0.47
1:D:143:PHE:N	1:D:144:PRO:HD2	2.30	0.47
1:A:50[B]:GLN:HE21	1:A:50[B]:GLN:C	2.23	0.47
1:C:173[B]:LYS:NZ	1:C:391:LEU:O	2.48	0.47
1:C:313:ASP:O	1:C:314:LEU:HD23	2.16	0.46
1:C:173[A]:LYS:HD3	1:C:173[A]:LYS:HA	1.69	0.46
1:B:433:LYS:HE3	1:B:441:ASP:OD2	2.16	0.46
1:C:106[B]:THR:CB	1:C:139:SER:HB3	2.46	0.46
1:A:448:ARG:NH2	4:A:620:HOH:O	2.48	0.46
1:A:50[B]:GLN:NE2	1:A:50[B]:GLN:HA	2.31	0.46
1:C:445:ILE:HD11	1:C:454:LEU:HD22	1.99	0.45
1:B:11:ARG:NH1	1:B:26:ALA:O	2.46	0.45
1:B:408:ILE:O	1:B:411:PRO:HD2	2.17	0.45
1:B:452:GLU:CD	1:B:452:GLU:C	2.85	0.44
1:D:218:LEU:N	1:D:219:PRO:CD	2.81	0.44
1:D:408:ILE:HG12	1:D:463:MET:HE1	1.99	0.44
1:C:408:ILE:O	1:C:411:PRO:HD2	2.18	0.44
1:A:143:PHE:N	1:A:144:PRO:HD2	2.33	0.44
1:B:415:ALA:O	1:B:448:ARG:HD2	2.17	0.43
1:C:218:LEU:N	1:C:219:PRO:CD	2.81	0.43
1:B:102[B]:THR:HG21	1:B:106:THR:HG22	2.00	0.43
1:B:55:GLY:O	1:B:83:ALA:HB1	2.19	0.43
1:C:173[B]:LYS:HA	1:C:173[B]:LYS:HD3	1.81	0.43
1:D:15:ASP:O	1:D:17:MET:N	2.52	0.43
1:D:410:THR:HB	1:D:411:PRO:HD3	2.00	0.43
1:D:15:ASP:O	1:D:18:GLY:N	2.46	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:15:ASP:HB3	1:B:34:ARG:HH21	1.83	0.42
1:B:451:ILE:HD12	1:B:451:ILE:HA	1.93	0.42
1:A:329[A]:LEU:HA	1:A:329[A]:LEU:HD23	1.79	0.42
1:B:11:ARG:NH2	1:B:13:GLU:OE2	2.49	0.42
1:C:336:VAL:HG13	1:C:375:LEU:HB3	2.02	0.42
1:A:301:GLY:HA3	1:A:302:PRO:HA	1.85	0.42
1:B:102[B]:THR:CG2	1:B:106:THR:HG22	2.49	0.42
1:C:199:GLU:OE2	1:D:357:GLU:OE2	2.38	0.42
1:D:10:TYR:CD1	1:D:10:TYR:N	2.88	0.41
1:D:215[A]:ARG:HD2	4:D:544:HOH:O	2.20	0.41
1:C:329[A]:LEU:HA	1:C:329[A]:LEU:HD12	1.65	0.41
1:B:324:LYS:HG3	4:D:540:HOH:O	2.20	0.41
1:A:260[B]:THR:HG22	4:A:819:HOH:O	2.20	0.41
1:A:341[B]:ILE:HD13	1:D:349:TRP:CZ2	2.56	0.41
1:B:143:PHE:N	1:B:144:PRO:HD2	2.35	0.41
1:B:215[C]:ARG:NH1	4:B:606:HOH:O	2.43	0.41
1:C:74:GLU:H	1:C:74:GLU:CD	2.29	0.41
1:C:102:THR:HB	4:C:667:HOH:O	2.19	0.41
1:C:243[B]:VAL:HG23	4:C:799:HOH:O	2.21	0.41
1:B:463[A]:MET:HE2	1:B:463[A]:MET:HB3	1.94	0.40
1:C:106[B]:THR:HB	1:C:139:SER:HB3	2.02	0.40
1:B:349:TRP:CH2	1:C:341:ILE:HD13	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/474 (97%)	449 (98%)	8 (2%)	1 (0%)	43	28
1	B	472/474 (100%)	459 (97%)	12 (2%)	1 (0%)	43	28
1	C	462/474 (98%)	449 (97%)	13 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	453/474 (96%)	441 (97%)	11 (2%)	1 (0%)	43	28
All	All	1845/1896 (97%)	1798 (98%)	44 (2%)	3 (0%)	43	28

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	449	LEU
1	B	449	LEU
1	D	16	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/364 (98%)	349 (98%)	7 (2%)	48	32
1	B	367/364 (101%)	363 (99%)	4 (1%)	65	54
1	C	360/364 (99%)	355 (99%)	5 (1%)	59	46
1	D	352/364 (97%)	348 (99%)	4 (1%)	65	54
All	All	1435/1456 (99%)	1415 (99%)	20 (1%)	68	46

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	SER
1	A	215[A]	ARG
1	A	215[D]	ARG
1	A	215[C]	ARG
1	A	341[A]	ILE
1	A	341[B]	ILE
1	A	430	LYS
1	B	12	ILE
1	B	138	SER
1	B	329	LEU
1	B	379	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	139	SER
1	C	329[A]	LEU
1	C	329[B]	LEU
1	C	369[A]	ILE
1	C	369[B]	ILE
1	D	310[A]	GLN
1	D	310[B]	GLN
1	D	341	ILE
1	D	379	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	164	GLN
1	A	165	GLN
1	A	268	GLN
1	A	339	GLN
1	A	368	ASN
1	A	385	GLN
1	B	31	GLN
1	B	236	ASN
1	B	339	GLN
1	B	343	ASN
1	B	368	ASN
1	C	252	GLN
1	C	339	GLN
1	C	343	ASN
1	C	368	ASN
1	D	158	HIS
1	D	164	GLN
1	D	339	GLN
1	D	368	ASN
1	D	394	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	KZ8	C	501	-	36,36,36	0.48	0	46,51,51	0.65	0
3	FMT	B	501	-	2,2,2	0.34	0	1,1,1	0.08	0
2	KZ8	A	501	-	36,36,36	0.45	0	46,51,51	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	KZ8	C	501	-	-	4/21/21/21	0/4/4/4
2	KZ8	A	501	-	-	4/21/21/21	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	KZ8	C16-C-S-O4

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	501	KZ8	C16-C-S-O4
2	A	501	KZ8	C1-C-S-N3
2	C	501	KZ8	C1-C-S-O4
2	A	501	KZ8	C1-C-S-O4
2	A	501	KZ8	C16-C-S-N3
2	C	501	KZ8	C1-C-S-N3
2	C	501	KZ8	C16-C-S-N3


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	501	KZ8	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	450/474 (94%)	0.12	16 (3%) 46 50	9, 24, 51, 113	11 (2%)
1	B	458/474 (96%)	0.25	36 (7%) 18 20	7, 25, 68, 114	15 (3%)
1	C	451/474 (95%)	0.24	21 (4%) 36 40	10, 25, 59, 107	15 (3%)
1	D	448/474 (94%)	0.24	25 (5%) 30 33	9, 25, 63, 111	9 (2%)
All	All	1807/1896 (95%)	0.21	98 (5%) 31 34	7, 25, 63, 114	50 (2%)

All (98) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	215[B]	ARG	7.7
1	A	10	TYR	6.6
1	D	10	TYR	6.5
1	C	314	LEU	6.1
1	B	10	TYR	5.5
1	A	314	LEU	5.3
1	C	10	TYR	5.2
1	C	322	PRO	5.1
1	D	12	ILE	5.0
1	D	418	TYR	5.0
1	A	12	ILE	4.9
1	B	317	GLY	4.6
1	B	445	ILE	4.5
1	C	16	THR	4.4
1	C	12	ILE	4.2
1	B	17	MET	4.0
1	B	15	ASP	4.0
1	B	325	VAL	3.9
1	C	323	GLY	3.8
1	A	20	VAL	3.8
1	B	14	HIS	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	24	ALA	3.7
1	A	14	HIS	3.6
1	B	12	ILE	3.6
1	A	16	THR	3.6
1	D	14	HIS	3.6
1	D	415	ALA	3.5
1	C	14	HIS	3.5
1	B	16	THR	3.5
1	B	320	ILE	3.5
1	D	20	VAL	3.5
1	C	139	SER	3.5
1	D	16	THR	3.5
1	A	323	GLY	3.4
1	D	17	MET	3.3
1	A	418	TYR	3.2
1	B	316	PRO	3.2
1	A	17	MET	3.1
1	C	20	VAL	3.1
1	B	418	TYR	3.1
1	C	25	LYS	3.1
1	D	314	LEU	3.1
1	B	432	ARG	3.0
1	A	139	SER	3.0
1	B	322	PRO	3.0
1	C	22	VAL	2.9
1	A	18	GLY	2.9
1	A	440	ILE	2.9
1	D	22	VAL	2.9
1	C	24	ALA	2.9
1	B	448	ARG	2.8
1	D	15	ASP	2.8
1	C	448	ARG	2.8
1	B	466	ALA	2.8
1	B	440	ILE	2.8
1	D	18	GLY	2.8
1	B	429	LEU	2.7
1	B	321	MET	2.7
1	C	15	ASP	2.7
1	B	24	ALA	2.7
1	D	123	GLY	2.6
1	D	416	ILE	2.6
1	D	426	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	447	ASP	2.5
1	B	451	ILE	2.5
1	B	446	GLY	2.5
1	B	467	GLU	2.4
1	B	430	LYS	2.4
1	D	120	ALA	2.4
1	B	20	VAL	2.4
1	C	125	VAL	2.4
1	A	15	ASP	2.3
1	B	449	LEU	2.3
1	C	17	MET	2.3
1	D	429	LEU	2.3
1	D	122	LYS	2.3
1	A	467	GLU	2.2
1	A	24	ALA	2.2
1	B	318	SER	2.2
1	B	323	GLY	2.2
1	C	313	ASP	2.2
1	D	445	ILE	2.2
1	D	451	ILE	2.2
1	B	22	VAL	2.1
1	B	18	GLY	2.1
1	B	313	ASP	2.1
1	B	452	GLU	2.1
1	C	121	ALA	2.1
1	B	319	SER	2.1
1	B	21	ARG	2.1
1	C	23	PRO	2.1
1	D	119	ILE	2.1
1	C	123	GLY	2.0
1	A	448	ARG	2.0
1	D	23	PRO	2.0
1	B	80	ILE	2.0
1	D	121	ALA	2.0
1	D	25	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FMT	B	501	3/3	0.94	0.16	35,35,35,38	0
2	KZ8	C	501	33/33	0.97	0.07	21,24,47,48	0
2	KZ8	A	501	33/33	0.97	0.07	20,26,46,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around KZ8 C 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around KZ8 A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.