



wwPDB EM Validation Summary Report ⓘ

Mar 20, 2026 – 05:04 AM UTC

PDB ID : 7S78 / pdb_00007s78
EMDB ID : EMD-24881
Title : Structure of a cell-entry defective human adenovirus provides insights into precursor proteins and capsid maturation
Authors : Reddy, V.S.; Yu, X.
Deposited on : 2021-09-15
Resolution : 3.72 Å(reported)
Based on initial model : 3IYN

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	10474 (3.22 - 4.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	952	
1	B	952	
1	C	952	
1	D	952	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	952	 62%30% . . .
1	F	952	 63%29%6% . .
1	G	952	 66%27% . .
1	H	952	 65%29% . .
1	I	952	 65%27%5% .
1	J	952	 63%30%5% . .
1	K	952	 65%28% . . .
1	L	952	 62%31% . .
2	N	571	 45%27%8% . 18%
3	M	585	 44%15% . 39%
4	P	140	 11%64%25% . . .
4	Q	140	 12%70%17%6%6%
4	R	140	 6%58%12% . 28%
4	S	140	 53%16% . 31%
5	U	227	 44%23%6%27%
5	V	227	 7%46%26%8%19%
6	0	250	 94% . .
6	1	250	 9%11% . 77%
6	2	250	 92% . .
6	3	250	 14%8% . 76%
6	4	250	 94% . .
6	W	250	 5%5% . . 86%
6	X	250	 19%9% . 69%
6	Y	250	 6%11%6% . 76%
6	Z	250	 5% . . . 90%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	5	16	50% 100%
8	6	10	20% 100%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 104715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hexon protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	929	Total	C	N	O	S	0	0
			7427	4718	1258	1415	36		
1	B	929	Total	C	N	O	S	0	0
			7429	4719	1258	1416	36		
1	C	933	Total	C	N	O	S	0	0
			7456	4736	1262	1422	36		
1	D	929	Total	C	N	O	S	0	0
			7427	4718	1258	1415	36		
1	E	926	Total	C	N	O	S	0	0
			7408	4708	1255	1409	36		
1	F	929	Total	C	N	O	S	0	0
			7430	4721	1258	1415	36		
1	G	931	Total	C	N	O	S	0	0
			7441	4726	1260	1419	36		
1	H	933	Total	C	N	O	S	0	0
			7455	4736	1262	1420	37		
1	I	927	Total	C	N	O	S	0	0
			7417	4713	1256	1412	36		
1	J	928	Total	C	N	O	S	0	0
			7419	4713	1256	1414	36		
1	K	931	Total	C	N	O	S	0	0
			7442	4728	1260	1418	36		
1	L	929	Total	C	N	O	S	0	0
			7427	4718	1258	1415	36		

- Molecule 2 is a protein called Penton protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	N	466	Total	C	N	O	S	0	0
			3734	2365	646	711	12		

- Molecule 3 is a protein called Pre-hexon-linking protein IIIa.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	M	359	Total	C	N	O	S	0	0
			2813	1752	517	535	9		

- Molecule 4 is a protein called Hexon-interlacing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	P	134	Total	C	N	O	S	0	0
			972	600	171	199	2		
4	Q	131	Total	C	N	O	S	0	0
			952	587	168	195	2		
4	R	101	Total	C	N	O	S	0	0
			751	470	128	151	2		
4	S	97	Total	C	N	O	S	0	0
			727	448	128	149	2		

- Molecule 5 is a protein called Pre-hexon-linking protein VIII.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	U	165	Total	C	N	O	S	0	0
			1268	797	223	243	5		
5	V	184	Total	C	N	O	S	0	0
			1414	890	248	272	4		

- Molecule 6 is a protein called Pre-protein VI.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	W	35	Total	C	N	O	S	0	0
			262	165	48	47	2		
6	X	78	Total	C	N	O	S	0	0
			577	361	107	106	3		
6	Y	61	Total	C	N	O	S	0	0
			485	305	90	87	3		
6	Z	25	Total	C	N	O	S	0	0
			179	109	35	34	1		
6	0	16	Total	C	N	O	S	0	0
			114	68	23	22	1		
6	1	57	Total	C	N	O	S	0	0
			449	279	84	83	3		
6	2	19	Total	C	N	O	S	0	0
			133	81	26	25	1		
6	3	60	Total	C	N	O	S	0	0
			471	293	87	88	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
6	4	15	Total	C	N	O	S	0	0
			106	64	21	20	1		

- Molecule 7 is a protein called Unknown-1.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	5	16	Total	C	N	O	0	0
			80	48	16	16		


- Molecule 8 is a protein called Unknown-2.

Mol	Chain	Residues	Atoms				AltConf	Trace
8	6	10	Total	C	N	O	0	0
			50	30	10	10		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hexon protein




- Molecule 1: Hexon protein






Metabolites	V513	L416	R322	K197	K113	Met
	D514	P323	T198	A1		
	C515	R324	F199	W9	Y120	
	V516	R325	P200		K126	
	V517	T326	Q201			
	V518	A327	E202		M130	
	L519	F328	Q203		H13	
	A521	R329	Q204		P131	
	A522	K440	D330		C132	
	R521	N331	W210		E133	
Metabolites	M528	T443	G334	H217	W134	
		E444			D135	
	N532	L445	L335		E136	
	P533	S446	R336	K224	A137	
	F534	Y338	W337		L138	
	N535	M449	N339			
	H536	E450				
	R537	T451		T227	E142	
	F538	R452	N343	C232	I1E	
	N539	M459	N355	Y236	ASN	
Metabolites	R543	E460	A356	A237	GLU	
	V544	T461	V357	K238	T36	
	R545	M462	V358	P239	GLU	
	S546	L463	D359		ASP	
				N243	ASP	
	L549	N466	R363		ASN	
	G550	R469	N364	Q246	GLU	
	N551		T365		ASP	
	G552	Y473	E366	L249	GLU	
	V554	S474	Q370	T250	VAL	
Metabolites	V555	M475	L371	K251	ASP	
		T476	L372	Q252	GLU	
	Q560	A477	L373	N254	GLN	
		L478	D374	C255	A160	
	Q563	Y479	S375	R256	Q163	
	K564	L480	L376	L257	K164	
	F565	P481	G377	E258	T165	
	F566	D482	R378	S259	H165	
	A567	K483	R379	Q260	V167	
	L568	L484	T380	V261	F168	
Metabolites	K569	K486	E262		G169	
	N570	Y486	F383		Q170	
	L571			N276		
	L572	L493	A389	G276	M177	
	L573	S494	N390	D277	L178	
	L574	D495	D391	N278		
	F584	N496	S392		I183	
		P497		S287		
	D587	Y500		E288	V187	
	V588	D501	T400	D289	E188	
Metabolites	N589	Y502	E401	V290	G189	
	N590		N402	T304	Q190	
	V591	R506	H403	L305	P191	
	L592	Y507	K306	E307	P192	
	Q593	V508	E406		K193	
	S594				Y194	
	F595	L512	G412	N291	A195	
					F196	
					T111	
					F112	




• Molecule 1: Hexon protein

Chain D: 65% 27% 5% •




• Molecule 1: Hexon protein

Chain E: 62% 30% • • •



• Molecule 1: Hexon protein


Chain G: 66% 27%






• Molecule 1: Hexon protein


Chain H: 65% 29%



• Molecule 1: Hexon protein


Chain I: 65% 27% 5%






• Molecule 1: Hexon protein


Chain J: 63% 30% 5% ..






• Molecule 1: Hexon protein


Chain K: 65% 28%




• Molecule 1: Hexon protein

Chain L: 62% 31%







- Molecule 4: Hexon-interlacing protein




- Molecule 4: Hexon-interlacing protein



- Molecule 5: Pre-hexon-linking protein VIII




- Molecule 5: Pre-hexon-linking protein VIII







• Molecule 6: Pre-protein VI




• Molecule 6: Pre-protein VI




• Molecule 6: Pre-protein VI




- Molecule 6: Pre-protein VI



- Molecule 6: Pre-protein VI



- Molecule 6: Pre-protein VI




GLU
GLY
GLY
GLY
GLN
LYS
ARG
PRO
PRO
ARG
PRO
ASP
ARG
GLU
GLU
THR
LEU
VAL
THR
THR
GLN
TLE
ASP
GLU
PRO
PRO
PRO
TYR
GLU
GLU
ALA
ALA
LEU
LYS
GLN
GLY
PRO
ARG
LEU
SER
PRO
THR
THR
ARG
PRO
TLE
TLE
ALA
GLY
PRO
GLY
LEU
ASN
MET
ALA
THR
THR
GLY
VAL
LEU
GLY
GLN
HIS
THR
PRO
VAL
THR
LEU
ASP
LEU
PRO
PRO

PRO
ALA
ASP
THR
GLN
LYS
PRO
VAL
ARG
PRO
GLY
GLY
THR
ALA
VAL
VAL
THR
ARG
ASP
SER
ARG
SER
LEU
ARG
ALA
ALA
SER
GLY
PRO
ARG
SER
LEU
ARG
PRO
VAL
ALA
SER
GLY
ASN
TRP
GLN
SER
THR
LEU
ASN
SER
TLE
VAL
GLY
LEU
VAL
GLY
VAL
GLN
SER
LEU
PRO
LYS

ARG
ARG
ARG
CYS
PHE

● Molecule 7: Unknown-1



● Molecule 8: Unknown-2



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	11277	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	12	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	22500	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	36.862	Depositor
Minimum map value	-27.573	Depositor
Average map value	-0.024	Depositor
Map value standard deviation	3.164	Depositor
Recommended contour level	3.15	Depositor
Map size (Å)	1089.9199, 1089.9199, 1089.9199	wwPDB
Map dimensions	832, 832, 832	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.31, 1.31, 1.31	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	48/7626 (0.6%)	0.81	43/10371 (0.4%)
1	B	0.85	43/7628 (0.6%)	0.78	50/10374 (0.5%)
1	C	0.81	37/7655 (0.5%)	0.75	35/10411 (0.3%)
1	D	0.92	43/7626 (0.6%)	0.83	47/10371 (0.5%)
1	E	0.87	47/7606 (0.6%)	0.72	22/10343 (0.2%)
1	F	0.89	55/7629 (0.7%)	0.78	36/10374 (0.3%)
1	G	0.75	33/7640 (0.4%)	0.70	27/10391 (0.3%)
1	H	0.74	29/7654 (0.4%)	0.70	30/10409 (0.3%)
1	I	0.71	28/7615 (0.4%)	0.63	17/10355 (0.2%)
1	J	0.88	50/7617 (0.7%)	0.73	31/10357 (0.3%)
1	K	0.76	34/7641 (0.4%)	0.71	33/10392 (0.3%)
1	L	0.82	42/7626 (0.6%)	0.73	35/10371 (0.3%)
2	N	1.29	43/3827 (1.1%)	1.23	59/5215 (1.1%)
3	M	0.61	4/2869 (0.1%)	0.63	3/3908 (0.1%)
4	P	0.85	6/986 (0.6%)	0.98	13/1347 (1.0%)
4	Q	0.49	0/964	0.63	3/1315 (0.2%)
4	R	0.77	2/762 (0.3%)	0.65	4/1039 (0.4%)
4	S	0.23	0/736	0.43	0/1002
5	U	1.14	15/1300 (1.2%)	0.86	5/1764 (0.3%)
5	V	1.13	15/1452 (1.0%)	1.15	17/1977 (0.9%)
6	0	1.05	0/115	1.15	2/152 (1.3%)
6	1	1.67	5/460 (1.1%)	2.16	20/615 (3.3%)
6	2	1.17	1/135 (0.7%)	1.35	3/178 (1.7%)
6	3	0.81	0/482	1.28	7/646 (1.1%)
6	4	1.26	1/107 (0.9%)	1.34	2/141 (1.4%)
6	W	1.72	3/271 (1.1%)	2.32	16/365 (4.4%)
6	X	1.08	1/590 (0.2%)	1.46	14/794 (1.8%)
6	Y	1.62	8/498 (1.6%)	2.07	20/667 (3.0%)
6	Z	1.69	4/182 (2.2%)	1.98	11/242 (4.5%)
All	All	0.87	597/107299 (0.6%)	0.81	605/145886 (0.4%)

The worst 5 of 597 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	39	SER	CA-C	-11.57	1.37	1.52
6	W	30	MET	CA-C	-11.17	1.39	1.52
1	H	594	SER	CA-C	-10.50	1.39	1.52
1	J	915	LEU	CA-C	-9.54	1.41	1.52
1	F	649	ALA	CA-CB	-9.46	1.44	1.53

The worst 5 of 605 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	1	26	GLY	N-CA-C	24.37	144.68	111.54
6	3	26	GLY	N-CA-C	20.30	150.30	113.76
6	Y	26	GLY	N-CA-C	19.80	144.65	110.80
6	W	26	GLY	N-CA-C	18.11	137.94	110.96
1	A	45	ARG	N-CA-C	16.76	135.50	113.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7427	0	7124	218	0
1	B	7429	0	7126	212	0
1	C	7456	0	7156	224	0
1	D	7427	0	7124	213	0
1	E	7408	0	7112	226	0
1	F	7430	0	7131	222	0
1	G	7441	0	7138	205	0
1	H	7455	0	7159	218	0
1	I	7417	0	7118	228	0
1	J	7419	0	7116	216	0
1	K	7442	0	7142	211	0
1	L	7427	0	7127	236	0
2	N	3734	0	3656	107	0
3	M	2813	0	2767	58	0
4	P	972	0	978	31	0
4	Q	952	0	959	27	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	R	751	0	757	13	0
4	S	727	0	732	11	0
5	U	1268	0	1227	46	0
5	V	1414	0	1360	45	0
6	0	114	0	104	4	0
6	1	449	0	419	16	0
6	2	133	0	119	4	0
6	3	471	0	441	13	0
6	4	106	0	98	4	0
6	W	262	0	238	22	0
6	X	577	0	526	26	0
6	Y	485	0	458	31	0
6	Z	179	0	166	11	0
7	5	80	0	19	0	0
8	6	50	0	12	0	0
All	All	104715	0	100609	2692	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 2692 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:251:LYS:HG2	1:C:258:GLU:HB2	1.48	0.96
4:P:58:GLU:HA	4:P:61:ALA:HB3	1.52	0.92
6:W:32:GLY:C	6:W:34:ALA:H	1.71	0.91
2:N:51:GLY:HA2	2:N:116:ARG:NH2	1.87	0.90
1:B:635:ASN:HD22	3:M:18:PRO:HB3	1.37	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	925/952 (97%)	858 (93%)	65 (7%)	2 (0%)	43	71
1	B	925/952 (97%)	855 (92%)	69 (8%)	1 (0%)	48	78
1	C	929/952 (98%)	859 (92%)	69 (7%)	1 (0%)	48	78
1	D	925/952 (97%)	855 (92%)	69 (8%)	1 (0%)	48	78
1	E	922/952 (97%)	854 (93%)	63 (7%)	5 (0%)	24	56
1	F	925/952 (97%)	854 (92%)	66 (7%)	5 (0%)	24	56
1	G	927/952 (97%)	850 (92%)	76 (8%)	1 (0%)	48	78
1	H	929/952 (98%)	857 (92%)	71 (8%)	1 (0%)	48	78
1	I	923/952 (97%)	842 (91%)	78 (8%)	3 (0%)	36	65
1	J	924/952 (97%)	853 (92%)	66 (7%)	5 (0%)	24	56
1	K	927/952 (97%)	850 (92%)	75 (8%)	2 (0%)	43	71
1	L	925/952 (97%)	855 (92%)	67 (7%)	3 (0%)	36	65
2	N	462/571 (81%)	427 (92%)	31 (7%)	4 (1%)	14	45
3	M	355/585 (61%)	328 (92%)	26 (7%)	1 (0%)	36	65
4	P	132/140 (94%)	117 (89%)	14 (11%)	1 (1%)	16	47
4	Q	129/140 (92%)	112 (87%)	17 (13%)	0	100	100
4	R	97/140 (69%)	82 (84%)	15 (16%)	0	100	100
4	S	93/140 (66%)	87 (94%)	6 (6%)	0	100	100
5	U	161/227 (71%)	140 (87%)	21 (13%)	0	100	100
5	V	178/227 (78%)	148 (83%)	26 (15%)	4 (2%)	5	30
6	0	12/250 (5%)	11 (92%)	0	1 (8%)	0	9
6	1	53/250 (21%)	45 (85%)	8 (15%)	0	100	100
6	2	15/250 (6%)	10 (67%)	4 (27%)	1 (7%)	1	13
6	3	56/250 (22%)	52 (93%)	3 (5%)	1 (2%)	6	33
6	4	11/250 (4%)	6 (54%)	5 (46%)	0	100	100
6	W	33/250 (13%)	28 (85%)	4 (12%)	1 (3%)	3	26
6	X	74/250 (30%)	56 (76%)	15 (20%)	3 (4%)	2	20
6	Y	57/250 (23%)	46 (81%)	8 (14%)	3 (5%)	1	17
6	Z	21/250 (8%)	17 (81%)	1 (5%)	3 (14%)	0	3
All	All	13045/15844 (82%)	11954 (92%)	1038 (8%)	53 (0%)	31	60

5 of 53 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	733	ASN
1	D	733	ASN
1	F	197	LYS
1	F	203	PRO
2	N	495	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	806/829 (97%)	743 (92%)	63 (8%)	11	37
1	B	807/829 (97%)	748 (93%)	59 (7%)	13	39
1	C	809/829 (98%)	746 (92%)	63 (8%)	11	37
1	D	806/829 (97%)	760 (94%)	46 (6%)	18	45
1	E	804/829 (97%)	744 (92%)	60 (8%)	12	39
1	F	807/829 (97%)	745 (92%)	62 (8%)	12	37
1	G	808/829 (98%)	760 (94%)	48 (6%)	18	44
1	H	809/829 (98%)	760 (94%)	49 (6%)	17	44
1	I	805/829 (97%)	746 (93%)	59 (7%)	13	39
1	J	804/829 (97%)	755 (94%)	49 (6%)	17	44
1	K	808/829 (98%)	753 (93%)	55 (7%)	14	41
1	L	806/829 (97%)	748 (93%)	58 (7%)	13	40
2	N	423/489 (86%)	370 (88%)	53 (12%)	4	21
3	M	304/500 (61%)	276 (91%)	28 (9%)	8	32
4	P	107/112 (96%)	92 (86%)	15 (14%)	3	18
4	Q	104/112 (93%)	93 (89%)	11 (11%)	6	27
4	R	86/112 (77%)	78 (91%)	8 (9%)	8	32
4	S	85/112 (76%)	77 (91%)	8 (9%)	8	31
5	U	136/186 (73%)	121 (89%)	15 (11%)	6	26
5	V	152/186 (82%)	134 (88%)	18 (12%)	5	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
6	0	12/210 (6%)	10 (83%)	2 (17%)	2	14
6	1	47/210 (22%)	40 (85%)	7 (15%)	3	16
6	2	13/210 (6%)	12 (92%)	1 (8%)	12	37
6	3	50/210 (24%)	43 (86%)	7 (14%)	3	18
6	4	11/210 (5%)	11 (100%)	0	100	100
6	W	26/210 (12%)	20 (77%)	6 (23%)	1	6
6	X	54/210 (26%)	49 (91%)	5 (9%)	8	32
6	Y	50/210 (24%)	38 (76%)	12 (24%)	1	5
6	Z	18/210 (9%)	15 (83%)	3 (17%)	2	14
All	All	11357/13647 (83%)	10487 (92%)	870 (8%)	14	37

5 of 870 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	297	THR
1	K	339	ASN
5	U	55	LEU
1	I	441	ASP
1	I	284	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 116 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	260	GLN
6	Y	13	HIS
1	I	246	GLN
6	X	74	GLN
2	N	199	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-24881. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 416



Y Index: 416




Z Index: 416


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 431



Y Index: 431




Z Index: 431


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.15. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 68380 nm³; this corresponds to an approximate mass of 61769 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.269 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.269 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.72	-	-
Author-provided FSC curve	3.72	4.02	3.75
Unmasked-calculated*	-	-	-


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-24881 and PDB model 7S78. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlays

9.1.1 Map-model overlay [i](#)




9.1.2 Map-model assembly overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 3.15 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3.15).

9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (3.15) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9000	 0.4710
0	 0.7140	 0.4920
1	 0.7680	 0.4890
2	 0.6560	 0.5190
3	 0.7170	 0.4610
4	 0.6440	 0.4730
5	 0.5120	 0.4670
6	 0.6800	 0.4480
A	 0.9180	 0.4710
B	 0.9160	 0.4650
C	 0.9160	 0.4680
D	 0.9140	 0.4770
E	 0.9140	 0.4770
F	 0.9180	 0.4790
G	 0.9140	 0.4810
H	 0.9140	 0.4810
I	 0.9160	 0.4790
J	 0.9160	 0.4770
K	 0.9190	 0.4740
L	 0.9130	 0.4770
M	 0.8070	 0.4370
N	 0.8880	 0.4510
P	 0.7860	 0.4080
Q	 0.7960	 0.4230
R	 0.8270	 0.4160
S	 0.8740	 0.4330
U	 0.8340	 0.4650
V	 0.7900	 0.4650
W	 0.5590	 0.4460
X	 0.7530	 0.4480
Y	 0.6800	 0.4580
Z	 0.6860	 0.4860

