



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:22 PM UTC

PDB ID : 6S8L / pdb_00006s8l
Title : Structure, Thermodynamics, and Kinetics of Plinabulin Binding to two Tubulin Isotypes
Authors : Sharma, A.; Olieric, N.; Steinmetz, M.
Deposited on : 2019-07-10
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	
2	B	450	
3	F	157	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 8691 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	424	Total	C	N	O	S	0	1	0
			3334	2115	565	632	22			

- Molecule 2 is a protein called Tubulin beta-3 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	420	Total	C	N	O	S	0	4	0
			3320	2097	563	635	25			

- Molecule 3 is a protein called Designed ankyrin repeat protein (DARPIN) D1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	155	Total	C	N	O	S	0	2	0
			1163	736	198	226	3			

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		

- Molecule 6 is (3Z,6Z)-3-benzylidene-6-[(5-tert-butyl-1H-imidazol-4-yl)methylidene]piperazine-2,5-dione (CCD ID: PN6) (formula: C₁₉H₂₀N₄O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			25	19	4	2		

- Molecule 7 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tubulin alpha-1B chain



• Molecule 2: Tubulin beta-3 chain



• Molecule 3: Designed ankyrin repeat protein (DARPIN) D1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.63Å 91.20Å 82.64Å 90.00° 97.49° 90.00°	Depositor
Resolution (Å)	45.60 – 1.80 45.60 – 1.80	Depositor EDS
% Data completeness (in resolution range)	93.3 (45.60-1.80) 93.5 (45.60-1.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 1.79Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.215 , 0.266 0.218 , 0.268	Depositor DCC
R_{free} test set	4692 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	16.0	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8691	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PN6, GDP, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.19	0/3409	0.40	0/4627
2	B	0.18	0/3405	0.38	0/4611
3	F	0.19	0/1185	0.37	0/1611
All	All	0.19	0/7999	0.39	0/10849

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3334	0	3243	24	0
2	B	3320	0	3219	32	0
3	F	1163	0	1176	14	0
4	A	32	0	10	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	B	25	0	20	4	0
7	B	28	0	11	1	0
8	A	353	0	0	4	0
8	B	248	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	F	186	0	0	4	0
All	All	8691	0	7679	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338:LYS:O	1:A:340:SER:N	2.24	0.70
3:F:13:ASP:N	8:F:203:HOH:O	2.24	0.70
1:A:209:ILE:HG23	1:A:230:LEU:HD23	1.79	0.64
2:B:275:LEU:HD11	2:B:300:ASN:HA	1.79	0.63
1:A:413:MET:HE3	1:A:417:GLU:HB3	1.81	0.63
2:B:326:LYS:HD3	2:B:326:LYS:H	1.65	0.61
3:F:107:ASN:HD21	3:F:138:ASP:H	1.47	0.61
2:B:48:ARG:HG3	2:B:245:PRO:HD3	1.85	0.59
1:A:364:PRO:O	8:A:601:HOH:O	2.17	0.59
2:B:72:PRO:HG3	2:B:96:GLN:HA	1.87	0.57
2:B:4:ILE:HD11	2:B:242:LEU:HD23	1.87	0.57
3:F:107:ASN:HD21	3:F:137:ALA:HA	1.72	0.55
1:A:2:ARG:NH1	8:A:616:HOH:O	2.40	0.54
3:F:13:ASP:OD1	8:F:201:HOH:O	2.18	0.54
1:A:229:ARG:NH1	8:A:602:HOH:O	2.24	0.53
2:B:239:THR:HG22	6:B:501:PN6:C17	2.39	0.52
1:A:430:LYS:NZ	1:A:434:GLU:OE2	2.43	0.51
1:A:175:PRO:HA	1:A:178:SER:HB3	1.92	0.51
3:F:107:ASN:ND2	3:F:138:ASP:H	2.09	0.51
2:B:42:LEU:HA	2:B:47:GLU:HG2	1.94	0.50
2:B:215:ARG:NH1	3:F:160:ASP:OD1	2.45	0.49
2:B:27:GLU:OE1	2:B:236:SER:OG	2.27	0.48
1:A:311:LYS:NZ	8:A:624:HOH:O	2.46	0.47
2:B:395:PHE:CE1	2:B:422:GLU:HB2	2.49	0.47
3:F:121:ALA:HB1	3:F:161:LEU:HD21	1.96	0.47
2:B:2:ARG:N	8:B:621:HOH:O	2.48	0.47
2:B:124:LYS:HB2	2:B:124:LYS:HE3	1.61	0.45
2:B:147:SER:HB2	2:B:190:SER:OG	2.16	0.45
1:A:27:GLU:CD	1:A:320:ARG:HH22	2.24	0.45
2:B:315:VAL:HB	2:B:351:VAL:HG22	1.98	0.45
3:F:107:ASN:HD21	3:F:138:ASP:N	2.12	0.45
2:B:6:HIS:CD2	2:B:21:TRP:HE1	2.36	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:93:ILE:HD11	2:B:121:VAL:HG21	2.00	0.44
1:A:265:ILE:HD11	1:A:431:ASP:HB3	1.99	0.44
3:F:102:HIS:HD2	8:F:340:HOH:O	2.00	0.44
2:B:311:ARG:NH2	2:B:345:GLU:OE1	2.42	0.44
2:B:139:HIS:HE1	2:B:170:SER:OG	2.00	0.43
1:A:265:ILE:HG23	1:A:432:TYR:CE1	2.53	0.43
2:B:182:VAL:HG21	2:B:407[B]:TRP:HZ3	1.83	0.43
2:B:242:LEU:HD22	6:B:501:PN6:C16	2.49	0.43
2:B:255:LEU:HB3	6:B:501:PN6:C11	2.48	0.43
2:B:69:ASP:O	2:B:94:PHE:HA	2.18	0.43
1:A:142:GLY:HA3	1:A:183:GLU:OE1	2.19	0.43
1:A:430:LYS:O	1:A:434:GLU:HG3	2.19	0.43
3:F:159:GLU:O	3:F:163:GLU:HG2	2.19	0.42
2:B:308:ARG:NH1	8:B:617:HOH:O	2.47	0.42
1:A:385:ALA:HB2	1:A:432:TYR:CG	2.55	0.42
2:B:269:MET:HE1	2:B:307:PRO:HG3	2.00	0.42
2:B:332:MET:HG3	2:B:353:VAL:HG21	2.02	0.42
1:A:341:ILE:O	1:A:341:ILE:HG13	2.20	0.42
3:F:65:VAL:O	3:F:69:HIS:HD2	2.02	0.42
2:B:12:CYS:HB2	7:B:502:GDP:C8	2.54	0.42
2:B:250:ALA:HA	2:B:254:LYS:HD2	2.01	0.42
1:A:7:ILE:HG21	1:A:153:LEU:HD21	2.02	0.41
1:A:12:ALA:HB3	1:A:140:SER:HB3	2.01	0.41
2:B:255:LEU:HB3	6:B:501:PN6:N12	2.35	0.41
1:A:265:ILE:HG21	1:A:313:MET:HE1	2.02	0.41
3:F:23:ARG:NH2	8:F:204:HOH:O	2.30	0.41
3:F:61:GLU:H	3:F:61:GLU:CD	2.29	0.41
2:B:105:LYS:HA	2:B:109:THR:OG1	2.20	0.41
1:A:161:TYR:HB3	1:A:164:LYS:CG	2.50	0.40
1:A:221:ARG:CG	2:B:326:LYS:HG3	2.51	0.40
3:F:61:GLU:O	3:F:65:VAL:HG23	2.22	0.40
1:A:320:ARG:HA	1:A:356:ASN:O	2.20	0.40
1:A:25:CYS:HB3	1:A:30:ILE:O	2.21	0.40
1:A:150:THR:O	1:A:154:MET:HG2	2.21	0.40
2:B:75:MET:HG3	2:B:94:PHE:CD2	2.56	0.40
2:B:136:GLN:HA	2:B:167:ASN:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	417/451 (92%)	409 (98%)	8 (2%)	0	100	100
2	B	416/450 (92%)	408 (98%)	8 (2%)	0	100	100
3	F	155/157 (99%)	155 (100%)	0	0	100	100
All	All	988/1058 (93%)	972 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	360/379 (95%)	360 (100%)	0	100	100
2	B	365/386 (95%)	361 (99%)	4 (1%)	65	60
3	F	122/122 (100%)	121 (99%)	1 (1%)	73	70
All	All	847/887 (96%)	842 (99%)	5 (1%)	78	77

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	58	SER
2	B	238	VAL
2	B	248	LEU
2	B	326	LYS
3	F	127	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	101	ASN
1	A	197	HIS
1	A	342	GLN
1	A	393	HIS
2	B	91	ASN
2	B	139	HIS
2	B	167	ASN
2	B	426	ASN
3	F	36	ASN
3	F	107	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GTP	A	501	5	33,34,34	3.24	17 (51%)	50,54,54	2.25	14 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GDP	B	502	5	29,30,30	3.60	18 (62%)	45,47,47	1.94	12 (26%)
6	PN6	B	501	-	26,27,27	2.59	13 (50%)	34,39,39	2.32	11 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	A	501	5	-	7/22/38/38	0/3/3/3
7	GDP	B	502	5	-	5/16/32/32	0/3/3/3
6	PN6	B	501	-	-	3/14/14/14	0/3/3/3

All (48) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	502	GDP	C2'-C3'	-10.22	1.25	1.53
4	A	501	GTP	O4'-C1'	8.34	1.61	1.42
4	A	501	GTP	C2'-C1'	-7.26	1.30	1.53
6	B	501	PN6	C6-C2	6.22	1.55	1.40
7	B	502	GDP	O4'-C4'	-5.98	1.31	1.45
7	B	502	GDP	C4-N3	5.93	1.47	1.34
4	A	501	GTP	O4'-C4'	-5.72	1.32	1.45
4	A	501	GTP	C4-N3	5.64	1.47	1.34
7	B	502	GDP	C2-N3	5.28	1.46	1.33
7	B	502	GDP	C3'-C4'	5.10	1.65	1.53
7	B	502	GDP	C1'-N9	-5.10	1.33	1.47
4	A	501	GTP	C2-N2	4.76	1.45	1.34
4	A	501	GTP	C2-N3	4.64	1.44	1.33
7	B	502	GDP	C2-N2	4.44	1.44	1.34
6	B	501	PN6	C1-N5	-4.35	1.31	1.37
4	A	501	GTP	PB-O3B	4.30	1.64	1.59
4	A	501	GTP	PA-O3A	4.17	1.64	1.59
7	B	502	GDP	O4'-C1'	4.11	1.51	1.42
7	B	502	GDP	PA-O3A	4.10	1.63	1.59
6	B	501	PN6	C14-C13	4.06	1.54	1.46
6	B	501	PN6	C8-N9	3.98	1.46	1.38
6	B	501	PN6	C10-N9	3.78	1.46	1.38
4	A	501	GTP	PB-O3A	3.32	1.63	1.59
6	B	501	PN6	C7-N12	3.31	1.45	1.38
6	B	501	PN6	C22-C1	3.26	1.56	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	501	PN6	C11-N12	3.20	1.44	1.38
6	B	501	PN6	C2-N3	-3.07	1.31	1.39
7	B	502	GDP	O2'-C2'	3.00	1.50	1.43
7	B	502	GDP	C5-N7	-2.99	1.33	1.39
4	A	501	GTP	C4-N9	-2.88	1.30	1.38
4	A	501	GTP	C5-C6	2.83	1.55	1.44
4	A	501	GTP	C2-N1	2.83	1.44	1.37
4	A	501	GTP	O2'-C2'	2.78	1.49	1.43
6	B	501	PN6	O20-C8	-2.69	1.18	1.23
6	B	501	PN6	O21-C11	-2.66	1.18	1.23
4	A	501	GTP	O3'-C3'	-2.64	1.36	1.43
4	A	501	GTP	C6-N1	2.63	1.43	1.38
7	B	502	GDP	C2'-C1'	2.63	1.61	1.53
7	B	502	GDP	C5-C6	2.62	1.54	1.44
7	B	502	GDP	C2-N1	2.58	1.43	1.37
6	B	501	PN6	C10-C11	2.53	1.53	1.48
7	B	502	GDP	O6-C6	-2.42	1.19	1.23
6	B	501	PN6	C7-C8	2.37	1.53	1.48
4	A	501	GTP	O6-C6	-2.21	1.19	1.23
7	B	502	GDP	C6-N1	2.07	1.42	1.38
4	A	501	GTP	C8-N7	2.04	1.38	1.32
7	B	502	GDP	O3'-C3'	2.02	1.48	1.43
7	B	502	GDP	C4-N9	-2.01	1.32	1.38

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	501	GTP	C1'-N9-C8	-7.79	104.59	126.73
4	A	501	GTP	C1'-N9-C4	7.45	148.49	126.49
7	B	502	GDP	C5-C4-N3	-5.65	119.40	128.39
6	B	501	PN6	C10-N9-C8	-5.18	119.64	125.51
7	B	502	GDP	C2-N3-C4	4.87	120.69	112.30
6	B	501	PN6	C7-N12-C11	-4.81	120.06	125.51
6	B	501	PN6	C10-C11-N12	4.79	120.73	116.02
6	B	501	PN6	N5-C4-N3	-4.79	107.42	112.98
6	B	501	PN6	C7-C8-N9	4.40	120.35	116.02
4	A	501	GTP	C2-N3-C4	3.86	118.94	112.30
4	A	501	GTP	C6-C5-N7	3.58	136.80	130.29
6	B	501	PN6	C8-C7-N12	3.52	119.81	116.65
7	B	502	GDP	C1'-N9-C8	3.43	136.48	126.73
7	B	502	GDP	C2-N1-C6	-3.42	118.90	125.11
7	B	502	GDP	C1'-N9-C4	-3.41	116.41	126.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	501	GTP	C5-C4-N9	3.15	111.29	105.66
6	B	501	PN6	C11-C10-N9	3.07	119.41	116.65
6	B	501	PN6	C22-C1-C2	-3.03	128.70	135.52
4	A	501	GTP	N2-C2-N1	2.91	122.89	116.76
4	A	501	GTP	O5'-C5'-C4'	-2.89	99.17	108.99
6	B	501	PN6	C7-C6-C2	-2.88	120.99	128.06
7	B	502	GDP	O3'-C3'-C4'	-2.87	102.83	111.08
4	A	501	GTP	C4-C5-N7	-2.83	106.19	110.67
4	A	501	GTP	N9-C8-N7	-2.80	108.20	113.40
7	B	502	GDP	N9-C4-N3	2.75	131.45	125.95
4	A	501	GTP	C5-C4-N3	-2.73	124.05	128.39
7	B	502	GDP	N9-C8-N7	-2.68	108.43	113.40
7	B	502	GDP	C5-C6-N1	2.68	120.07	113.25
4	A	501	GTP	C4'-O4'-C1'	-2.38	104.22	109.47
7	B	502	GDP	C8-N7-C5	2.26	108.28	104.26
4	A	501	GTP	C5-C6-N1	2.25	118.99	113.25
6	B	501	PN6	O20-C8-C7	-2.25	118.81	121.91
7	B	502	GDP	C4-C5-N7	-2.23	107.13	110.67
7	B	502	GDP	C3'-C2'-C1'	2.15	105.54	101.46
4	A	501	GTP	C2-N1-C6	-2.07	121.36	125.11
6	B	501	PN6	C22-C1-N5	2.04	123.73	119.48
4	A	501	GTP	N1-C2-N3	-2.00	119.66	123.32

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	501	GTP	C5'-O5'-PA-O3A
4	A	501	GTP	C5'-O5'-PA-O1A
6	B	501	PN6	C2-C6-C7-C8
6	B	501	PN6	N9-C10-C13-C14
7	B	502	GDP	C5'-O5'-PA-O3A
7	B	502	GDP	C5'-O5'-PA-O1A
7	B	502	GDP	C5'-O5'-PA-O2A
4	A	501	GTP	PB-O3B-PG-O1G
7	B	502	GDP	PB-O3A-PA-O1A
4	A	501	GTP	C5'-O5'-PA-O2A
6	B	501	PN6	C11-C10-C13-C14
4	A	501	GTP	PB-O3B-PG-O2G
4	A	501	GTP	PB-O3B-PG-O3G
4	A	501	GTP	PB-O3A-PA-O2A
7	B	502	GDP	PB-O3A-PA-O2A


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	502	GDP	1	0
6	B	501	PN6	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	424/451 (94%)	0.49	37 (8%) 16 14	10, 22, 49, 83	1 (0%)
2	B	420/450 (93%)	0.74	55 (13%) 7 6	11, 26, 55, 85	4 (0%)
3	F	155/157 (98%)	0.07	3 (1%) 66 66	9, 19, 41, 63	2 (1%)
All	All	999/1058 (94%)	0.53	95 (9%) 14 12	9, 23, 52, 85	7 (0%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	262	TYR	9.4
1	A	179	THR	8.9
2	B	250	ALA	5.8
2	B	57	SER	5.6
2	B	440	ALA	5.2
1	A	341	ILE	5.2
1	A	348	PRO	5.1
2	B	245	PRO	5.1
2	B	325	MET	5.1
2	B	248	LEU	4.9
2	B	407[A]	TRP	4.6
2	B	39	ASP	4.5
1	A	349	THR	4.5
1	A	364	PRO	4.2
1	A	245	ASP	4.1
2	B	75	MET	4.1
1	A	340	SER	4.0
2	B	58	SER	4.0
2	B	59	HIS	3.9
1	A	337	THR	3.8
1	A	57	GLY	3.8
1	A	309	HIS	3.7
2	B	371	LEU	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	97	SER	3.7
2	B	335	ILE	3.7
2	B	42	LEU	3.5
1	A	350	GLY	3.5
1	A	436	GLY	3.4
1	A	36	MET	3.4
1	A	365	GLY	3.3
1	A	338	LYS	3.3
2	B	249	ASN	3.3
2	B	29	GLY	3.3
1	A	362	VAL	3.2
1	A	336	LYS	3.2
2	B	358	ILE	3.1
2	B	244	PHE	3.1
1	A	263	PRO	3.0
3	F	32	ILE	3.0
2	B	74	THR	3.0
2	B	96	GLN	3.0
2	B	369	ARG	2.9
1	A	347[A]	CYS	2.9
1	A	178	SER	2.9
2	B	406	HIS	2.9
2	B	334	ALA	2.9
3	F	167	LYS	2.9
2	B	415	GLU	2.9
2	B	412	GLY	2.9
1	A	247	ALA	2.8
2	B	128	ASN	2.8
2	B	86	LEU	2.8
2	B	333	LEU	2.8
2	B	322	ARG	2.8
2	B	242	LEU	2.7
1	A	59	GLY	2.7
2	B	338	LYS	2.7
1	A	335	ILE	2.6
2	B	41	ASP	2.6
2	B	56	ALA	2.6
2	B	72	PRO	2.6
2	B	251	ASP	2.5
2	B	127	GLU	2.5
2	B	370	GLY	2.5
1	A	58	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	355	VAL	2.5
1	A	430	LYS	2.5
1	A	37	PRO	2.4
1	A	284	GLU	2.4
1	A	261	PRO	2.4
2	B	36	TYR	2.4
2	B	76	ASP	2.4
3	F	29	GLU	2.4
2	B	337	SER	2.4
1	A	308	ARG	2.3
2	B	285	ALA	2.3
1	A	370	LYS	2.3
2	B	255	LEU	2.3
2	B	326	LYS	2.3
1	A	435	VAL	2.2
2	B	372	LYS	2.2
1	A	2	ARG	2.2
2	B	33	SER	2.2
2	B	272	PHE	2.2
1	A	297	GLU	2.2
2	B	38	GLY	2.2
2	B	278	ARG	2.2
2	B	77	SER	2.1
2	B	178	SER	2.1
1	A	47	ASP	2.1
2	B	2	ARG	2.0
2	B	82	ALA	2.0
1	A	279	GLU	2.0
1	A	296	PHE	2.0
2	B	276	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	PN6	B	501	25/25	0.84	0.17	19,27,32,34	25
5	MG	B	503	1/1	0.95	0.06	27,27,27,27	0
4	GTP	A	501	32/32	0.97	0.06	5,12,19,21	0
7	GDP	B	502	28/28	0.97	0.06	10,15,20,21	0
5	MG	A	502	1/1	0.98	0.03	15,15,15,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PN6 B 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GDP B 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.