



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:37 AM UTC

PDB ID : 8S98 / pdb_00008s98
Title : Crystal structure of the TYK2 pseudokinase domain in complex with compound 8
Authors : Toms, A.V.; Leit, S.; Greenwood, J.R.; Carriero, S.; Mondal, S.; Abel, R.; Ashwell, M.; Blanchette, H.; Boyles, N.; Cartwright, M.; Collis, A.; Feng, S.; Ghanakota, P.; Harriman, G.C.; Hosagrahara, V.; Kaila, N.; Kapeller, R.; Rafi, S.; Romero, D.L.; Tarantino, P.; Timaniya, J.; Wester, R.T.; Westlin, W.; Srivastava, B.; Miao, W.; Tummino, P.; McElwee, J.J.; Edmondson, S.D.; Massee, C.E.
Deposited on : 2023-03-27
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	: 4-5-2 with Phenix2.0
Mogul	: 2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	: 2.0
EDS	: 3.0
Buster-report	: wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	: 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	: 9.0.010 (Gargrove)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	5% 78% 9% 13%
1	B	295	7% 80% 7% 12%

Continued on next page...

Density-Fitness : 1.0.12
 Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.49

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	C	295	 A horizontal bar chart showing the quality of chain C. The bar is divided into four segments: a small red segment at the beginning labeled '6%', a large green segment labeled '78%', a small yellow segment labeled '10%', and a small grey segment at the end labeled '12%'.

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6885 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-receptor tyrosine-protein kinase TYK2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	1	0
			2025	1283	363	369	10			
1	B	259	Total	C	N	O	S	0	5	0
			2081	1315	376	379	11			
1	C	261	Total	C	N	O	S	0	5	0
			2075	1316	370	378	11			

- Molecule 2 is (8S)-N-cyclopropyl-5-[(2-methoxypyridin-3-yl)amino]-7-(methylamino)pyrazolo[1,5-a]pyrimidine-3-carboxamide (CCD ID: ZRU) (formula: C₁₇H₁₉N₇O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			26	17	7	2		
2	B	1	Total	C	N	O	0	0
			26	17	7	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			26	17	7	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	199	Total	O	0	0
			199	199		
3	B	221	Total	O	0	1
			222	222		
3	C	205	Total	O	0	0
			205	205		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Non-receptor tyrosine-protein kinase TYK2



- Molecule 1: Non-receptor tyrosine-protein kinase TYK2



- Molecule 1: Non-receptor tyrosine-protein kinase TYK2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.11Å 112.60Å 156.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.47 – 1.87 47.47 – 1.87	Depositor EDS
% Data completeness (in resolution range)	95.2 (47.47-1.87) 95.2 (47.47-1.87)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 1.87Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.225 , 0.274 (Not available) , 0.244	Depositor DCC
R_{free} test set	2213 reflections (3.10%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.753	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6885	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, ZRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	2/2062 (0.1%)	1.40	5/2795 (0.2%)
1	B	1.13	3/2121 (0.1%)	1.38	6/2876 (0.2%)
1	C	1.16	5/2114 (0.2%)	1.38	4/2868 (0.1%)
All	All	1.14	10/6297 (0.2%)	1.39	15/8539 (0.2%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	736	CYS	C-O	6.90	1.31	1.23
1	A	761	GLY	C-O	6.17	1.32	1.23
1	C	690	VAL	C-O	6.00	1.30	1.24
1	C	853	PRO	C-O	-5.69	1.17	1.24
1	B	690	VAL	C-O	5.56	1.30	1.24
1	C	658	THR	C-O	5.53	1.30	1.24
1	B	694	PRO	C-O	-5.27	1.17	1.23
1	B	688	GLU	C-O	5.14	1.30	1.23
1	C	592	LEU	C-O	5.06	1.30	1.24
1	C	639	VAL	C-O	5.00	1.29	1.23

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	777	PRO	N-CA-CB	-5.86	96.15	102.60
1	A	776	ILE	CB-CA-C	5.82	116.70	111.00
1	B	668	THR	CA-CB-OG1	-5.46	101.41	109.60
1	A	605	GLU	N-CA-C	-5.43	102.84	110.50
1	C	859	PHE	CA-CB-CG	-5.41	108.39	113.80
1	A	776	ILE	N-CA-C	-5.31	104.08	109.02
1	C	693	GLY	CA-C-N	5.28	125.19	119.76
1	C	693	GLY	C-N-CA	5.28	125.19	119.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	818	ARG	CA-C-N	5.20	129.18	120.86
1	B	818	ARG	C-N-CA	5.20	129.18	120.86
1	C	668	THR	CA-CB-OG1	-5.17	101.84	109.60
1	B	645	ASP	CA-CB-CG	5.16	117.76	112.60
1	A	859	PHE	CA-CB-CG	-5.13	108.67	113.80
1	A	636	GLU	CB-CG-CD	-5.13	103.88	112.60
1	B	599	THR	CA-CB-OG1	-5.09	101.97	109.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2025	0	2004	16	0
1	B	2081	0	2051	8	0
1	C	2075	0	2041	14	0
2	A	26	0	0	0	0
2	B	26	0	0	0	0
2	C	26	0	0	0	0
3	A	199	0	0	4	0
3	B	222	0	0	2	0
3	C	205	0	0	2	0
All	All	6885	0	6096	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:712:MET:HE1	1:A:841:LEU:HD11	1.45	0.98
1:A:791:LEU:N	3:A:1001:HOH:O	2.25	0.70
1:C:816:GLN:HG3	3:C:1193:HOH:O	1.96	0.66
1:A:712:MET:HE1	1:A:841:LEU:CD1	2.27	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:732:HIS:CE1	1:C:760:PRO:HB3	2.36	0.60
1:C:585:ASP:OD1	1:C:587:LYS:HG2	2.03	0.57
1:A:816:GLN:HE22	1:A:823:LYS:NZ	2.04	0.55
1:A:816:GLN:HG3	3:A:1162:HOH:O	2.08	0.53
1:C:607:ARG:HB3	1:C:636:GLU:HG3	1.90	0.52
1:B:609:ARG:HG3	1:B:609:ARG:HH11	1.76	0.51
1:A:816:GLN:NE2	1:A:816:GLN:HA	2.26	0.50
1:A:808:CSO:SG	1:A:841:LEU:HD12	2.52	0.50
1:C:685:MET:HG2	1:C:687:THR:CG2	2.42	0.49
1:A:712:MET:HE2	1:A:712:MET:HA	1.96	0.48
1:C:784:LEU:HB3	1:C:785:PRO:HD3	1.95	0.47
1:A:638:ARG:HD3	1:A:689:TYR:CE1	2.50	0.47
1:A:797:LYS:NZ	3:A:1006:HOH:O	2.39	0.46
1:B:784:LEU:HB3	1:B:785:PRO:HD3	1.97	0.46
1:C:685:MET:HE3	1:C:685:MET:HB2	1.86	0.45
1:B:672:PHE:CE1	1:C:701:ARG:CZ	3.00	0.45
1:C:838:CYS:HB3	3:C:1174:HOH:O	2.17	0.44
1:C:794:ALA:HA	1:C:797:LYS:HD2	1.99	0.44
1:A:816:GLN:HE22	1:A:823:LYS:HZ3	1.64	0.43
1:B:712:MET:HA	1:B:712:MET:HE2	1.99	0.43
3:B:1078:HOH:O	1:C:745:LEU:HD21	2.18	0.43
1:B:582[B]:HIS:CD2	1:C:705:HIS:NE2	2.87	0.42
1:A:740[A]:ILE:HD11	1:A:803:THR:HG21	2.01	0.42
1:A:645:ASP:O	1:A:646:PRO:C	2.63	0.42
1:B:656:TYR:OH	1:B:683:ASN:ND2	2.51	0.42
1:A:727:ASN:ND2	3:A:1013:HOH:O	2.53	0.41
1:C:747:LEU:HD23	1:C:747:LEU:HA	1.91	0.41
1:C:740:ILE:HD11	1:C:803:THR:HG21	2.01	0.41
1:A:647:SER:HB3	1:A:651:ILE:HB	2.03	0.41
1:B:685:MET:HB2	1:B:685:MET:HE3	1.83	0.41
1:A:813:ALA:HB3	1:A:816:GLN:HG2	2.03	0.40
1:B:852:GLU:N	3:B:1006:HOH:O	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/295 (84%)	243 (98%)	5 (2%)	0	100	100
1	B	257/295 (87%)	249 (97%)	8 (3%)	0	100	100
1	C	259/295 (88%)	252 (97%)	7 (3%)	0	100	100
All	All	764/885 (86%)	744 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/253 (86%)	215 (99%)	3 (1%)	59	48
1	B	224/253 (88%)	218 (97%)	6 (3%)	39	23
1	C	221/253 (87%)	219 (99%)	2 (1%)	70	63
All	All	663/759 (87%)	652 (98%)	11 (2%)	53	40

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	777	PRO
1	A	804	LEU
1	A	821	SER
1	B	649	HIS
1	B	679	ARG
1	B	752	SER
1	B	777	PRO
1	B	804	LEU
1	B	821	SER
1	C	841	LEU
1	C	868	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	597	GLN
1	A	674	HIS
1	A	727	ASN
1	A	734	ASN
1	A	816	GLN
1	A	855	GLN
1	B	597	GLN
1	B	602	ASN
1	B	674	HIS
1	B	683	ASN
1	B	734	ASN
1	C	727	ASN
1	C	734	ASN
1	C	739	ASN
1	C	830	GLN
1	C	847	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	A	808	1	3,6,7	0.76	0	1,6,8	0.43	0
1	CSO	C	808	1	3,6,7	0.73	0	1,6,8	0.56	0
1	CSO	B	808	1	3,6,7	0.72	0	1,6,8	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	808	1	-	0/1/5/7	-
1	CSO	C	808	1	-	0/1/5/7	-
1	CSO	B	808	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	808	CSO	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ZRU	C	901	-	29,29,29	1.97	10 (34%)	35,41,41	2.36	16 (45%)
2	ZRU	A	901	-	29,29,29	2.06	9 (31%)	35,41,41	2.72	17 (48%)
2	ZRU	B	901	-	29,29,29	2.00	10 (34%)	35,41,41	2.20	13 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ZRU	C	901	-	-	0/14/18/18	0/4/4/4
2	ZRU	A	901	-	-	0/14/18/18	0/4/4/4
2	ZRU	B	901	-	-	2/14/18/18	0/4/4/4

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	ZRU	C12-N15	-5.03	1.32	1.37
2	C	901	ZRU	C17-C18	-4.36	1.35	1.41
2	B	901	ZRU	C17-C18	-4.12	1.35	1.41
2	B	901	ZRU	C18-C21	-4.09	1.37	1.49
2	A	901	ZRU	C18-C19	-3.99	1.34	1.45
2	A	901	ZRU	C18-C21	-3.95	1.38	1.49
2	C	901	ZRU	C18-C21	-3.93	1.38	1.49
2	A	901	ZRU	C17-C18	-3.77	1.36	1.41
2	C	901	ZRU	C18-C19	-3.49	1.36	1.45
2	B	901	ZRU	C18-C19	-3.48	1.36	1.45
2	B	901	ZRU	C12-N15	-3.36	1.34	1.37
2	B	901	ZRU	C8-N9	-3.23	1.35	1.41
2	B	901	ZRU	C11-C12	3.16	1.43	1.38
2	A	901	ZRU	C11-C12	3.00	1.42	1.38
2	C	901	ZRU	C14-N13	2.94	1.50	1.45
2	C	901	ZRU	C17-N16	2.86	1.37	1.32
2	C	901	ZRU	C11-C12	2.82	1.42	1.38
2	B	901	ZRU	C17-N16	2.74	1.37	1.32
2	B	901	ZRU	C14-N13	2.54	1.50	1.45
2	B	901	ZRU	C10-N20	2.52	1.37	1.32
2	C	901	ZRU	C10-N20	2.46	1.37	1.32
2	C	901	ZRU	C5-N4	2.44	1.39	1.34
2	C	901	ZRU	C8-N9	-2.41	1.37	1.41
2	A	901	ZRU	C17-N16	2.41	1.36	1.32
2	C	901	ZRU	C25-C24	2.40	1.54	1.48
2	A	901	ZRU	C8-N9	-2.22	1.37	1.41
2	B	901	ZRU	C5-N4	2.19	1.39	1.34
2	A	901	ZRU	C5-N4	2.17	1.39	1.34
2	A	901	ZRU	C10-N20	2.11	1.36	1.32

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	ZRU	C18-C19-N15	6.74	109.11	105.58
2	A	901	ZRU	C26-C24-N23	-6.28	110.29	118.52
2	C	901	ZRU	C18-C19-N15	5.26	108.34	105.58
2	A	901	ZRU	C25-C24-N23	-4.91	112.08	118.52
2	B	901	ZRU	C17-N16-N15	4.72	106.75	103.43
2	B	901	ZRU	C25-C24-N23	-4.57	112.53	118.52
2	B	901	ZRU	C18-C19-N15	4.52	107.95	105.58
2	A	901	ZRU	C17-N16-N15	4.42	106.53	103.43
2	C	901	ZRU	N9-C10-N20	4.04	124.06	114.41
2	A	901	ZRU	C19-N15-N16	-4.03	107.99	112.24
2	C	901	ZRU	C19-N15-N16	-4.03	107.99	112.24
2	B	901	ZRU	N9-C10-N20	4.01	123.99	114.41
2	C	901	ZRU	C17-C18-C19	3.96	107.53	104.57
2	C	901	ZRU	C17-N16-N15	3.92	106.19	103.43
2	C	901	ZRU	C25-C24-N23	-3.73	113.63	118.52
2	C	901	ZRU	C26-C24-N23	-3.72	113.64	118.52
2	B	901	ZRU	C17-C18-C19	3.63	107.28	104.57
2	A	901	ZRU	C1-O2-C3	3.57	120.54	117.20
2	B	901	ZRU	C19-N15-N16	-3.56	108.49	112.24
2	A	901	ZRU	N9-C10-N20	3.43	122.60	114.41
2	A	901	ZRU	C10-N20-C19	3.36	121.95	117.18
2	C	901	ZRU	C12-N15-N16	3.22	129.25	125.79
2	A	901	ZRU	C18-C19-N20	3.03	131.77	123.54
2	B	901	ZRU	C26-C24-N23	-2.98	114.61	118.52
2	C	901	ZRU	C18-C19-N20	2.86	131.32	123.54
2	A	901	ZRU	N15-C19-N20	-2.74	119.11	122.15
2	B	901	ZRU	C18-C19-N20	2.69	130.84	123.54
2	A	901	ZRU	C17-C18-C19	2.65	106.55	104.57
2	C	901	ZRU	C11-C10-N20	-2.58	118.58	122.75
2	C	901	ZRU	C10-N20-C19	2.54	120.79	117.18
2	C	901	ZRU	C11-C12-N13	-2.53	121.97	127.81
2	A	901	ZRU	C11-C10-N20	-2.47	118.75	122.75
2	B	901	ZRU	C12-N15-N16	2.43	128.41	125.79
2	B	901	ZRU	C18-C17-N16	-2.36	109.52	112.45
2	C	901	ZRU	C7-C8-C3	2.30	119.06	116.84
2	B	901	ZRU	C11-C10-N20	-2.29	119.04	122.75
2	A	901	ZRU	C11-C12-N15	2.25	117.69	114.85
2	A	901	ZRU	C26-C25-C24	-2.24	57.85	59.80
2	B	901	ZRU	C7-C8-C3	2.23	119.00	116.84
2	A	901	ZRU	C18-C17-N16	-2.19	109.73	112.45
2	B	901	ZRU	C11-C12-N13	-2.13	122.89	127.81
2	C	901	ZRU	C1-O2-C3	-2.13	115.21	117.20
2	A	901	ZRU	C11-C12-N13	-2.12	122.93	127.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	901	ZRU	C11-C12-N15	2.11	117.51	114.85
2	A	901	ZRU	C12-N15-N16	2.05	128.00	125.79
2	C	901	ZRU	C18-C17-N16	-2.04	109.92	112.45

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	901	ZRU	C8-C3-O2-C1
2	B	901	ZRU	N4-C3-O2-C1

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ZRU C 901



Ligand ZRU A 901





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	255/295 (86%)	0.39	14 (5%)	30 33	7, 21, 38, 49	1 (0%)
1	B	258/295 (87%)	0.50	21 (8%)	18 20	7, 22, 38, 48	5 (1%)
1	C	260/295 (88%)	0.44	17 (6%)	25 27	7, 22, 37, 60	5 (1%)
All	All	773/885 (87%)	0.44	52 (6%)	24 26	7, 21, 38, 60	11 (1%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	681	PRO	4.0
1	B	817	SER	3.9
1	C	838	CYS	3.7
1	A	791	LEU	3.6
1	B	839	PRO	3.4
1	B	837	SER	3.4
1	C	678	VAL	3.3
1	A	841	LEU	3.3
1	B	838	CYS	3.2
1	C	840	GLN	3.1
1	C	837	SER	3.1
1	B	609	ARG	3.0
1	B	816	GLN	2.9
1	C	681	PRO	2.9
1	A	579	LEU	2.9
1	C	680	GLY	2.9
1	A	750	GLY	2.8
1	C	843	THR	2.8
1	C	836	PRO	2.7
1	B	809	PHE	2.7
1	A	865	ASP	2.6
1	A	816	GLN	2.6
1	C	839	PRO	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	578	GLN	2.6
1	B	680	GLY	2.6
1	A	842	ALA	2.6
1	B	842	ALA	2.5
1	B	815	LEU	2.5
1	C	868	ARG	2.5
1	A	635	GLN	2.4
1	A	836	PRO	2.4
1	B	812	GLU	2.4
1	B	813	ALA	2.4
1	C	842	ALA	2.4
1	A	578	GLN	2.3
1	B	579	LEU	2.3
1	B	841	LEU	2.3
1	B	701[A]	ARG	2.2
1	C	582	HIS	2.2
1	A	751	THR	2.2
1	A	747	LEU	2.2
1	B	678	VAL	2.2
1	C	656	TYR	2.2
1	C	817	SER	2.2
1	C	841	LEU	2.1
1	B	705	HIS	2.1
1	A	709	ALA	2.1
1	A	701	ARG	2.1
1	B	836	PRO	2.1
1	C	809	PHE	2.0
1	B	649	HIS	2.0
1	C	867	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	A	808	7/8	0.82	0.13	21,24,28,28	0
1	CSO	B	808	7/8	0.88	0.12	23,26,29,29	0
1	CSO	C	808	7/8	0.91	0.10	22,25,26,27	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZRU	A	901	26/26	0.89	0.10	12,14,15,15	0
2	ZRU	B	901	26/26	0.90	0.10	12,13,17,17	0
2	ZRU	C	901	26/26	0.90	0.10	12,15,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZRU A 901:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZRU B 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.