



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:56 AM UTC

PDB ID : 8S9D / pdb_00008s9d
Title : C143S variant of Citrate Synthase (CitA) in Mycobacterium tuberculosis
Authors : Pathirage, R.; Ronning, D.
Deposited on : 2023-03-27
Resolution : 2.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	379	17% 86% 11% ...

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	402	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FLC	A	404	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called citrate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	368	2788	1754	506	516	12	0	0	0

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	143	SER	CYS	engineered mutation	UNP A0A045JB88
A	374	HIS	-	expression tag	UNP A0A045JB88
A	375	HIS	-	expression tag	UNP A0A045JB88
A	376	HIS	-	expression tag	UNP A0A045JB88
A	377	HIS	-	expression tag	UNP A0A045JB88
A	378	HIS	-	expression tag	UNP A0A045JB88
A	379	HIS	-	expression tag	UNP A0A045JB88

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is CITRATE ANION (CCD ID: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	27	Total	O	0	0
			27	27		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: citrate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	150.25Å 150.25Å 231.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.85 – 2.57 50.85 – 2.57	Depositor EDS
% Data completeness (in resolution range)	99.6 (50.85-2.57) 89.6 (50.85-2.57)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.23 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158:000	Depositor
R, R_{free}	0.313 , 0.329 0.315 , 0.330	Depositor DCC
R_{free} test set	2003 reflections (4.36%)	wwPDB-VP
Wilson B-factor (Å ²)	49.5	Xtrriage
Anisotropy	0.116	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 70.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	2848	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.59 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.5266e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.18	0/2851	0.32	0/3883

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2788	0	2759	22	0
2	A	20	0	30	3	0
3	A	13	0	5	1	0
4	A	27	0	0	1	0
All	All	2848	0	2794	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:110:GLN:NE2	4:A:501:HOH:O	2.20	0.74
1:A:289:LEU:HB3	1:A:298:ILE:HD13	1.73	0.70
1:A:125:GLN:HG2	2:A:406:EDO:H12	1.71	0.69
1:A:158:PRO:HB2	1:A:163:ILE:HD11	1.74	0.68
1:A:149:ARG:HD2	2:A:403:EDO:H21	1.77	0.64
1:A:259:ARG:HH22	3:A:404:FLC:HG1	1.63	0.62
1:A:230:ASP:OD1	1:A:230:ASP:N	2.27	0.61
1:A:279:VAL:O	1:A:283:GLN:HG3	2.08	0.53
1:A:208:GLY:O	1:A:211:HIS:ND1	2.37	0.53
1:A:161:ARG:NH1	1:A:270:LEU:O	2.43	0.49
1:A:367:GLU:H	1:A:367:GLU:CD	2.21	0.47
1:A:170:TRP:HE1	1:A:323:MET:HE3	1.79	0.46
1:A:234:VAL:O	1:A:238:ILE:HG13	2.16	0.46
1:A:256:GLU:HA	1:A:299:GLU:HG2	1.99	0.45
1:A:252:VAL:HG23	1:A:253:TYR:H	1.82	0.44
1:A:185:ALA:HB2	1:A:202:ALA:HB2	2.00	0.44
1:A:252:VAL:HG23	1:A:253:TYR:N	2.33	0.43
1:A:187:VAL:HG22	1:A:346:VAL:HG21	2.01	0.43
1:A:314:ARG:HD3	1:A:314:ARG:HA	1.79	0.43
1:A:68:ALA:H	2:A:403:EDO:H22	1.85	0.41
1:A:236:LYS:HZ2	1:A:288:GLU:CD	2.28	0.41
1:A:32:LEU:HD23	1:A:39:ILE:HD13	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/379 (97%)	352 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	280/291 (96%)	270 (96%)	10 (4%)	31 56

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	SER
1	A	88	LEU
1	A	116	SER
1	A	138	ARG
1	A	144	SER
1	A	218	VAL
1	A	230	ASP
1	A	262	VAL
1	A	346	VAL
1	A	358	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	HIS
1	A	155	GLN
1	A	301	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	403	-	3,3,3	0.52	0	2,2,2	0.27	0
2	EDO	A	406	-	3,3,3	0.37	0	2,2,2	0.32	0
2	EDO	A	401	-	3,3,3	0.45	0	2,2,2	0.33	0
2	EDO	A	405	-	3,3,3	0.39	0	2,2,2	0.53	0
3	FLC	A	404	-	12,12,12	1.14	2 (16%)	17,17,17	1.17	2 (11%)
2	EDO	A	402	-	3,3,3	0.49	0	2,2,2	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	403	-	-	0/1/1/1	-
2	EDO	A	406	-	-	1/1/1/1	-
2	EDO	A	401	-	-	0/1/1/1	-
2	EDO	A	405	-	-	0/1/1/1	-
3	FLC	A	404	-	-	9/16/16/16	-
2	EDO	A	402	-	-	0/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	404	FLC	OA2-CAC	-2.80	1.21	1.30
3	A	404	FLC	OA1-CAC	2.68	1.30	1.22

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	404	FLC	OA1-CAC-CA	-3.54	112.91	122.95
3	A	404	FLC	OA2-CAC-CA	2.87	123.44	114.35

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	404	FLC	CAC-CA-CB-OHB
3	A	404	FLC	CG-CB-CBC-OB1
3	A	404	FLC	CG-CB-CBC-OB2
3	A	404	FLC	OHB-CB-CBC-OB1
3	A	404	FLC	OHB-CB-CBC-OB2
3	A	404	FLC	CAC-CA-CB-CG
3	A	404	FLC	CAC-CA-CB-CBC
2	A	406	EDO	O1-C1-C2-O2
3	A	404	FLC	CB-CA-CAC-OA2
3	A	404	FLC	CB-CA-CAC-OA1

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	403	EDO	2	0
2	A	406	EDO	1	0
3	A	404	FLC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	368/379 (97%)	1.19	65 (17%) 4 3	43, 65, 112, 148	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	372	THR	11.4
1	A	373	ALA	9.9
1	A	371	THR	9.3
1	A	9	VAL	7.9
1	A	252	VAL	7.1
1	A	7	ASN	6.2
1	A	6	GLU	6.1
1	A	250	HIS	5.8
1	A	295	ASP	5.8
1	A	158	PRO	5.7
1	A	255	ALA	5.5
1	A	212	GLY	5.4
1	A	253	TYR	5.3
1	A	157	GLU	5.2
1	A	211	HIS	5.2
1	A	368	ARG	5.1
1	A	8	PHE	5.0
1	A	254	ARG	4.9
1	A	370	LEU	4.7
1	A	367	GLU	4.4
1	A	226	GLU	4.1
1	A	251	ARG	4.1
1	A	227	ARG	4.0
1	A	13	ASP	3.9
1	A	296	ARG	3.9
1	A	244	LYS	3.7
1	A	347	ARG	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	28	ASP	3.6
1	A	293	ARG	3.5
1	A	11	GLY	3.5
1	A	63	SER	3.4
1	A	294	PRO	3.3
1	A	159	ASP	3.3
1	A	14	GLY	3.2
1	A	256	GLU	3.1
1	A	261	ARG	3.0
1	A	179	ASN	3.0
1	A	297	ALA	3.0
1	A	291	GLU	2.9
1	A	143	SER	2.8
1	A	314	ARG	2.7
1	A	242	GLY	2.7
1	A	228	ALA	2.7
1	A	240	ASP	2.6
1	A	229	GLY	2.6
1	A	319	MET	2.6
1	A	178	MET	2.5
1	A	230	ASP	2.4
1	A	144	SER	2.4
1	A	62	GLY	2.4
1	A	10	PRO	2.3
1	A	27	LYS	2.3
1	A	292	ARG	2.3
1	A	224	GLU	2.3
1	A	44	SER	2.2
1	A	217	ARG	2.2
1	A	369	VAL	2.2
1	A	155	GLN	2.1
1	A	110	GLN	2.1
1	A	30	GLY	2.1
1	A	213	GLY	2.1
1	A	46	ARG	2.1
1	A	45	GLN	2.1
1	A	210	LEU	2.0
1	A	288	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FLC	A	404	13/13	0.49	0.51	15,16,19,21	13
2	EDO	A	402	4/4	0.66	0.47	20,20,20,20	4
2	EDO	A	405	4/4	0.74	0.40	20,20,20,20	4
2	EDO	A	403	4/4	0.78	0.35	20,20,20,20	4
2	EDO	A	406	4/4	0.84	0.25	20,20,20,20	4
2	EDO	A	401	4/4	0.95	0.22	20,20,20,20	4

6.5 Other polymers [i](#)

There are no such residues in this entry.