



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 12:29 AM UTC

PDB ID : 3SBF / pdb_00003sbf
Title : Crystal structure of the mutant P311A of enolase superfamily member from VIBRIONALES BACTERIUM complexed with Mg and D-Arabinonate
Authors : Fedorov, A.A.; Fedorov, E.V.; Wichelecki, D.; Gerlt, J.A.; Almo, S.C.
Deposited on : 2011-06-04
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	3% 90% 8% ..
1	B	401	% 91% 6% .
1	C	401	2% 91% 8% .
1	D	401	5% 91% 8% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 14132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called mandelate racemase / muconate lactonizing enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	397	Total	C	N	O	S	0	1	0
			3172	2017	549	589	17			
1	B	390	Total	C	N	O	S	0	0	0
			3112	1978	541	576	17			
1	C	397	Total	C	N	O	S	0	1	0
			3174	2018	551	588	17			
1	D	399	Total	C	N	O	S	0	2	0
			3198	2034	554	593	17			


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP A5KUH4
A	2	SER	-	expression tag	UNP A5KUH4
A	3	LEU	-	expression tag	UNP A5KUH4
A	311	ALA	PRO	engineered mutation	UNP A5KUH4
B	1	MET	-	expression tag	UNP A5KUH4
B	2	SER	-	expression tag	UNP A5KUH4
B	3	LEU	-	expression tag	UNP A5KUH4
B	311	ALA	PRO	engineered mutation	UNP A5KUH4
C	1	MET	-	expression tag	UNP A5KUH4
C	2	SER	-	expression tag	UNP A5KUH4
C	3	LEU	-	expression tag	UNP A5KUH4
C	311	ALA	PRO	engineered mutation	UNP A5KUH4
D	1	MET	-	expression tag	UNP A5KUH4
D	2	SER	-	expression tag	UNP A5KUH4
D	3	LEU	-	expression tag	UNP A5KUH4
D	311	ALA	PRO	engineered mutation	UNP A5KUH4

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	2	Total Mg 2 2	0	0
2	C	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O S 15 8 2 4 1	0	0
3	C	1	Total C N O S 15 8 2 4 1	0	0

- Molecule 4 is D-arabinonic acid (CCD ID: D8T) (formula: $C_5H_{10}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	5	6		
4	B	1	Total	C	O	0	0
			11	5	6		
4	C	1	Total	C	O	0	0
			11	5	6		
4	D	1	Total	C	O	0	0
			11	5	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	359	Total	O	0	0
			359	359		
5	B	347	Total	O	0	0
			347	347		
5	C	361	Total	O	0	0
			361	361		
5	D	330	Total	O	0	0
			330	330		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: mandelate racemase / muconate lactonizing enzyme




- Molecule 1: mandelate racemase / muconate lactonizing enzyme




- Molecule 1: mandelate racemase / muconate lactonizing enzyme



- Molecule 1: mandelate racemase / muconate lactonizing enzyme





4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	163.37Å 163.37Å 110.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.79 – 1.50 46.79 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.3 (46.79-1.50) 98.3 (46.79-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.01 (at 1.50Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.180 , 0.206 0.176 , 0.202	Depositor DCC
R_{free} test set	11638 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 27.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	14132	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EPE, D8T

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/3257	0.74	0/4431
1	B	0.43	0/3193	0.71	0/4340
1	C	0.41	0/3259	0.73	2/4433 (0.0%)
1	D	0.40	0/3283	0.71	0/4465
All	All	0.42	0/12992	0.72	2/17669 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	17	LYS	CA-C-N	5.16	124.78	119.05
1	C	17	LYS	C-N-CA	5.16	124.78	119.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3172	0	3068	22	0
1	B	3112	0	3022	14	0
1	C	3174	0	3073	19	0
1	D	3198	0	3099	20	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	2	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	15	0	17	1	0
3	C	15	0	17	1	0
4	A	11	0	8	0	0
4	B	11	0	8	1	0
4	C	11	0	8	0	0
4	D	11	0	8	0	0
5	A	359	0	0	4	0
5	B	347	0	0	2	0
5	C	361	0	0	7	0
5	D	330	0	0	2	0
All	All	14132	0	12328	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:21:HIS:HB3	1:D:43:GLN:HG3	1.69	0.73
1:B:60:ILE:O	1:B:64:LYS:HE2	1.92	0.69
3:C:403:EPE:H22	5:C:1335:HOH:O	1.96	0.66
1:C:240:GLN:HG3	5:C:510:HOH:O	1.98	0.63
1:B:246:ASN:CG	5:B:1365:HOH:O	2.44	0.60
1:A:34:VAL:HG21	1:A:106:LEU:HD21	1.84	0.59
1:B:19:ASP:OD2	1:B:20:ARG:HG2	2.04	0.58
1:A:163:PRO:HB3	1:A:177:TYR:CG	2.40	0.57
1:D:163:PRO:HB2	1:D:168:THR:HG21	1.87	0.57
1:B:64:LYS:HD3	5:B:759:HOH:O	2.05	0.56
1:B:372:ASP:OD2	1:B:374:GLU:HG2	2.08	0.54
1:C:240:GLN:HG3	5:C:458:HOH:O	2.07	0.54
1:D:311:ALA:HB1	1:D:312:PRO:CD	2.38	0.53
1:D:218:ILE:HD11	1:D:250:GLN:NE2	2.24	0.52
1:A:244:LEU:O	1:A:248:ARG:HG2	2.09	0.52
1:C:127:THR:HG21	1:C:144:PHE:CG	2.44	0.52
1:D:20:ARG:HD2	1:D:384:ARG:NH1	2.24	0.52
1:C:300:GLN:NE2	5:C:1414:HOH:O	2.40	0.52
1:A:60:ILE:O	1:A:64:LYS:HE2	2.09	0.51
1:A:124:PRO:HG3	5:A:696:HOH:O	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:ARG:NH1	3:A:403:EPE:O2S	2.44	0.50
1:B:340:TYR:CD1	1:B:344:THR:HG21	2.47	0.50
1:A:73:TRP:CH2	1:A:77:MET:HG3	2.48	0.49
1:B:20:ARG:HA	1:B:20:ARG:HH11	1.78	0.49
1:A:24:ILE:HD11	1:A:46:PRO:HA	1.95	0.48
1:C:138:TYR:CE1	1:C:197:LYS:HD3	2.48	0.48
1:C:340:TYR:CD1	1:C:344:THR:HG21	2.47	0.48
1:B:24:ILE:HD11	1:B:46:PRO:HA	1.96	0.48
1:D:311:ALA:HB1	1:D:312:PRO:HD2	1.96	0.48
1:C:339:GLU:OE2	1:C:339:GLU:HA	2.13	0.48
1:C:16:THR:C	1:C:18:PRO:HD3	2.39	0.47
1:A:274:ARG:HD3	5:A:536:HOH:O	2.15	0.47
1:B:58:LYS:HB3	1:B:59:PRO:HD3	1.96	0.46
1:A:314:MET:HG3	5:A:428:HOH:O	2.15	0.46
1:D:346:LYS:NZ	5:D:986:HOH:O	2.44	0.46
1:B:209:HIS:NE2	4:B:404:D8T:HAG	2.31	0.46
1:B:310:CYS:HA	1:B:314:MET:HE1	1.98	0.46
1:D:134:MET:SD	1:D:156:LEU:HD22	2.56	0.46
1:B:306:ILE:HB	1:B:332:ALA:HA	1.97	0.45
1:A:43:GLN:O	1:A:46:PRO:HD3	2.16	0.45
1:A:41:THR:HG21	1:A:313:ASP:O	2.17	0.45
1:D:58:LYS:HB3	1:D:59:PRO:HD3	1.98	0.44
1:C:146:GLU:HG3	5:C:1184:HOH:O	2.17	0.43
1:D:306:ILE:HB	1:D:332:ALA:HA	2.00	0.43
1:D:163:PRO:CB	1:D:168:THR:HG21	2.47	0.43
1:D:24:ILE:HD11	1:D:46:PRO:HA	2.00	0.43
1:A:21:HIS:CE1	1:A:312:PRO:HB2	2.53	0.43
1:A:374:GLU:HG2	5:A:1393:HOH:O	2.17	0.43
1:A:73:TRP:CZ2	1:A:77:MET:HG3	2.54	0.43
1:A:207:ASP:HA	1:A:233:GLU:HB3	2.01	0.42
1:C:21:HIS:CE1	1:C:312:PRO:HB2	2.54	0.42
1:A:306:ILE:HB	1:A:332:ALA:HA	2.01	0.42
1:C:24:ILE:HD11	1:C:46:PRO:HA	2.01	0.42
1:C:138:TYR:CZ	1:C:197:LYS:HD3	2.54	0.42
1:D:60:ILE:O	1:D:64:LYS:HE2	2.19	0.42
1:D:210:GLU:CD	1:D:400:THR:HB	2.44	0.42
1:D:244:LEU:O	1:D:248:ARG:HG2	2.19	0.42
1:D:4:LYS:HD2	5:D:903:HOH:O	2.19	0.42
1:A:14:ILE:HD11	1:A:373:ARG:HD3	2.01	0.42
1:C:153:ARG:CZ	5:C:1201:HOH:O	2.68	0.42
1:D:17:LYS:HD3	1:D:382:MET:HB3	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:341:ASN:OD1	1:D:341:ASN:C	2.63	0.42
1:C:306:ILE:HB	1:C:332:ALA:HA	2.01	0.42
1:B:207:ASP:HA	1:B:233:GLU:HB3	2.00	0.41
1:A:165:ASP:OD1	1:A:165:ASP:N	2.54	0.41
1:C:274:ARG:HD3	5:C:522:HOH:O	2.19	0.41
1:C:310:CYS:HB2	1:C:336:GLU:HB3	2.03	0.41
1:A:314:MET:HE2	1:A:314:MET:HB3	1.76	0.41
1:C:104:ALA:HB3	1:C:366:GLY:HA2	2.03	0.41
1:D:104:ALA:HB3	1:D:366:GLY:HA2	2.02	0.41
1:A:104:ALA:HB3	1:A:366:GLY:HA2	2.03	0.41
1:B:244:LEU:O	1:B:248:ARG:HG2	2.21	0.41
1:C:73:TRP:CH2	1:C:77:MET:HG3	2.56	0.41
1:C:43:GLN:HG3	1:C:313:ASP:HB2	2.03	0.40
1:A:183:TYR:OH	1:A:208:VAL:HA	2.21	0.40
1:D:203:HIS:HB3	1:D:230:TYR:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/401 (99%)	389 (98%)	7 (2%)	0	100	100
1	B	386/401 (96%)	380 (98%)	6 (2%)	0	100	100
1	C	396/401 (99%)	389 (98%)	7 (2%)	0	100	100
1	D	399/401 (100%)	390 (98%)	9 (2%)	0	100	100
All	All	1577/1604 (98%)	1548 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/344 (99%)	336 (98%)	5 (2%)	57	31
1	B	335/344 (97%)	331 (99%)	4 (1%)	63	38
1	C	341/344 (99%)	335 (98%)	6 (2%)	51	24
1	D	344/344 (100%)	340 (99%)	4 (1%)	63	38
All	All	1361/1376 (99%)	1342 (99%)	19 (1%)	59	33

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	ARG
1	A	165	ASP
1	A	166	LEU
1	A	257	LEU
1	A	374	GLU
1	B	4	LYS
1	B	20	ARG
1	B	166	LEU
1	B	257	LEU
1	C	31	ASN
1	C	43	GLN
1	C	162	VAL
1	C	201	GLN
1	C	257	LEU
1	C	387	GLU
1	D	20	ARG
1	D	43	GLN
1	D	139	ASP
1	D	166	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	HIS
1	A	201	GLN
1	A	226	GLN
1	B	201	GLN
1	B	219	GLN
1	B	226	GLN
1	B	240	GLN
1	B	273	ASN
1	C	21	HIS
1	C	89	ASN
1	C	90	ASN
1	C	240	GLN
1	C	273	ASN
1	C	300	GLN
1	C	390	GLN
1	D	201	GLN
1	D	273	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	D8T	A	404	2	10,10,10	0.76	0	12,13,13	0.98	1 (8%)
3	EPE	C	403	-	15,15,15	0.93	1 (6%)	19,20,20	1.62	6 (31%)
4	D8T	D	403	2	10,10,10	0.78	0	12,13,13	1.05	2 (16%)
4	D8T	C	404	2	10,10,10	0.79	0	12,13,13	1.15	2 (16%)
3	EPE	A	403	-	15,15,15	0.79	1 (6%)	19,20,20	1.80	4 (21%)
4	D8T	B	404	2	10,10,10	0.79	0	12,13,13	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	D8T	A	404	2	-	0/14/14/14	-
3	EPE	C	403	-	-	3/9/19/19	0/1/1/1
4	D8T	D	403	2	-	2/14/14/14	-
4	D8T	C	404	2	-	2/14/14/14	-
3	EPE	A	403	-	-	6/9/19/19	0/1/1/1
4	D8T	B	404	2	-	2/14/14/14	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	403	EPE	C10-S	3.21	1.82	1.77
3	A	403	EPE	C10-S	2.76	1.81	1.77

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	EPE	C5-N4-C3	5.12	119.87	108.84
3	C	403	EPE	C5-N4-C3	3.81	117.05	108.84
3	A	403	EPE	C3-C2-N1	-2.68	105.25	110.65
3	C	403	EPE	C7-N4-C5	2.58	118.12	111.24
4	C	404	D8T	OAJ-CAD-CAE	2.27	119.61	113.31
4	D	403	D8T	OAJ-CAD-CAE	2.25	119.55	113.31
3	C	403	EPE	C7-N4-C3	2.23	117.18	111.24
3	C	403	EPE	O2S-S-C10	2.21	110.07	106.73
3	C	403	EPE	O3S-S-C10	2.20	110.31	106.00
3	A	403	EPE	O3S-S-C10	2.18	110.27	106.00
3	A	403	EPE	C7-N4-C5	2.18	117.05	111.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	403	D8T	OAA-CAD-CAE	-2.11	116.01	121.62
3	C	403	EPE	C6-N1-C2	2.09	113.34	108.84
4	C	404	D8T	OAA-CAD-CAE	-2.01	116.25	121.62
4	A	404	D8T	OAJ-CAD-CAE	2.00	118.88	113.31

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	EPE	C10-C9-N1-C6
3	C	403	EPE	C10-C9-N1-C6
3	C	403	EPE	C8-C7-N4-C3
3	A	403	EPE	C8-C7-N4-C5
3	A	403	EPE	C10-C9-N1-C2
3	C	403	EPE	C10-C9-N1-C2
3	A	403	EPE	C9-C10-S-O3S
3	A	403	EPE	C9-C10-S-O1S
3	A	403	EPE	C9-C10-S-O2S
4	D	403	D8T	OAA-CAD-CAE-CAF
4	D	403	D8T	OAJ-CAD-CAE-CAF
4	C	404	D8T	OAA-CAD-CAE-CAF
4	B	404	D8T	OAA-CAD-CAE-CAF
4	C	404	D8T	OAJ-CAD-CAE-CAF
4	B	404	D8T	OAJ-CAD-CAE-CAF

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	403	EPE	1	0
3	A	403	EPE	1	0
4	B	404	D8T	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	397/401 (99%)	-0.02	12 (3%) 52 56	9, 19, 32, 59	1 (0%)
1	B	390/401 (97%)	-0.06	5 (1%) 75 78	15, 19, 31, 44	0
1	C	397/401 (99%)	-0.04	8 (2%) 65 68	9, 19, 32, 49	1 (0%)
1	D	399/401 (99%)	0.12	21 (5%) 32 35	10, 20, 36, 51	2 (0%)
All	All	1583/1604 (98%)	-0.00	46 (2%) 53 58	9, 19, 34, 59	4 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	338	VAL	5.1
1	A	34	VAL	4.7
1	A	33	GLY	4.4
1	A	164	THR	4.3
1	B	166	LEU	4.1
1	D	164	THR	3.9
1	D	158	PHE	3.7
1	B	20	ARG	3.6
1	B	157	GLY	3.5
1	A	311	ALA	3.4
1	A	163	PRO	3.3
1	D	3	LEU	3.3
1	A	337	HIS	3.2
1	D	20	ARG	3.2
1	D	159	TYR	3.2
1	D	375	ALA	3.2
1	D	387	GLU	3.1
1	A	158	PHE	3.1
1	C	338	VAL	3.0
1	D	162	VAL	2.9
1	D	373	ARG	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	160	GLY	2.9
1	B	340	TYR	2.8
1	A	162	VAL	2.7
1	D	82	TRP	2.7
1	C	162	VAL	2.7
1	D	374	GLU	2.5
1	D	163	PRO	2.5
1	D	135	GLU	2.4
1	D	133	THR	2.4
1	C	341	ASN	2.4
1	C	201	GLN	2.4
1	C	340	TYR	2.3
1	C	337	HIS	2.3
1	A	166	LEU	2.3
1	C	387	GLU	2.3
1	D	340	TYR	2.2
1	D	377	ALA	2.2
1	D	378	GLU	2.2
1	D	166	LEU	2.1
1	C	164	THR	2.1
1	A	165	ASP	2.1
1	D	346	LYS	2.1
1	B	341	ASN	2.1
1	D	132	ASP	2.0
1	A	31	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EPE	C	403	15/15	0.83	0.13	27,35,52,53	0
3	EPE	A	403	15/15	0.85	0.12	27,30,51,60	0
4	D8T	D	403	11/11	0.85	0.11	22,32,39,39	0
4	D8T	B	404	11/11	0.91	0.09	19,26,33,36	0
4	D8T	C	404	11/11	0.94	0.07	19,24,29,32	0
4	D8T	A	404	11/11	0.94	0.07	18,21,26,28	0
2	MG	C	402	1/1	0.99	0.03	17,17,17,17	0
2	MG	D	402	1/1	0.99	0.03	19,19,19,19	0
2	MG	A	402	1/1	0.99	0.03	17,17,17,17	0
2	MG	B	403	1/1	0.99	0.11	24,24,24,24	0
2	MG	B	402	1/1	1.00	0.04	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.