



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:57 AM UTC

PDB ID : 8SBF / pdb_00008sbf
Title : Full-length structure of the LysR-type transcriptional regulator, ACIAD0746, from *Acinetobacter baylyi*
Authors : Momany, C.; Nune, M.; Brondani, J.C.; Afful, D.; Neidle, E.; Galloway, N.R.
Deposited on : 2023-04-03
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	 89% 9% .
1	B	300	 85% 13% ..
1	C	300	 87% 11% .
1	D	300	 78% 19% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 19224 atoms, of which 9401 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative transcriptional regulator (LysR family).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	294	Total	C	H	N	O	S	0	3	0
			4655	1461	2349	410	421	14			
1	C	295	Total	C	H	N	O	S	0	1	0
			4646	1459	2343	409	421	14			
1	B	295	Total	C	H	N	O	S	0	1	0
			4656	1461	2350	411	420	14			
1	D	293	Total	C	H	N	O	S	0	1	0
			4626	1452	2335	408	417	14			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	296	HIS	-	expression tag	UNP Q6FE56
A	297	HIS	-	expression tag	UNP Q6FE56
A	298	HIS	-	expression tag	UNP Q6FE56
A	299	HIS	-	expression tag	UNP Q6FE56
A	300	HIS	-	expression tag	UNP Q6FE56
C	296	HIS	-	expression tag	UNP Q6FE56
C	297	HIS	-	expression tag	UNP Q6FE56
C	298	HIS	-	expression tag	UNP Q6FE56
C	299	HIS	-	expression tag	UNP Q6FE56
C	300	HIS	-	expression tag	UNP Q6FE56
B	296	HIS	-	expression tag	UNP Q6FE56
B	297	HIS	-	expression tag	UNP Q6FE56
B	298	HIS	-	expression tag	UNP Q6FE56
B	299	HIS	-	expression tag	UNP Q6FE56
B	300	HIS	-	expression tag	UNP Q6FE56
D	296	HIS	-	expression tag	UNP Q6FE56
D	297	HIS	-	expression tag	UNP Q6FE56
D	298	HIS	-	expression tag	UNP Q6FE56
D	299	HIS	-	expression tag	UNP Q6FE56
D	300	HIS	-	expression tag	UNP Q6FE56


- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	C	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	2	Total	Mg	0	0
			2	2		
3	B	2	Total	Mg	0	0
			2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	159	Total 159	O 159	0	0
5	C	184	Total 187	O 187	0	3
5	B	136	Total 138	O 138	0	2
5	D	105	Total 108	O 108	0	3


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative transcriptional regulator (LysR family)




- Molecule 1: Putative transcriptional regulator (LysR family)




- Molecule 1: Putative transcriptional regulator (LysR family)



- Molecule 1: Putative transcriptional regulator (LysR family)





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.60Å 109.69Å 110.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.45 – 2.29 26.45 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.7 (26.45-2.29) 95.0 (26.45-2.29)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.56 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.211 , 0.258 0.212 , 0.258	Depositor DCC
R_{free} test set	2912 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	45.0	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for -h,l,k 0.015 for -l,-k,-h 0.015 for k,h,-l 0.004 for k,l,h 0.004 for l,h,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	19224	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/2356	0.26	0/3193
1	B	0.10	0/2348	0.28	0/3182
1	C	0.10	0/2345	0.26	0/3179
1	D	0.11	0/2333	0.28	0/3161
All	All	0.10	0/9382	0.27	0/12715

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2306	2349	2337	14	0
1	B	2306	2350	2350	34	0
1	C	2303	2343	2343	20	0
1	D	2291	2335	2335	36	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	18	24	24	5	0
5	A	159	0	0	0	0
5	B	138	0	0	0	0
5	C	187	0	0	0	0
5	D	108	0	0	1	0
All	All	9823	9401	9389	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:165:LEU:HD11	1:D:204:VAL:HG11	1.49	0.94
1:B:188:LEU:HD13	1:B:213:VAL:HG13	1.70	0.72
1:B:133:ILE:HG21	1:B:152:LEU:HD13	1.71	0.71
1:B:193:ILE:HD12	1:B:220:LEU:HD12	1.72	0.70
1:D:165:LEU:HD12	1:D:265:LEU:HB3	1.74	0.70
1:C:66:LEU:HD11	1:B:84:ALA:HB2	1.74	0.69
1:B:110:MET:HE1	1:B:124:LEU:HB2	1.75	0.69
1:A:175:LEU:HD12	1:A:238:MET:HE1	1.76	0.68
1:D:185:GLN:O	1:D:188:LEU:HD22	1.94	0.68
1:B:134:GLU:OE2	1:B:138:GLN:NE2	2.27	0.67
1:B:211:GLN:OE1	1:B:211:GLN:N	2.28	0.67
1:D:238:MET:HE2	1:D:258:LEU:HD11	1.77	0.66
1:A:143:LEU:HD23	1:A:144:ALA:N	2.10	0.66
1:B:187:PHE:CE2	1:B:238:MET:HE1	2.31	0.65
1:D:23:ALA:O	1:D:27:VAL:HG22	1.97	0.65
1:C:45:LEU:HD22	1:B:84:ALA:HB3	1.78	0.65
1:B:190:VAL:HG11	1:B:238:MET:HE2	1.78	0.65
1:B:187:PHE:CD2	1:B:238:MET:HE1	2.33	0.62
1:D:165:LEU:HD11	1:D:204:VAL:CG1	2.28	0.62
1:D:201:THR:HG21	1:D:241:ILE:HD12	1.81	0.62
1:C:52:ARG:O	1:C:53:LEU:HD23	2.01	0.59
1:A:59:LEU:HD11	1:A:67:LEU:HD22	1.86	0.57
1:C:1:MET:HE2	1:C:3:MET:HB3	1.86	0.57
1:B:133:ILE:CG2	1:B:152:LEU:HD13	2.35	0.57
1:D:152:LEU:HD23	1:D:154:GLN:NE2	2.21	0.56
1:C:155:LEU:HD13	1:C:272:LEU:HD11	1.88	0.55
1:D:193:ILE:HD11	1:D:236:LEU:HD12	1.89	0.54
1:D:49:LEU:HD23	1:D:62:LEU:HD23	1.88	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:30:THR:HG21	4:B:402:GOL:H32	1.90	0.54
1:C:136:VAL:HG11	1:C:272:LEU:HD13	1.90	0.54
1:D:165:LEU:HD13	1:D:240:CYS:SG	2.48	0.53
1:B:246:ILE:HG22	1:B:250:ILE:HG12	1.91	0.53
1:B:246:ILE:HG22	1:B:250:ILE:CG1	2.39	0.53
1:D:157:CYS:SG	1:D:270:HIS:ND1	2.64	0.52
1:C:45:LEU:HD22	1:B:84:ALA:CB	2.40	0.51
1:B:78:GLU:HG3	4:B:403:GOL:H31	1.91	0.51
1:D:260:THR:HG21	1:D:263:TRP:HE3	1.76	0.51
1:B:59:LEU:HD11	1:B:67:LEU:HD22	1.92	0.51
1:D:3:MET:HE3	1:D:8:LEU:HD21	1.93	0.51
1:D:52:ARG:O	1:D:53:LEU:HD22	2.10	0.51
1:B:219:ARG:NH2	1:D:118:PRO:O	2.43	0.51
1:A:118:PRO:O	1:C:219:ARG:NH2	2.41	0.50
1:D:238:MET:CE	1:D:258:LEU:HD11	2.40	0.50
1:A:153:PRO:HB3	1:B:218:ILE:HD12	1.94	0.49
1:D:241:ILE:HG22	1:D:246:ILE:HD11	1.93	0.49
1:C:28:ARG:NE	1:C:28:ARG:O	2.45	0.49
1:B:121:HIS:C	1:B:122:LEU:HD23	2.38	0.49
1:D:36:MET:HE2	1:D:40:ASP:OD2	2.13	0.48
1:D:260:THR:HG21	1:D:263:TRP:CE3	2.48	0.48
1:C:104:TYR:OH	1:C:226:GLU:OE2	2.25	0.48
1:D:279:GLN:HG2	1:D:283:LEU:HD23	1.96	0.48
1:A:119:GLN:OE1	1:A:119:GLN:N	2.35	0.48
1:B:95:LEU:C	1:B:141:LEU:HD22	2.39	0.48
1:B:135:ALA:CB	1:B:141:LEU:HD12	2.45	0.47
1:B:135:ALA:HB3	1:B:141:LEU:HD12	1.97	0.47
1:B:193:ILE:HD11	4:B:401:GOL:H11	1.96	0.47
1:C:84:ALA:HB3	1:B:45:LEU:HD22	1.97	0.47
1:A:250:ILE:HD11	1:A:257:ILE:HG13	1.97	0.46
1:B:222:LEU:HD22	1:D:110:MET:HE1	1.98	0.46
1:B:183:THR:HG22	1:B:184:ALA:N	2.31	0.46
1:D:165:LEU:HB3	1:D:242:SER:HA	1.97	0.46
1:A:67:LEU:HD23	1:A:139:PHE:CD2	2.51	0.46
1:A:183:THR:OG1	1:A:185:GLN:OE1	2.34	0.45
1:A:94:ARG:NH1	1:A:140:GLN:O	2.49	0.45
1:B:89:GLN:HG3	1:B:90:SER:N	2.32	0.45
1:D:12:VAL:HG21	1:D:74:LEU:HD12	1.99	0.45
1:A:169:CYS:HB3	1:A:238:MET:HG2	1.99	0.45
1:D:2:ARG:H	1:D:2:ARG:HD2	1.81	0.45
1:C:23:ALA:O	1:C:27:VAL:HG22	2.16	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:221:THR:O	1:D:222:LEU:HD23	2.17	0.44
1:C:1:MET:HE3	1:C:44:GLY:HA3	1.99	0.44
1:D:133:ILE:HD13	1:D:272:LEU:HD21	2.00	0.44
1:D:147:GLU:HG3	1:D:200:GLY:HA3	2.00	0.44
1:D:183:THR:HG23	1:D:186:ASP:H	1.82	0.43
1:D:188:LEU:HD23	1:D:189:ASN:H	1.83	0.43
1:D:221:THR:C	1:D:222:LEU:HD23	2.43	0.43
1:C:194:LEU:HD12	1:C:218:ILE:HD12	2.00	0.43
1:A:98:THR:HA	1:A:127:GLY:O	2.19	0.43
1:D:82:GLN:HG3	1:D:83:ALA:N	2.33	0.42
1:D:165:LEU:HD21	1:D:204:VAL:HG21	2.02	0.42
1:A:164:GLU:OE2	1:A:266:THR:HG22	2.19	0.42
1:D:155:LEU:HD23	1:D:272:LEU:HD11	2.02	0.42
1:B:30:THR:HG21	4:B:402:GOL:H11	2.01	0.41
1:D:277:LYS:NZ	5:D:420:HOH:O	2.53	0.41
1:C:52:ARG:HD3	1:C:57:LEU:HD21	2.02	0.41
1:C:169:CYS:HB3	1:C:238:MET:HG2	2.02	0.41
1:B:138:GLN:O	1:B:139:PHE:HB2	2.21	0.41
1:C:250:ILE:HD11	1:C:257:ILE:HG13	2.02	0.41
1:A:152:LEU:HD23	1:A:155:LEU:HG	2.02	0.41
1:B:172:ASP:OD1	1:B:172:ASP:N	2.45	0.41
1:C:147:GLU:O	1:C:267:ARG:HD2	2.20	0.40
1:B:163:ASP:OD2	1:B:267:ARG:NH1	2.54	0.40
1:C:2:ARG:N	1:C:2:ARG:HD3	2.35	0.40
1:C:145:LEU:HD12	1:C:145:LEU:N	2.36	0.40
1:B:30:THR:CB	4:B:402:GOL:H32	2.51	0.40
1:D:145:LEU:N	1:D:145:LEU:HD12	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/300 (98%)	285 (97%)	10 (3%)	0	100	100
1	B	294/300 (98%)	284 (97%)	9 (3%)	1 (0%)	36	46
1	C	294/300 (98%)	282 (96%)	12 (4%)	0	100	100
1	D	292/300 (97%)	280 (96%)	10 (3%)	2 (1%)	18	23
All	All	1175/1200 (98%)	1131 (96%)	41 (4%)	3 (0%)	36	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	89	GLN
1	D	54	GLY
1	D	214	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/251 (99%)	244 (98%)	4 (2%)	55	73
1	B	247/251 (98%)	242 (98%)	5 (2%)	48	67
1	C	247/251 (98%)	241 (98%)	6 (2%)	43	62
1	D	245/251 (98%)	236 (96%)	9 (4%)	30	45
All	All	987/1004 (98%)	963 (98%)	24 (2%)	43	62

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	ARG
1	A	177	LYS
1	A	180	ARG
1	A	188	LEU
1	C	56	ARG
1	C	149	SER
1	C	157	CYS
1	C	188	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	230	LYS
1	C	272	LEU
1	B	88	LEU
1	B	143	LEU
1	B	152	LEU
1	B	241	ILE
1	B	267	ARG
1	D	25	ASP
1	D	29	LEU
1	D	31	GLN
1	D	55	LYS
1	D	108	ASP
1	D	165	LEU
1	D	180	ARG
1	D	215	ASP
1	D	241	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	HIS
1	A	138	GLN
1	A	211	GLN
1	A	270	HIS
1	C	82	GLN
1	B	159	HIS
1	D	217	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 7 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	B	403	-	5,5,5	0.89	0	5,5,5	1.19	1 (20%)
4	GOL	B	401	-	5,5,5	0.93	0	5,5,5	1.15	1 (20%)
4	GOL	B	402	-	5,5,5	0.88	0	5,5,5	1.20	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	403	-	-	2/4/4/4	-
4	GOL	B	401	-	-	2/4/4/4	-
4	GOL	B	402	-	-	2/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	402	GOL	C3-C2-C1	-2.18	103.80	111.80
4	B	403	GOL	C3-C2-C1	-2.10	104.08	111.80
4	B	401	GOL	C3-C2-C1	-2.08	104.19	111.80

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	402	GOL	O1-C1-C2-C3
4	B	403	GOL	O1-C1-C2-C3
4	B	401	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	403	GOL	O1-C1-C2-O2
4	B	402	GOL	O1-C1-C2-O2
4	B	401	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	403	GOL	1	0
4	B	401	GOL	1	0
4	B	402	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/300 (98%)	0.03	4 (1%) 73 75	35, 62, 96, 123	2 (0%)
1	B	295/300 (98%)	0.29	3 (1%) 79 80	26, 70, 111, 134	1 (0%)
1	C	295/300 (98%)	0.06	4 (1%) 73 75	35, 62, 96, 115	1 (0%)
1	D	293/300 (97%)	0.67	20 (6%) 23 25	34, 96, 174, 203	1 (0%)
All	All	1177/1200 (98%)	0.26	31 (2%) 57 59	26, 69, 130, 203	5 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	182	LEU	3.3
1	D	171	PRO	3.2
1	D	214	PRO	3.2
1	D	179	GLY	3.0
1	D	1	MET	3.0
1	A	294	VAL	3.0
1	D	190	VAL	2.8
1	D	255	LEU	2.8
1	D	253	GLY	2.7
1	D	192	TRP	2.7
1	A	266	THR	2.7
1	A	88	LEU	2.6
1	D	148	GLY	2.6
1	C	56	ARG	2.4
1	B	198	GLY	2.4
1	B	209	ILE	2.4
1	D	213	VAL	2.4
1	D	210	LEU	2.3
1	B	54	GLY	2.3
1	C	128	ASN	2.2
1	D	178	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	263	TRP	2.2
1	D	136	VAL	2.2
1	D	55	LYS	2.2
1	C	53	LEU	2.1
1	C	295	ASN	2.1
1	A	219[A]	ARG	2.1
1	D	215	ASP	2.1
1	D	233	ALA	2.0
1	D	256	VAL	2.0
1	D	264	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	402	6/6	0.68	0.20	95,133,157,169	0
3	MG	B	404	1/1	0.75	0.22	85,85,85,85	0
4	GOL	B	401	6/6	0.81	0.15	103,133,160,160	0
3	MG	C	402	1/1	0.85	0.18	90,90,90,90	0
4	GOL	B	403	6/6	0.85	0.19	99,135,156,163	0
3	MG	C	401	1/1	0.89	0.10	69,69,69,69	0
2	NA	B	406	1/1	0.90	0.14	74,74,74,74	0
2	NA	A	401	1/1	0.91	0.10	76,76,76,76	0
3	MG	B	405	1/1	0.94	0.07	58,58,58,58	0
2	NA	C	403	1/1	0.94	0.18	59,59,59,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.