



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 09:28 AM UTC

PDB ID : 3SDK / pdb\_00003sdk  
Title : Structure of yeast 20S open-gate proteasome with Compound 34  
Authors : Sintchak, M.D.  
Deposited on : 2011-06-09  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	
1	O	250	
2	B	235	
2	P	235	
3	C	241	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	241	<p>24% 78% 20%</p>
4	D	260	<p>5% 77% 12% 10%</p>
4	R	260	<p>5% 71% 17% 11%</p>
5	E	233	<p>6% 78% 19%</p>
5	S	233	<p>12% 76% 22%</p>
6	F	242	<p>3% 78% 16%</p>
6	T	242	<p>2% 83% 13%</p>
7	G	243	<p>3% 79% 19%</p>
7	U	243	<p>4% 81% 16%</p>
8	H	222	<p>1% 83% 17%</p>
8	V	222	<p>3% 83% 17%</p>
9	I	204	<p>84% 15%</p>
9	W	204	<p>85% 14%</p>
10	J	198	<p>4% 81% 18%</p>
10	X	198	<p>5% 80% 17%</p>
11	K	212	<p>1% 83% 15%</p>
11	Y	212	<p>90% 8%</p>
12	L	222	<p>86% 12%</p>
12	Z	222	<p>2% 84% 13%</p>
13	1	233	<p>83% 16%</p>
13	M	233	<p>3% 85% 14%</p>
14	2	196	<p>87% 12%</p>
14	N	196	<p>86% 12%</p>

## 2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 49255 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome component Y7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	1915	1219	315	377	4	0	0	0
1	O	250	1915	1219	315	377	4	0	0	0

- Molecule 2 is a protein called Proteasome component Y13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	235	1829	1158	303	365	3	0	0	0
2	P	235	1829	1158	303	365	3	0	0	0

- Molecule 3 is a protein called Proteasome component PRE6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	241	1891	1181	331	375	4	0	0	0
3	Q	241	1891	1181	331	375	4	0	0	0

- Molecule 4 is a protein called Proteasome component PUP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	233	1794	1123	302	362	7	0	0	0
4	R	232	1786	1120	298	361	7	0	0	0

- Molecule 5 is a protein called Proteasome component PRE5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			
5	S	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			

- Molecule 6 is a protein called Proteasome component C1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	236	Total	C	N	O	S	0	0	0
			1840	1171	321	344	4			
6	T	236	Total	C	N	O	S	0	0	0
			1840	1171	321	344	4			

- Molecule 7 is a protein called Proteasome component C7-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			
7	U	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			

- Molecule 8 is a protein called Proteasome component PUP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	222	Total	C	N	O	S	0	0	0
			1685	1061	293	324	7			
8	V	222	Total	C	N	O	S	0	0	0
			1685	1061	293	324	7			

- Molecule 9 is a protein called Proteasome component PUP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome component C11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	198	Total	C	N	O	S	0	0	0
			1582	1003	269	305	5			

- Molecule 11 is a protein called Proteasome component PRE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome component C5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome component PRE4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	1	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome component PRE3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	2	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	F	2	Total	Mg	0	0
			2	2		
15	G	1	Total	Mg	0	0
			1	1		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
15	H	1	Total Mg 1 1	0	0
15	I	2	Total Mg 2 2	0	0
15	K	1	Total Mg 1 1	0	0
15	L	2	Total Mg 2 2	0	0
15	N	1	Total Mg 1 1	0	0

- Molecule 16 is N-[(2S)-3-(3-tert-butyl-1,2,4-oxadiazol-5-yl)-1-((2S)-1-[(4-methylbenzyl)amino]-1-oxo-4-phenylbutan-2-yl)amino)-1-oxopropan-2-yl]-5-methyl-1,2-oxazole-3-carboxamide (CCD ID: P3N) (formula: C<sub>32</sub>H<sub>38</sub>N<sub>6</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	K	1	Total C N O 43 32 6 5	0	0
16	Y	1	Total C N O 43 32 6 5	0	0

- Molecule 17 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
17	K	1	12	6	1	4	1	0	0
17	Y	1	12	6	1	4	1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome component Y7




- Molecule 1: Proteasome component Y7




- Molecule 2: Proteasome component Y13





- Molecule 2: Proteasome component Y13









• Molecule 3: Proteasome component PRE6



• Molecule 3: Proteasome component PRE6




• Molecule 4: Proteasome component PUP2





• Molecule 4: Proteasome component PUP2









- Molecule 5: Proteasome component PRE5




- Molecule 5: Proteasome component PRE5



- Molecule 6: Proteasome component C1





- Molecule 6: Proteasome component C1








- Molecule 7: Proteasome component C7-alpha





- Molecule 7: Proteasome component C7-alpha




- Molecule 8: Proteasome component PUP1




- Molecule 8: Proteasome component PUP1




- Molecule 9: Proteasome component PUP3




- Molecule 9: Proteasome component PUP3




- Molecule 10: Proteasome component C11




- Molecule 10: Proteasome component C11



- Molecule 11: Proteasome component PRE2









- Molecule 11: Proteasome component PRE2





- Molecule 12: Proteasome component C5





- Molecule 12: Proteasome component C5




- Molecule 13: Proteasome component PRE4





- Molecule 13: Proteasome component PRE4



## ● Molecule 14: Proteasome component PRE3

Chain N:  86% 12%

## ● Molecule 14: Proteasome component PRE3

Chain 2:  % 87% 12%

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	136.12Å 299.40Å 146.24Å 90.00° 112.92° 90.00°	Depositor
Resolution (Å)	50.00 – 2.70 50.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	95.0 (50.00-2.70) 95.1 (50.00-2.70)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.43 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.223 , 0.259 0.219 , 0.251	Depositor DCC
$R_{free}$ test set	5652 reflections (2.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.4	Xtrriage
Anisotropy	0.156	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 25.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	49255	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: P3N, MG, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.55	0/1952	0.82	0/2642
1	O	0.63	2/1952 (0.1%)	0.84	3/2642 (0.1%)
2	B	0.56	0/1858	0.85	1/2516 (0.0%)
2	P	0.56	0/1858	0.83	0/2516
3	C	0.57	0/1920	0.85	0/2598
3	Q	0.60	2/1920 (0.1%)	0.88	0/2598
4	D	0.55	0/1817	0.81	0/2449
4	R	0.52	0/1810	0.82	0/2440
5	E	0.64	3/1823 (0.2%)	0.85	2/2463 (0.1%)
5	S	0.57	2/1823 (0.1%)	0.83	4/2463 (0.2%)
6	F	0.56	1/1879 (0.1%)	0.83	0/2535
6	T	0.57	1/1879 (0.1%)	0.84	0/2535
7	G	0.61	1/1959 (0.1%)	0.85	1/2652 (0.0%)
7	U	0.58	1/1959 (0.1%)	0.85	0/2652
8	H	0.55	0/1716	0.83	0/2326
8	V	0.59	0/1716	0.81	0/2326
9	I	0.65	1/1611 (0.1%)	0.83	3/2174 (0.1%)
9	W	0.67	0/1611	0.86	1/2174 (0.0%)
10	J	0.63	0/1613	0.82	0/2173
10	X	0.58	0/1610	0.83	1/2170 (0.0%)
11	K	0.58	0/1681	0.86	2/2274 (0.1%)
11	Y	0.53	0/1681	0.81	0/2274
12	L	0.57	0/1795	0.80	0/2420
12	Z	0.59	0/1795	0.82	0/2420
13	1	0.62	0/1855	0.83	2/2514 (0.1%)
13	M	0.61	1/1855 (0.1%)	0.82	0/2514
14	2	0.68	0/1541	0.84	0/2087
14	N	0.71	1/1541 (0.1%)	0.82	0/2087
All	All	0.59	16/50030 (0.0%)	0.83	20/67634 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if

the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	O	0	1
5	E	0	2
5	S	0	1
14	N	0	1
All	All	0	5

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	207(E)	ASN	C-O	12.41	1.39	1.24
1	O	200	SER	C-O	11.93	1.39	1.24
3	Q	180(D)	GLU	C-O	9.51	1.36	1.24
14	N	181	ALA	C-O	8.14	1.34	1.24
1	O	200	SER	CA-C	7.29	1.62	1.52

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	S	152	GLN	CA-C-N	7.36	127.03	119.82
5	S	152	GLN	C-N-CA	7.36	127.03	119.82
5	E	202	ARG	N-CA-C	-6.10	105.87	113.38
1	O	152	ASP	CA-C-N	5.89	127.20	119.84
1	O	152	ASP	C-N-CA	5.89	127.20	119.84

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
5	E	204	GLU	Peptide
5	E	207(E)	ASN	Mainchain
14	N	181	ALA	Mainchain
1	O	200	SER	Mainchain
5	S	204	GLU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1926	28	0
1	O	1915	0	1926	24	0
2	B	1829	0	1829	25	0
2	P	1829	0	1829	27	0
3	C	1891	0	1900	27	0
3	Q	1891	0	1900	38	0
4	D	1794	0	1778	21	0
4	R	1786	0	1763	19	0
5	E	1795	0	1797	22	0
5	S	1795	0	1797	27	0
6	F	1840	0	1838	30	0
6	T	1840	0	1838	21	0
7	G	1921	0	1910	34	0
7	U	1921	0	1910	34	0
8	H	1685	0	1688	17	0
8	V	1685	0	1688	21	0
9	I	1581	0	1574	17	0
9	W	1581	0	1574	23	0
10	J	1585	0	1590	14	0
10	X	1582	0	1583	20	0
11	K	1644	0	1595	16	0
11	Y	1644	0	1595	13	0
12	L	1757	0	1711	21	0
12	Z	1757	0	1711	25	0
13	1	1824	0	1832	24	0
13	M	1824	0	1832	27	0
14	2	1512	0	1481	16	0
14	N	1512	0	1481	14	0
15	F	2	0	0	0	0
15	G	1	0	0	0	0
15	H	1	0	0	0	0
15	I	2	0	0	0	0
15	K	1	0	0	0	0
15	L	2	0	0	0	0
15	N	1	0	0	0	0
16	K	43	0	38	2	0
16	Y	43	0	38	1	0
17	K	12	0	13	1	0
17	Y	12	0	13	1	0
All	All	49255	0	48978	549	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 549 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:127:LEU:HD12	4:D:127:LEU:N	1.53	1.02
4:D:127:LEU:H	4:D:127:LEU:CD1	1.70	1.01
1:A:130:ARG:HH21	7:G:124:THR:CG2	1.78	0.97
4:D:127:LEU:C	4:D:127:LEU:HD13	1.89	0.96
4:D:127:LEU:HD12	4:D:127:LEU:H	0.81	0.95

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	239 (96%)	6 (2%)	3 (1%)	10	27
1	O	248/250 (99%)	234 (94%)	12 (5%)	2 (1%)	16	37
2	B	233/235 (99%)	223 (96%)	7 (3%)	3 (1%)	9	25
2	P	233/235 (99%)	218 (94%)	11 (5%)	4 (2%)	7	19
3	C	239/241 (99%)	229 (96%)	6 (2%)	4 (2%)	7	19
3	Q	239/241 (99%)	230 (96%)	6 (2%)	3 (1%)	9	25
4	D	229/260 (88%)	218 (95%)	11 (5%)	0	100	100
4	R	228/260 (88%)	219 (96%)	9 (4%)	0	100	100
5	E	231/233 (99%)	217 (94%)	9 (4%)	5 (2%)	5	14
5	S	231/233 (99%)	216 (94%)	11 (5%)	4 (2%)	7	19
6	F	234/242 (97%)	223 (95%)	10 (4%)	1 (0%)	30	54
6	T	234/242 (97%)	221 (94%)	13 (6%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
7	G	241/243 (99%)	234 (97%)	6 (2%)	1 (0%)	30	54
7	U	241/243 (99%)	232 (96%)	7 (3%)	2 (1%)	16	37
8	H	220/222 (99%)	211 (96%)	8 (4%)	1 (0%)	24	48
8	V	220/222 (99%)	213 (97%)	6 (3%)	1 (0%)	24	48
9	I	202/204 (99%)	192 (95%)	9 (4%)	1 (0%)	24	48
9	W	202/204 (99%)	198 (98%)	4 (2%)	0	100	100
10	J	196/198 (99%)	188 (96%)	7 (4%)	1 (0%)	24	48
10	X	196/198 (99%)	185 (94%)	8 (4%)	3 (2%)	8	22
11	K	210/212 (99%)	201 (96%)	8 (4%)	1 (0%)	24	48
11	Y	210/212 (99%)	204 (97%)	6 (3%)	0	100	100
12	L	220/222 (99%)	213 (97%)	7 (3%)	0	100	100
12	Z	220/222 (99%)	213 (97%)	7 (3%)	0	100	100
13	1	231/233 (99%)	221 (96%)	9 (4%)	1 (0%)	30	54
13	M	231/233 (99%)	214 (93%)	17 (7%)	0	100	100
14	2	194/196 (99%)	186 (96%)	8 (4%)	0	100	100
14	N	194/196 (99%)	189 (97%)	5 (3%)	0	100	100
All	All	6255/6382 (98%)	5981 (96%)	233 (4%)	41 (1%)	18	41

5 of 41 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	54	VAL
3	C	203	THR
3	C	207	ALA
5	E	6	ASN
6	F	184	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	184 (88%)	25 (12%)	5	12
1	O	209/209 (100%)	197 (94%)	12 (6%)	18	43
2	B	195/195 (100%)	180 (92%)	15 (8%)	12	30
2	P	195/195 (100%)	179 (92%)	16 (8%)	10	27
3	C	213/213 (100%)	195 (92%)	18 (8%)	10	25
3	Q	213/213 (100%)	202 (95%)	11 (5%)	21	47
4	D	192/215 (89%)	179 (93%)	13 (7%)	14	35
4	R	191/215 (89%)	172 (90%)	19 (10%)	7	19
5	E	192/192 (100%)	171 (89%)	21 (11%)	6	16
5	S	192/192 (100%)	178 (93%)	14 (7%)	13	32
6	F	195/200 (98%)	176 (90%)	19 (10%)	8	20
6	T	195/200 (98%)	180 (92%)	15 (8%)	12	30
7	G	207/207 (100%)	194 (94%)	13 (6%)	16	39
7	U	207/207 (100%)	198 (96%)	9 (4%)	26	54
8	H	181/181 (100%)	169 (93%)	12 (7%)	15	36
8	V	181/181 (100%)	171 (94%)	10 (6%)	19	45
9	I	172/172 (100%)	164 (95%)	8 (5%)	23	51
9	W	172/172 (100%)	165 (96%)	7 (4%)	27	56
10	J	175/175 (100%)	161 (92%)	14 (8%)	11	28
10	X	174/175 (99%)	162 (93%)	12 (7%)	14	34
11	K	169/169 (100%)	156 (92%)	13 (8%)	12	30
11	Y	169/169 (100%)	162 (96%)	7 (4%)	27	56
12	L	185/185 (100%)	178 (96%)	7 (4%)	29	58
12	Z	185/185 (100%)	176 (95%)	9 (5%)	22	49
13	1	199/199 (100%)	189 (95%)	10 (5%)	22	48
13	M	199/199 (100%)	191 (96%)	8 (4%)	28	56
14	2	162/162 (100%)	157 (97%)	5 (3%)	35	65
14	N	162/162 (100%)	152 (94%)	10 (6%)	16	39
All	All	5290/5348 (99%)	4938 (93%)	352 (7%)	15	36

5 of 352 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	Q	156	ILE
7	U	124	THR
4	R	48	LEU
5	S	65	LYS
8	V	153	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 195 such sidechains are listed below:

Mol	Chain	Res	Type
3	Q	163	GLN
7	U	121	GLN
4	R	23	GLN
5	S	125	GLN
8	V	57	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
17	MES	Y	302	-	12,12,12	2.31	1 (8%)	15,16,16	1.30	1 (6%)
16	P3N	K	302	-	46,46,46	1.29	5 (10%)	61,64,64	1.81	9 (14%)
17	MES	K	303	-	12,12,12	2.28	1 (8%)	15,16,16	1.16	1 (6%)
16	P3N	Y	301	-	46,46,46	1.26	6 (13%)	61,64,64	1.75	7 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
17	MES	Y	302	-	-	3/6/14/14	0/1/1/1
16	P3N	K	302	-	-	0/40/40/40	0/4/4/4
17	MES	K	303	-	-	1/6/14/14	0/1/1/1
16	P3N	Y	301	-	-	7/40/40/40	0/4/4/4

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
17	Y	302	MES	C8-S	-7.73	1.66	1.77
17	K	303	MES	C8-S	-7.61	1.66	1.77
16	K	302	P3N	C1-N36	5.62	1.35	1.29
16	Y	301	P3N	C1-N36	5.09	1.34	1.29
16	Y	301	P3N	C32-C31	-2.85	1.34	1.41

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
16	K	302	P3N	O29-C28-C35	6.33	124.60	115.89
16	Y	301	P3N	O29-C28-C35	5.84	123.92	115.89
16	K	302	P3N	O39-C1-N36	-5.63	109.06	113.62
16	Y	301	P3N	O39-C1-N36	-5.58	109.11	113.62
16	K	302	P3N	C35-C28-C32	-4.97	125.46	134.05

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
17	Y	302	MES	C7-C8-S-O1S
17	Y	302	MES	C7-C8-S-O3S

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
16	Y	301	P3N	N38-C37-C40-C41
16	Y	301	P3N	N38-C37-C40-C42
16	Y	301	P3N	N38-C37-C40-C43


There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
17	Y	302	MES	1	0
16	K	302	P3N	2	0
17	K	303	MES	1	0
16	Y	301	P3N	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/250 (100%)	0.24	10 (4%) 42 38	35, 50, 71, 89	0
1	O	250/250 (100%)	0.33	13 (5%) 33 29	37, 55, 83, 94	0
2	B	235/235 (100%)	0.53	21 (8%) 15 13	34, 53, 86, 91	0
2	P	235/235 (100%)	0.53	19 (8%) 18 15	33, 55, 85, 94	0
3	C	241/241 (100%)	0.71	25 (10%) 11 10	38, 61, 100, 118	0
3	Q	241/241 (100%)	1.21	59 (24%) 2 1	42, 65, 110, 128	0
4	D	233/260 (89%)	0.44	12 (5%) 33 29	36, 56, 81, 97	0
4	R	232/260 (89%)	0.50	12 (5%) 33 29	35, 57, 80, 94	0
5	E	233/233 (100%)	0.60	14 (6%) 27 24	42, 58, 84, 94	0
5	S	233/233 (100%)	0.81	28 (12%) 9 7	40, 60, 90, 102	0
6	F	236/242 (97%)	0.28	8 (3%) 48 44	33, 51, 80, 88	0
6	T	236/242 (97%)	0.37	6 (2%) 58 55	32, 52, 77, 98	0
7	G	243/243 (100%)	0.09	8 (3%) 49 45	30, 47, 72, 96	0
7	U	243/243 (100%)	0.09	9 (3%) 45 41	31, 47, 70, 86	0
8	H	222/222 (100%)	-0.00	2 (0%) 81 80	34, 44, 61, 88	0
8	V	222/222 (100%)	0.13	6 (2%) 56 53	37, 47, 64, 87	0
9	I	204/204 (100%)	-0.14	0 100 100	31, 43, 57, 64	0
9	W	204/204 (100%)	-0.02	3 (1%) 72 70	31, 44, 61, 66	0
10	J	198/198 (100%)	0.07	8 (4%) 42 38	33, 45, 58, 110	0
10	X	198/198 (100%)	0.15	9 (4%) 38 34	36, 46, 62, 111	0
11	K	212/212 (100%)	-0.06	3 (1%) 73 72	33, 42, 59, 72	0
11	Y	212/212 (100%)	-0.03	1 (0%) 87 86	32, 44, 60, 65	0
12	L	222/222 (100%)	-0.01	1 (0%) 87 86	32, 44, 65, 72	0
12	Z	222/222 (100%)	0.01	4 (1%) 67 65	32, 44, 64, 73	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
13	1	233/233 (100%)	-0.02	1 (0%) 88 87	31, 43, 59, 64	0
13	M	233/233 (100%)	0.07	6 (2%) 57 54	33, 45, 59, 62	0
14	2	196/196 (100%)	-0.12	2 (1%) 79 78	33, 41, 58, 72	0
14	N	196/196 (100%)	-0.13	0 100 100	32, 41, 57, 71	0
All	All	6315/6382 (98%)	0.25	290 (4%) 37 34	30, 48, 83, 128	0

The worst 5 of 290 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
10	X	-1	MET	9.6
3	Q	56	LEU	9.6
2	B	217	ALA	6.9
5	E	4	PHE	6.8
3	Q	55	THR	6.7

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
15	MG	I	201	1/1	0.69	0.20	58,58,58,58	0
15	MG	L	201	1/1	0.74	0.19	56,56,56,56	0
15	MG	L	202	1/1	0.86	0.15	54,54,54,54	0
15	MG	H	301	1/1	0.91	0.13	66,66,66,66	0
15	MG	F	302	1/1	0.91	0.22	107,107,107,107	0
15	MG	F	301	1/1	0.93	0.09	61,61,61,61	0
16	P3N	K	302	43/43	0.93	0.11	38,45,57,58	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
16	P3N	Y	301	43/43	0.93	0.11	40,45,58,59	0
15	MG	G	301	1/1	0.94	0.11	59,59,59,59	0
15	MG	N	201	1/1	0.94	0.08	47,47,47,47	0
17	MES	K	303	12/12	0.94	0.12	62,66,69,69	0
17	MES	Y	302	12/12	0.94	0.12	66,69,71,71	0
15	MG	I	202	1/1	0.96	0.13	57,57,57,57	0
15	MG	K	301	1/1	0.97	0.08	53,53,53,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.