



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:25 PM UTC

PDB ID : 3SDO / pdb_00003sdo
Title : Structure of a Nitrilotriacetate monooxygenase from Burkholderia pseudomallei
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2011-06-09
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	453	4% 93% 5%
1	B	453	7% 79% 8% 13%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6879 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitrilotriacetate monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	443	Total	C	N	O	S	0	1	0
			3392	2144	606	638	4			
1	B	396	Total	C	N	O	S	0	1	0
			2985	1890	537	554	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q3JIP8
A	-2	PRO	-	expression tag	UNP Q3JIP8
A	-1	GLY	-	expression tag	UNP Q3JIP8
A	0	SER	-	expression tag	UNP Q3JIP8
B	-3	GLY	-	expression tag	UNP Q3JIP8
B	-2	PRO	-	expression tag	UNP Q3JIP8
B	-1	GLY	-	expression tag	UNP Q3JIP8
B	0	SER	-	expression tag	UNP Q3JIP8

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	273	Total	O	0	0
			273	273		
3	B	217	Total	O	0	0
			217	217		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nitrilotriacetate monooxygenase



• Molecule 1: Nitrilotriacetate monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.19Å 92.78Å 162.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.68 – 2.00 48.68 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.1 (48.68-2.00) 95.4 (48.68-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.59 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.172 , 0.213 0.187 , 0.228	Depositor DCC
R_{free} test set	2887 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.134	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6879	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/3484	0.81	0/4739
1	B	0.71	0/3064	0.81	1/4169 (0.0%)
All	All	0.72	0/6548	0.81	1/8908 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	365	GLU	N-CA-C	-5.62	98.82	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3392	0	3218	13	0
1	B	2985	0	2806	32	0
2	A	8	0	12	0	0
2	B	4	0	6	0	0
3	A	273	0	0	1	0
3	B	217	0	0	1	0
All	All	6879	0	6042	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:378:MET:HE1	1:B:408:VAL:HG11	1.52	0.90
1:B:374:VAL:HG12	1:B:378:MET:HE2	1.52	0.89
1:B:363:THR:HG22	1:B:368:ILE:HD11	1.76	0.67
1:A:126:THR:HG21	1:A:147:TYR:CD1	2.33	0.63
1:B:378:MET:HE1	1:B:408:VAL:CG1	2.28	0.62
1:B:363:THR:CG2	1:B:368:ILE:HD11	2.32	0.58
1:B:374:VAL:HG12	1:B:378:MET:CE	2.27	0.58
1:B:363:THR:HG22	1:B:368:ILE:CD1	2.33	0.57
1:A:126:THR:HG21	1:A:147:TYR:CE1	2.39	0.57
1:B:233:PHE:CE1	1:B:391:MET:HG3	2.39	0.57
1:B:363:THR:HG22	1:B:368:ILE:CG1	2.37	0.55
1:A:227:ARG:NH2	3:A:555:HOH:O	2.41	0.54
1:B:11:LEU:HD11	1:B:13[A]:GLN:HG2	1.89	0.53
1:B:375:ALA:HA	1:B:378:MET:HE3	1.90	0.52
1:A:95:MET:SD	1:A:105:VAL:HG22	2.50	0.52
1:B:122:TRP:CE2	1:B:124:VAL:HG22	2.45	0.50
1:B:363:THR:HG22	1:B:368:ILE:HG13	1.93	0.50
1:A:335:ASN:OD1	1:A:338:ARG:NH2	2.45	0.49
1:A:264:LYS:HE3	1:A:431:LEU:O	2.13	0.49
1:B:374:VAL:CG1	1:B:378:MET:HE2	2.34	0.48
1:A:378:MET:HE1	1:A:404:PHE:HE2	1.77	0.48
1:B:219:ASP:OD1	1:B:248:ARG:NH2	2.48	0.46
1:B:300:LEU:CD1	1:B:304:PHE:HE2	2.28	0.46
1:A:201:ASN:HB3	1:B:61:VAL:CG2	2.46	0.46
1:B:300:LEU:HD12	1:B:304:PHE:HE2	1.80	0.46
1:B:363:THR:CG2	1:B:368:ILE:CG1	2.95	0.45
1:A:291:LEU:CD2	1:A:355:SER:CB	2.95	0.45
1:A:340:ALA:O	1:A:343:ARG:O	2.36	0.44
1:B:186:ARG:NH1	1:B:188:ASP:OD2	2.50	0.44
1:B:207:GLN:HB3	1:B:426:THR:HB	1.98	0.44
1:A:108:GLN:HG2	1:B:74:GLU:CD	2.42	0.43
1:B:95:MET:SD	1:B:105:VAL:HG22	2.59	0.43
1:B:11:LEU:CG	1:B:13[A]:GLN:HG2	2.48	0.43
1:A:273:VAL:HG12	1:A:374:VAL:HG21	2.01	0.43
1:B:11:LEU:CD1	1:B:13[A]:GLN:HG2	2.49	0.42
1:B:267:PRO:HD2	1:B:389:GLY:O	2.19	0.42
1:B:236:GLY:N	3:B:642:HOH:O	2.43	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:11:LEU:HD21	1:B:13[A]:GLN:HG3	2.01	0.41
1:A:61:VAL:CG2	1:B:201:ASN:HB3	2.51	0.41
1:B:101:GLU:N	1:B:101:GLU:CD	2.79	0.41
1:B:197:GLU:HG2	1:B:198:GLY:O	2.21	0.41
1:B:186:ARG:HH21	1:B:197:GLU:CD	2.28	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/453 (98%)	426 (96%)	15 (3%)	1 (0%)	43	42
1	B	391/453 (86%)	372 (95%)	16 (4%)	3 (1%)	16	11
All	All	833/906 (92%)	798 (96%)	31 (4%)	4 (0%)	24	21

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	365	GLU
1	A	398	GLY
1	B	364	SER
1	B	141	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/361 (95%)	340 (99%)	2 (1%)	78	85
1	B	291/361 (81%)	288 (99%)	3 (1%)	68	75
All	All	633/722 (88%)	628 (99%)	5 (1%)	73	80

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	ASN
1	A	84	VAL
1	B	71	ASN
1	B	96	SER
1	B	405	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	GLN
1	A	123	ASN
1	A	140	HIS
1	A	145	GLN
1	A	209	GLN
1	A	306	GLN
1	B	108	GLN
1	B	123	ASN
1	B	262	HIS
1	B	407	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	450	-	3,3,3	0.62	0	2,2,2	0.21	0
2	EDO	B	450	-	3,3,3	0.70	0	2,2,2	0.51	0
2	EDO	A	451	-	3,3,3	0.58	0	2,2,2	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	450	-	-	0/1/1/1	-
2	EDO	B	450	-	-	1/1/1/1	-
2	EDO	A	451	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	450	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	443/453 (97%)	-0.13	17 (3%) 44 43	15, 24, 44, 57	1 (0%)
1	B	396/453 (87%)	0.05	32 (8%) 18 17	15, 25, 59, 79	1 (0%)
All	All	839/906 (92%)	-0.04	49 (5%) 29 27	15, 24, 50, 79	2 (0%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	297	LEU	6.7
1	B	364	SER	5.7
1	A	-1	GLY	5.1
1	B	139	PRO	4.7
1	B	298	ALA	4.6
1	B	305	GLN	4.4
1	B	299	TYR	4.1
1	B	143	HIS	4.1
1	A	361	ILE	4.1
1	A	359	SER	3.9
1	B	366	ALA	3.9
1	B	311	VAL	3.9
1	B	304	PHE	3.8
1	B	300	LEU	3.6
1	B	362	GLY	3.5
1	B	303	PHE	3.3
1	A	363	THR	3.1
1	A	360	ASN	3.1
1	A	0	SER	3.1
1	B	363	THR	3.0
1	A	323	THR	3.0
1	B	309	PHE	3.0
1	B	301	SER	2.9
1	A	299	TYR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	291	LEU	2.8
1	B	126	THR	2.6
1	B	351	ALA	2.6
1	B	350	VAL	2.5
1	B	348	ARG	2.5
1	B	142	ASP	2.5
1	A	236	GLY	2.5
1	B	140	HIS	2.4
1	A	321	ILE	2.4
1	A	356	THR	2.3
1	B	347	LEU	2.3
1	B	306	GLN	2.3
1	A	324	LEU	2.2
1	A	140	HIS	2.2
1	B	278	GLN	2.1
1	A	302	HIS	2.1
1	B	296	ALA	2.1
1	B	138	ARG	2.1
1	B	294	ARG	2.1
1	A	127	SER	2.1
1	A	357	ARG	2.1
1	A	235	ASN	2.1
1	B	141	PRO	2.1
1	B	144	ALA	2.1
1	B	353	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	A	451	4/4	0.93	0.10	30,32,32,34	0
2	EDO	B	450	4/4	0.96	0.09	28,28,29,29	0
2	EDO	A	450	4/4	0.97	0.04	24,26,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.