



wwPDB EM Validation Summary Report ⓘ

Mar 7, 2026 – 04:06 AM UTC

PDB ID : 6SD1 / pdb_00006sd1
EMDB ID : EMD-10145
Title : Structure of the RBM3/collar region of the Salmonella flagella MS-ring protein
FliF with 33-fold symmetry applied
Authors : Johnson, S.; Fong, Y.H.; Deme, J.C.; Furlong, E.J.; Kuhlen, L.; Lea, S.M.
Deposited on : 2019-07-26
Resolution : 2.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8728 (2.10 - 3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	560	21% 6% 73%
1	B	560	20% 7% 73%
1	C	560	22% 5% 73%
1	D	560	21% 6% 73%





Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	560	 22% 5% 73%
1	F	560	 21% 6% 73%
1	G	560	 21% 6% 73%
1	H	560	 20% 7% 73%
1	I	560	 21% 6% 73%
1	J	560	 21% 6% 73%
1	K	560	 22% 5% 73%
1	L	560	 20% 7% 73%
1	M	560	 20% 7% 73%
1	N	560	 21% 6% 73%
1	O	560	 20% 7% 73%
1	P	560	 20% 7% 73%
1	Q	560	 20% 7% 73%
1	R	560	 21% 6% 73%
1	S	560	 21% 6% 73%
1	T	560	 21% 6% 73%
1	U	560	 21% 6% 73%
1	V	560	 21% 6% 73%
1	W	560	 21% 6% 73%
1	X	560	 20% 7% 73%
1	Y	560	 20% 7% 73%
1	Z	560	 21% 6% 73%
1	a	560	 21% 6% 73%
1	b	560	 20% 7% 73%
1	c	560	 21% 6% 73%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	d	560	 21% 6% 73%
1	e	560	 21% 6% 73%
1	f	560	 21% 6% 73%
1	g	560	 21% 6% 73%

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 39369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Flagellar M-ring protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	151	1193	726	223	241	3	0	0
1	B	151	1193	726	223	241	3	0	0
1	C	151	1193	726	223	241	3	0	0
1	D	151	1193	726	223	241	3	0	0
1	E	151	1193	726	223	241	3	0	0
1	F	151	1193	726	223	241	3	0	0
1	G	151	1193	726	223	241	3	0	0
1	H	151	1193	726	223	241	3	0	0
1	I	151	1193	726	223	241	3	0	0
1	J	151	1193	726	223	241	3	0	0
1	K	151	1193	726	223	241	3	0	0
1	L	151	1193	726	223	241	3	0	0
1	M	151	1193	726	223	241	3	0	0
1	N	151	1193	726	223	241	3	0	0
1	O	151	1193	726	223	241	3	0	0
1	P	151	1193	726	223	241	3	0	0
1	Q	151	1193	726	223	241	3	0	0


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	S	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	T	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	U	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	V	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	W	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	X	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	Y	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	Z	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	a	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	b	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	c	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	d	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	e	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	f	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		
1	g	151	Total	C	N	O	S	0	0
			1193	726	223	241	3		




• Molecule 1: Flagellar M-ring protein




• Molecule 1: Flagellar M-ring protein







• Molecule 1: Flagellar M-ring protein




• Molecule 1: Flagellar M-ring protein







• Molecule 1: Flagellar M-ring protein




• Molecule 1: Flagellar M-ring protein





• Molecule 1: Flagellar M-ring protein



• Molecule 1: Flagellar M-ring protein



ASP	THR	THR	THR	THR	ASN	ASN	ASN	ASN	ALA	ALA	GLY	P3555	R360	R361	E362	I372	R373	S386	V387	A388	V389	V390	V391	N392	Y393	K394	THR	THR	ALA	ALA	ASP	GLY	LYS	PRO	L402	P403	L404	T405	A406	D407	K410	L415	E418	A419	M420	M431	V432	W433	N434	S435	P436	F437	S438	ALA	ALA	VAL									
ASN	THR	GLY	GLY	GLY	GLY	LEU	PRO	PHE	TRP	ARG	GLN	GLN	GLN	SER	PHE	ILE	ASP	GLN	LEU	LEU	ALA	ALA	GLY	ASN	TRP	LEU	LEU	VAL	VAL	VAL	ALA	TRP	ILE	LEU	ARG	TRP	ILE	LEU	TRP	ARG	LYS	ALA	VAL	ARG	GLN	PRO	ASN	PRO	ASP	ARG	VAL	LEU	GLU	GLU	ALA	ALA	ALA	ALA	GLN	GLN	GLN	GLN	ALA	ALA	VAL
GLN	GLU	THR	GLU	GLU	ALA	VAL	GLU	ARG	LEU	SER	LYS	ASP	GLU	GLN	GLN	GLN	GLN	ARG	ALA	ALA	ALA	ASN	GLN	ARG	LEU	LEU	GLY	ALA	GLU	VAL	MET	SER	GLN	ARG	ILE	ARG	GLU	MET	SER	ASP	ASN	ASP	PRO	ARG	VAL	VAL	ALA	LEU	VAL	ILE	ARG	GLN	TRP	MET	SER	ASN	ASP	HIS	GLU						

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C33	Depositor
Number of particles used	77849	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	48	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.100	Depositor
Minimum map value	-0.057	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.01	Depositor
Map size (Å)	355.104, 355.104, 355.104	wwPDB
Map dimensions	432, 432, 432	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.822, 0.822, 0.822	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.16	0/1205	0.34	0/1624
1	B	0.16	0/1205	0.33	0/1624
1	C	0.16	0/1205	0.34	0/1624
1	D	0.16	0/1205	0.33	0/1624
1	E	0.16	0/1205	0.34	0/1624
1	F	0.17	0/1205	0.33	0/1624
1	G	0.16	0/1205	0.34	0/1624
1	H	0.17	0/1205	0.33	0/1624
1	I	0.16	0/1205	0.33	0/1624
1	J	0.16	0/1205	0.33	0/1624
1	K	0.16	0/1205	0.33	0/1624
1	L	0.16	0/1205	0.33	0/1624
1	M	0.16	0/1205	0.32	0/1624
1	N	0.16	0/1205	0.33	0/1624
1	O	0.16	0/1205	0.33	0/1624
1	P	0.16	0/1205	0.32	0/1624
1	Q	0.16	0/1205	0.34	0/1624
1	R	0.17	0/1205	0.33	0/1624
1	S	0.16	0/1205	0.33	0/1624
1	T	0.17	0/1205	0.33	0/1624
1	U	0.16	0/1205	0.32	0/1624
1	V	0.16	0/1205	0.33	0/1624
1	W	0.16	0/1205	0.33	0/1624
1	X	0.16	0/1205	0.33	0/1624
1	Y	0.17	0/1205	0.33	0/1624
1	Z	0.16	0/1205	0.34	0/1624
1	a	0.16	0/1205	0.32	0/1624
1	b	0.16	0/1205	0.33	0/1624
1	c	0.16	0/1205	0.34	0/1624
1	d	0.16	0/1205	0.33	0/1624
1	e	0.17	0/1205	0.33	0/1624
1	f	0.17	0/1205	0.33	0/1624
1	g	0.16	0/1205	0.33	0/1624
All	All	0.16	0/39765	0.33	0/53592

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1193	0	1173	22	0
1	B	1193	0	1173	24	0
1	C	1193	0	1173	19	0
1	D	1193	0	1173	23	0
1	E	1193	0	1173	21	0
1	F	1193	0	1173	25	0
1	G	1193	0	1173	23	0
1	H	1193	0	1173	26	0
1	I	1193	0	1173	23	0
1	J	1193	0	1173	24	0
1	K	1193	0	1173	22	0
1	L	1193	0	1173	26	0
1	M	1193	0	1173	26	0
1	N	1193	0	1173	25	0
1	O	1193	0	1173	26	0
1	P	1193	0	1173	28	0
1	Q	1193	0	1173	27	0
1	R	1193	0	1173	25	0
1	S	1193	0	1173	26	0
1	T	1193	0	1173	22	0
1	U	1193	0	1173	24	0
1	V	1193	0	1173	23	0
1	W	1193	0	1173	24	0
1	X	1193	0	1173	27	0
1	Y	1193	0	1173	27	0
1	Z	1193	0	1173	23	0
1	a	1193	0	1173	24	0
1	b	1193	0	1173	26	0
1	c	1193	0	1173	23	0
1	d	1193	0	1173	23	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	e	1193	0	1173	24	0
1	f	1193	0	1173	23	0
1	g	1193	0	1173	26	0
All	All	39369	0	38709	676	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 676 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Q:280:GLU:HG3	1:Q:372:ILE:HG12	1.80	0.64
1:B:250:GLU:O	1:B:254:SER:OG	2.15	0.63
1:H:360:ARG:NH1	1:H:362:GLU:OE1	2.31	0.63
1:P:405:THR:HG22	1:P:407:ASP:H	1.64	0.62
1:G:405:THR:HG22	1:G:407:ASP:H	1.64	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/560 (26%)	138 (95%)	7 (5%)	0	100	100
1	B	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	C	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	D	145/560 (26%)	138 (95%)	7 (5%)	0	100	100
1	E	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	F	145/560 (26%)	138 (95%)	7 (5%)	0	100	100
1	G	145/560 (26%)	139 (96%)	6 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	145/560 (26%)	138 (95%)	7 (5%)	0	100	100
1	I	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	J	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	K	145/560 (26%)	136 (94%)	9 (6%)	0	100	100
1	L	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	M	145/560 (26%)	140 (97%)	5 (3%)	0	100	100
1	N	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	O	145/560 (26%)	140 (97%)	5 (3%)	0	100	100
1	P	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	Q	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	R	145/560 (26%)	138 (95%)	7 (5%)	0	100	100
1	S	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	T	145/560 (26%)	138 (95%)	7 (5%)	0	100	100
1	U	145/560 (26%)	140 (97%)	5 (3%)	0	100	100
1	V	145/560 (26%)	138 (95%)	7 (5%)	0	100	100
1	W	145/560 (26%)	140 (97%)	5 (3%)	0	100	100
1	X	145/560 (26%)	137 (94%)	8 (6%)	0	100	100
1	Y	145/560 (26%)	140 (97%)	5 (3%)	0	100	100
1	Z	145/560 (26%)	140 (97%)	5 (3%)	0	100	100
1	a	145/560 (26%)	138 (95%)	7 (5%)	0	100	100
1	b	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	c	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	d	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	e	145/560 (26%)	138 (95%)	7 (5%)	0	100	100
1	f	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
1	g	145/560 (26%)	139 (96%)	6 (4%)	0	100	100
All	All	4785/18480 (26%)	4579 (96%)	206 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/467 (29%)	134 (100%)	0	100	100
1	B	134/467 (29%)	134 (100%)	0	100	100
1	C	134/467 (29%)	134 (100%)	0	100	100
1	D	134/467 (29%)	134 (100%)	0	100	100
1	E	134/467 (29%)	134 (100%)	0	100	100
1	F	134/467 (29%)	134 (100%)	0	100	100
1	G	134/467 (29%)	134 (100%)	0	100	100
1	H	134/467 (29%)	134 (100%)	0	100	100
1	I	134/467 (29%)	134 (100%)	0	100	100
1	J	134/467 (29%)	134 (100%)	0	100	100
1	K	134/467 (29%)	134 (100%)	0	100	100
1	L	134/467 (29%)	134 (100%)	0	100	100
1	M	134/467 (29%)	134 (100%)	0	100	100
1	N	134/467 (29%)	134 (100%)	0	100	100
1	O	134/467 (29%)	134 (100%)	0	100	100
1	P	134/467 (29%)	134 (100%)	0	100	100
1	Q	134/467 (29%)	134 (100%)	0	100	100
1	R	134/467 (29%)	134 (100%)	0	100	100
1	S	134/467 (29%)	134 (100%)	0	100	100
1	T	134/467 (29%)	134 (100%)	0	100	100
1	U	134/467 (29%)	134 (100%)	0	100	100
1	V	134/467 (29%)	134 (100%)	0	100	100
1	W	134/467 (29%)	134 (100%)	0	100	100
1	X	134/467 (29%)	134 (100%)	0	100	100
1	Y	134/467 (29%)	134 (100%)	0	100	100
1	Z	134/467 (29%)	134 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	a	134/467 (29%)	134 (100%)	0	100	100
1	b	134/467 (29%)	134 (100%)	0	100	100
1	c	134/467 (29%)	134 (100%)	0	100	100
1	d	134/467 (29%)	134 (100%)	0	100	100
1	e	134/467 (29%)	134 (100%)	0	100	100
1	f	134/467 (29%)	134 (100%)	0	100	100
1	g	134/467 (29%)	134 (100%)	0	100	100
All	All	4422/15411 (29%)	4422 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 114 such sidechains are listed below:

Mol	Chain	Res	Type
1	R	434	ASN
1	g	261	ASN
1	V	261	ASN
1	g	259	ASN
1	d	261	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-10145. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 216




Y Index: 216



Z Index: 216


6.2.2 Raw map



X Index: 216



Y Index: 216




Z Index: 216


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 274




Y Index: 156




Z Index: 197


6.3.2 Raw map



X Index: 275



Y Index: 157




Z Index: 199


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_10145_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 254 nm³; this corresponds to an approximate mass of 229 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.385 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.385 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.60	-	-
Author-provided FSC curve	2.55	2.86	2.58
Unmasked-calculated*	2.91	3.24	2.93

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 2.91 differs from the reported value 2.6 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-10145 and PDB model 6SD1. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).





































































9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 92% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9180	 0.6230
A	 0.9170	 0.6230
B	 0.9200	 0.6220
C	 0.9150	 0.6210
D	 0.9190	 0.6220
E	 0.9170	 0.6210
F	 0.9190	 0.6230
G	 0.9190	 0.6260
H	 0.9190	 0.6240
I	 0.9180	 0.6220
J	 0.9180	 0.6240
K	 0.9140	 0.6230
L	 0.9230	 0.6230
M	 0.9190	 0.6210
N	 0.9180	 0.6230
O	 0.9150	 0.6230
P	 0.9180	 0.6240
Q	 0.9150	 0.6240
R	 0.9160	 0.6240
S	 0.9150	 0.6230
T	 0.9190	 0.6220
U	 0.9180	 0.6220
V	 0.9130	 0.6230
W	 0.9180	 0.6240
X	 0.9180	 0.6220
Y	 0.9200	 0.6260
Z	 0.9170	 0.6250
a	 0.9170	 0.6220
b	 0.9220	 0.6250
c	 0.9150	 0.6220
d	 0.9190	 0.6180
e	 0.9170	 0.6250
f	 0.9160	 0.6250
g	 0.9160	 0.6240

