



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:27 PM UTC

PDB ID : 1SGN / pdb_00001sgn
Title : ASN 18 VARIANT OF TURKEY OVOMUCOID INHIBITOR THIRD DOMAIN COMPLEXED WITH STREPTOMYCES GRISEUS PROTEINASE B
Authors : Huang, K.; Lu, W.; Anderson, S.; Laskowski Jr., M.; James, M.N.G.
Deposited on : 1999-03-25
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	E	185	 80% 17% .
2	I	51	 71% 25% . .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1862 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Streptogrisin B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	185	Total	C	N	O	S	0	0	0
			1309	799	228	276	6			

- Molecule 2 is a protein called Ovomucoid.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	51	Total	C	N	O	S	0	0	0
			387	236	66	79	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	18	ASN	LEU	engineered mutation	UNP P68390

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	124	Total	O	0	0
			124	124		
4	I	37	Total	O	0	0
			37	37		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Streptogrisin B

Chain E:  80% 17% .



• Molecule 2: Ovomucoid

Chain I:  71% 25% ..



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.58Å 54.67Å 45.61Å 90.00° 119.17° 90.00°	Depositor
Resolution (Å)	20.00 – 1.80	Depositor
% Data completeness (in resolution range)	89.0 (20.00-1.80)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT	Depositor
R, R_{free}	0.159 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1862	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	1.62	6/1334 (0.4%)	1.64	16/1818 (0.9%)
2	I	1.60	1/395 (0.3%)	1.77	5/533 (0.9%)
All	All	1.61	7/1729 (0.4%)	1.67	21/2351 (0.9%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	214	SER	C-N	-5.79	1.26	1.33
1	E	210	ILE	CA-C	5.56	1.58	1.52
2	I	48	LEU	N-CA	5.54	1.52	1.46
1	E	29	ASP	CG-OD1	5.52	1.35	1.25
1	E	114	PRO	CA-C	5.35	1.59	1.52
1	E	201	SER	C-O	-5.32	1.17	1.23
1	E	140	GLY	C-O	5.16	1.30	1.23

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	120	GLY	O-C-N	-8.16	113.38	122.61
2	I	47	THR	N-CA-CB	-7.08	99.70	110.33
1	E	129	ASN	CA-CB-CG	-6.53	106.07	112.60
1	E	109	THR	N-CA-C	-6.29	105.38	114.12
1	E	138	ARG	CD-NE-CZ	5.82	132.55	124.40
1	E	240	SER	CA-CB-OG	-5.73	99.64	111.10
1	E	198	PRO	CB-CA-C	-5.71	103.50	110.98
1	E	194	ASP	CA-CB-CG	5.67	118.27	112.60
1	E	33	SER	CA-CB-OG	-5.64	99.82	111.10
2	I	51	SER	CA-CB-OG	-5.63	99.83	111.10
1	E	100	ASN	N-CA-CB	5.46	119.72	110.49
1	E	63	ALA	CA-C-N	-5.44	114.40	122.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	63	ALA	C-N-CA	-5.44	114.40	122.56
1	E	81	ARG	NE-CZ-NH2	5.34	124.01	119.20
2	I	9	SER	CA-C-N	-5.33	113.98	122.66
2	I	9	SER	C-N-CA	-5.33	113.98	122.66
1	E	181	ILE	CB-CA-C	-5.13	104.55	110.91
2	I	36	ASN	CA-CB-CG	-5.10	107.50	112.60
1	E	120	GLY	CA-C-N	-5.03	108.03	122.06
1	E	120	GLY	C-N-CA	-5.03	108.03	122.06
1	E	51	TYR	CA-CB-CG	5.01	122.93	113.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1309	0	1227	15	0
2	I	387	0	353	11	0
3	E	5	0	0	0	0
4	E	124	0	0	0	1
4	I	37	0	0	0	1
All	All	1862	0	1580	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:166:ASN:HD22	1:E:179:GLY:HA2	1.42	0.84
1:E:110:ASN:C	1:E:110:ASN:HD22	2.00	0.69
2:I:21:ARG:HH11	2:I:21:ARG:HB3	1.56	0.69
1:E:110:ASN:ND2	1:E:113:ILE:H	1.97	0.62
2:I:21:ARG:HH11	2:I:21:ARG:CG	2.14	0.61
2:I:21:ARG:HH11	2:I:21:ARG:CB	2.17	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:166:ASN:ND2	1:E:179:GLY:HA2	2.18	0.57
1:E:199:LEU:HD13	1:E:228:PHE:CZ	2.41	0.55
1:E:49:THR:HG22	1:E:51:TYR:CE1	2.42	0.54
1:E:113:ILE:O	1:E:115:LYS:NZ	2.37	0.54
1:E:33:SER:HB3	1:E:66:TRP:CH2	2.46	0.51
2:I:11:TYR:CE2	2:I:13:LYS:HD2	2.45	0.51
2:I:11:TYR:HB3	2:I:12:PRO:HA	1.92	0.51
2:I:44:SER:O	2:I:47:THR:HB	2.11	0.49
2:I:20:TYR:OH	2:I:55:LYS:HE2	2.12	0.48
1:E:68:ALA:HB2	1:E:85:LEU:HD21	1.96	0.47
1:E:67:TRP:CD2	1:E:81:ARG:HG3	2.50	0.47
1:E:110:ASN:HD22	1:E:111:THR:N	2.13	0.46
1:E:49:THR:CG2	1:E:51:TYR:CE1	2.99	0.45
1:E:181:ILE:O	1:E:227:PHE:HA	2.18	0.44
2:I:21:ARG:CG	2:I:21:ARG:NH1	2.78	0.43
2:I:21:ARG:NH1	2:I:21:ARG:HG2	2.34	0.43
2:I:21:ARG:HH11	2:I:21:ARG:HG2	1.82	0.43
1:E:110:ASN:HD21	1:E:113:ILE:H	1.64	0.41
1:E:192(B):PRO:HB2	2:I:32:GLY:HA3	2.02	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:552:HOH:O	4:I:69:HOH:O[2_657]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	E	183/185 (99%)	175 (96%)	8 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	49/51 (96%)	48 (98%)	1 (2%)	0	100	100
All	All	232/236 (98%)	223 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	138/138 (100%)	134 (97%)	4 (3%)	37	25
2	I	45/45 (100%)	41 (91%)	4 (9%)	9	2
All	All	183/183 (100%)	175 (96%)	8 (4%)	25	13

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	34	SER
1	E	110	ASN
1	E	222	SER
1	E	233	GLU
2	I	7	ASP
2	I	10	GLU
2	I	21	ARG
2	I	47	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	110	ASN
1	E	166	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	E	500	-	4,4,4	1.55	1 (25%)	6,6,6	0.35	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	500	PO4	P-O4	-2.05	1.48	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.