



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:35 PM UTC

PDB ID : 3SGK / pdb_00003sgk
Title : Unique carbohydrate/carbohydrate interactions are required for high affinity binding of FcgIII and antibodies lacking core fucose
Authors : Ferrara, C.; Grau, S.; Jaeger, C.; Sondermann, P.; Bruenker, P.; Waldhauer, I.; Hennig, M.; Ruf, A.; Rufer, A.C.; Stihle, M.; Umana, P.; Benz, J.
Deposited on : 2011-06-15
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	0.2% 76% 16% 7%
1	B	225	2% 73% 20% 6%
2	C	204	2% 63% 15% 19%
3	D	8	38% 62%
3	E	8	12% 62% 25%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	8	
5	G	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	E	1	X	-	-	-
3	MAN	E	4	X	-	-	-
4	MAN	F	4	X	-	-	-
4	MAN	F	6	X	-	-	-
4	MAN	F	7	X	-	-	-
4	MAN	F	8	X	-	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5163 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fc fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	209	Total	C	N	O	S	0	0	0
			1668	1061	281	320	6			
1	B	212	Total	C	N	O	S	0	0	0
			1694	1080	284	324	6			


- Molecule 2 is a protein called human Fcg3a receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	165	Total	C	N	O	S	0	0	0
			1331	850	229	248	4			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	38	GLN	ASN	engineered mutation	UNP P08637
C	74	GLN	ASN	engineered mutation	UNP P08637
C	158	VAL	PHE	SEE REMARK 999	UNP P08637
C	169	GLN	ASN	engineered mutation	UNP P08637
C	191	GLY	-	expression tag	UNP P08637
C	192	LYS	-	expression tag	UNP P08637
C	193	LYS	-	expression tag	UNP P08637
C	194	LYS	-	expression tag	UNP P08637
C	195	LYS	-	expression tag	UNP P08637
C	196	LYS	-	expression tag	UNP P08637
C	197	LYS	-	expression tag	UNP P08637
C	198	GLY	-	expression tag	UNP P08637
C	199	HIS	-	expression tag	UNP P08637
C	200	HIS	-	expression tag	UNP P08637
C	201	HIS	-	expression tag	UNP P08637
C	202	HIS	-	expression tag	UNP P08637
C	203	HIS	-	expression tag	UNP P08637
C	204	HIS	-	expression tag	UNP P08637

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)][2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	8	Total	C	N	O	0	0	0
			103	58	5	40			
3	E	8	Total	C	N	O	0	0	0
			103	58	5	40			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	8	Total	C	N	O	0	0	0
			94	52	2	40			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	G	2	Total	C	N	O	0	0	0
			28	16	2	10			


- Molecule 6 is MALONATE ION (CCD ID: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	54	Total	O	0	0
			54	54		
7	B	30	Total	O	0	0
			30	30		
7	C	51	Total	O	0	0
			51	51		

Chain D:  38% 62%

NAG1	NAG2	BR43	MAN4	NAG5	MAN6	NAG7	NAG8
------	------	------	------	------	------	------	------

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)][2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  12% 62% 25%

NAG1	NAG2	BR43	MAN4	NAG5	MAN6	NAG7	NAG8
------	------	------	------	------	------	------	------

- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  12% 50% 38%

NAG1	NAG2	BR43	MAN4	NAG5	MAN6	NAG7	NAG8
------	------	------	------	------	------	------	------

- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

NAG1	NAG2
------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.34Å 88.23Å 141.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.40 50.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.40) 99.9 (50.00-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.6.0112	Depositor
R, R_{free}	0.205 , 0.250 0.208 , 0.250	Depositor DCC
R_{free} test set	1687 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	61.2	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5163	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/1714	0.78	0/2335
1	B	0.70	0/1741	0.79	0/2372
2	C	0.79	0/1366	0.83	0/1854
All	All	0.73	0/4821	0.79	0/6561

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1668	0	1633	22	0
1	B	1694	0	1664	27	0
2	C	1331	0	1287	18	0
3	D	103	0	88	0	0
3	E	103	0	88	1	0
4	F	94	0	79	4	0
5	G	28	0	24	0	0
6	A	7	0	2	0	0
7	A	54	0	0	2	0
7	B	30	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	51	0	0	3	0
All	All	5163	0	4865	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:C:302:HOH:O	4:F:1:NAG:C8	2.05	1.03
1:B:236:GLY:O	2:C:161:LYS:HE3	1.82	0.79
1:B:350:THR:HB	1:B:441:LEU:HG	1.63	0.79
1:B:338:LYS:HE3	1:B:430:GLU:OE1	1.84	0.77
3:E:5:NAG:H5	3:E:8:NAG:H5	1.65	0.77
7:C:302:HOH:O	4:F:1:NAG:H83	1.77	0.75
1:A:346:PRO:HB3	1:A:372:PHE:HB3	1.68	0.73
1:A:274:LYS:HE3	1:A:276:ASN:HD21	1.55	0.71
1:B:419:GLN:HG3	1:B:420:GLY:HA2	1.73	0.69
2:C:103:GLU:HA	2:C:143:LYS:HB2	1.74	0.69
1:B:238:PRO:HD2	1:B:328:LEU:HD21	1.76	0.68
7:C:302:HOH:O	4:F:1:NAG:H81	1.82	0.68
1:A:428:MET:HE2	7:A:642:HOH:O	1.94	0.66
1:A:326:LYS:HA	1:A:326:LYS:HE2	1.78	0.64
2:C:63:VAL:HG21	2:C:88:ILE:HD11	1.78	0.63
1:A:429:HIS:CD2	1:A:431:ALA:H	2.17	0.62
1:A:429:HIS:HD2	1:A:431:ALA:H	1.49	0.60
1:A:311:GLN:HE21	1:A:311:GLN:HA	1.66	0.60
1:B:379:VAL:HG22	1:B:427:VAL:HG22	1.83	0.60
1:B:238:PRO:HD2	1:B:328:LEU:CD2	2.33	0.56
1:B:374:PRO:O	1:B:429:HIS:HE1	1.87	0.56
1:A:320:LYS:HG3	1:A:335:THR:HG22	1.88	0.55
1:B:338:LYS:HE2	1:B:374:PRO:HG3	1.89	0.54
1:A:415:SER:O	1:A:419:GLN:HG3	2.07	0.54
1:B:360:LYS:O	1:B:414:LYS:HD3	2.07	0.54
1:A:374:PRO:O	1:A:429:HIS:HE1	1.92	0.53
1:B:429:HIS:CD2	1:B:431:ALA:H	2.27	0.53
1:B:429:HIS:HD2	1:B:431:ALA:H	1.58	0.51
1:B:351:LEU:HB2	1:B:366:THR:HB	1.91	0.51
2:C:63:VAL:CG2	2:C:88:ILE:HD11	2.43	0.49
1:B:365:LEU:HD12	1:B:410:LEU:HD23	1.95	0.48
1:A:409:LYS:HB2	1:B:407:TYR:OH	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:278:TYR:HB2	1:B:320:LYS:HB3	1.96	0.48
1:B:426:SER:HB3	1:B:438:GLN:HG3	1.95	0.48
2:C:20:LEU:HD21	2:C:92:LEU:HB2	1.96	0.48
2:C:43:PHE:CE2	2:C:48:LEU:HD13	2.49	0.48
2:C:93:LEU:HD23	2:C:168:VAL:HG22	1.97	0.47
2:C:73:THR:H	2:C:76:SER:HB2	1.80	0.47
1:B:269:GLU:HB2	7:B:512:HOH:O	2.13	0.47
1:B:419:GLN:HG3	1:B:420:GLY:CA	2.43	0.47
2:C:119:HIS:HD2	2:C:120:LYS:HG3	1.79	0.46
1:A:342:GLN:HB2	7:A:638:HOH:O	2.15	0.46
2:C:130:ARG:NH2	2:C:148:ASP:OD1	2.49	0.45
1:A:360:LYS:O	1:A:414:LYS:HD3	2.16	0.45
4:F:3:BMA:H62	4:F:6:MAN:H2	1.54	0.44
1:A:344:ARG:HG2	1:A:373:TYR:HB3	1.99	0.44
1:A:350:THR:HG23	1:A:439:LYS:HB3	1.99	0.44
1:B:292:ARG:NE	1:B:292:ARG:H	2.15	0.44
1:B:318:GLU:HA	1:B:337:SER:HB3	2.00	0.43
1:A:368:LEU:HD13	1:A:407:TYR:CZ	2.53	0.43
1:B:334:LYS:HA	1:B:334:LYS:HD3	1.83	0.43
1:A:242:LEU:HD11	1:A:259:VAL:CG1	2.49	0.43
1:B:235:LEU:HD21	2:C:117:ALA:HB3	2.01	0.43
2:C:31:GLY:O	2:C:32:ALA:HB3	2.18	0.43
1:A:351:LEU:HB2	1:A:366:THR:HB	2.01	0.42
1:A:406:LEU:C	1:A:406:LEU:HD12	2.43	0.42
1:B:318:GLU:OE1	1:B:335:THR:HG21	2.19	0.42
1:B:346:PRO:HB3	1:B:372:PHE:HB3	2.02	0.42
2:C:21:GLU:OE2	2:C:63:VAL:HG23	2.19	0.42
1:A:237:GLY:HA2	2:C:134:HIS:CD2	2.54	0.41
2:C:146:LEU:HD12	2:C:172:ILE:HG22	2.00	0.41
1:B:397:VAL:HB	1:B:405:PHE:CE2	2.55	0.41
2:C:108:LEU:HD12	2:C:139:PHE:HD2	1.85	0.41
1:A:390:ASN:HD22	1:A:390:ASN:HA	1.73	0.41
2:C:113:TRP:CH2	2:C:114:LYS:HE2	2.57	0.40
1:B:296:TYR:C	1:B:298:SER:H	2.29	0.40
2:C:38:GLN:HE21	2:C:38:GLN:HB3	1.70	0.40
1:A:414:LYS:HB3	1:A:414:LYS:HE2	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/225 (92%)	202 (98%)	4 (2%)	1 (0%)	24	37
1	B	210/225 (93%)	207 (99%)	3 (1%)	0	100	100
2	C	161/204 (79%)	157 (98%)	4 (2%)	0	100	100
All	All	578/654 (88%)	566 (98%)	11 (2%)	1 (0%)	43	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	237	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/208 (93%)	183 (94%)	11 (6%)	18	33
1	B	197/208 (95%)	184 (93%)	13 (7%)	15	26
2	C	148/183 (81%)	136 (92%)	12 (8%)	11	18
All	All	539/599 (90%)	503 (93%)	36 (7%)	15	26

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	269	GLU
1	A	282	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	294	GLU
1	A	302	VAL
1	A	307	THR
1	A	309	LEU
1	A	311	GLN
1	A	342	GLN
1	A	344	ARG
1	A	362	GLN
1	A	437	THR
1	B	242	LEU
1	B	262	VAL
1	B	264	VAL
1	B	272	GLU
1	B	292	ARG
1	B	295	GLN
1	B	302	VAL
1	B	340	LYS
1	B	358	LEU
1	B	430	GLU
1	B	439	LYS
1	B	441	LEU
1	B	442	SER
2	C	15	GLN
2	C	38	GLN
2	C	48	LEU
2	C	128	LYS
2	C	130	ARG
2	C	137	SER
2	C	146	LEU
2	C	149	SER
2	C	157	LEU
2	C	161	LYS
2	C	164	SER
2	C	167	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	268	HIS
1	A	276	ASN
1	A	311	GLN
1	A	315	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	342	GLN
1	A	362	GLN
1	A	390	ASN
1	A	421	ASN
1	A	429	HIS
1	A	438	GLN
1	B	342	GLN
1	B	386	GLN
1	B	390	ASN
1	B	418	GLN
1	B	429	HIS
1	B	435	HIS
2	C	38	GLN
2	C	74	GLN
2	C	119	HIS
2	C	134	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

26 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	D	1	3,1	14,14,15	0.61	0	17,19,21	1.13	1 (5%)
3	NAG	D	2	3	14,14,15	0.71	0	17,19,21	1.48	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BMA	D	3	3	11,11,12	0.22	0	15,15,17	0.99	1 (6%)
3	MAN	D	4	3	11,11,12	0.58	0	15,15,17	0.85	0
3	NAG	D	5	3	14,14,15	0.56	0	17,19,21	1.06	0
3	MAN	D	6	3	11,11,12	0.71	0	15,15,17	1.04	0
3	NAG	D	7	3	14,14,15	0.56	0	17,19,21	1.02	2 (11%)
3	NAG	D	8	3	14,14,15	0.57	0	17,19,21	1.15	2 (11%)
3	NAG	E	1	3,1	14,14,15	0.56	0	17,19,21	1.49	2 (11%)
3	NAG	E	2	3	14,14,15	0.61	0	17,19,21	1.51	3 (17%)
3	BMA	E	3	3	11,11,12	0.62	0	15,15,17	2.58	7 (46%)
3	MAN	E	4	3	11,11,12	0.82	0	15,15,17	1.36	2 (13%)
3	NAG	E	5	3	14,14,15	0.52	0	17,19,21	0.95	1 (5%)
3	MAN	E	6	3	11,11,12	0.71	0	15,15,17	0.88	0
3	NAG	E	7	3	14,14,15	0.54	0	17,19,21	1.64	2 (11%)
3	NAG	E	8	3	14,14,15	0.54	0	17,19,21	1.36	1 (5%)
4	NAG	F	1	2,4	14,14,15	0.86	0	17,19,21	3.34	7 (41%)
4	NAG	F	2	4	14,14,15	0.76	1 (7%)	17,19,21	0.68	0
4	BMA	F	3	4	11,11,12	0.63	0	15,15,17	2.70	5 (33%)
4	MAN	F	4	4	11,11,12	0.83	0	15,15,17	2.82	3 (20%)
4	MAN	F	5	4	11,11,12	0.58	0	15,15,17	1.04	1 (6%)
4	MAN	F	6	4	11,11,12	0.80	0	15,15,17	1.03	1 (6%)
4	MAN	F	7	4	11,11,12	0.62	0	15,15,17	1.20	1 (6%)
4	MAN	F	8	4	11,11,12	0.82	0	15,15,17	0.83	0
5	NAG	G	1	5,2	14,14,15	0.50	0	17,19,21	1.29	2 (11%)
5	NAG	G	2	5	14,14,15	0.52	0	17,19,21	1.51	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	1/2/19/22	0/1/1/1
3	NAG	D	5	3	-	2/6/23/26	0/1/1/1
3	MAN	D	6	3	-	0/2/19/22	0/1/1/1
3	NAG	D	7	3	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	8	3	-	2/6/23/26	0/1/1/1
3	NAG	E	1	3,1	1/1/5/7	2/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1
3	BMA	E	3	3	-	0/2/19/22	0/1/1/1
3	MAN	E	4	3	1/1/4/5	1/2/19/22	0/1/1/1
3	NAG	E	5	3	-	2/6/23/26	0/1/1/1
3	MAN	E	6	3	-	0/2/19/22	0/1/1/1
3	NAG	E	7	3	-	3/6/23/26	0/1/1/1
3	NAG	E	8	3	-	4/6/23/26	0/1/1/1
4	NAG	F	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	F	2	4	-	0/6/23/26	0/1/1/1
4	BMA	F	3	4	-	2/2/19/22	0/1/1/1
4	MAN	F	4	4	1/1/4/5	2/2/19/22	0/1/1/1
4	MAN	F	5	4	-	2/2/19/22	0/1/1/1
4	MAN	F	6	4	1/1/4/5	2/2/19/22	0/1/1/1
4	MAN	F	7	4	1/1/4/5	2/2/19/22	0/1/1/1
4	MAN	F	8	4	1/1/4/5	2/2/19/22	0/1/1/1
5	NAG	G	1	5,2	-	2/6/23/26	0/1/1/1
5	NAG	G	2	5	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	2	NAG	O5-C1	-2.10	1.40	1.43

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	4	MAN	C1-C2-C3	-8.18	97.73	109.64
4	F	1	NAG	C1-C2-N2	-7.83	98.10	110.43
4	F	1	NAG	C2-N2-C7	7.14	132.47	122.90
4	F	3	BMA	C1-O5-C5	5.86	120.05	112.19
3	E	3	BMA	C3-C4-C5	-5.78	99.75	110.23
4	F	4	MAN	C1-O5-C5	-5.51	104.80	112.19
4	F	3	BMA	O5-C5-C6	-5.12	97.70	107.66
4	F	3	BMA	O3-C3-C2	4.95	120.16	110.05
3	D	2	NAG	C1-O5-C5	4.84	118.67	112.19
3	E	1	NAG	C1-O5-C5	4.80	118.62	112.19
3	E	3	BMA	O3-C3-C2	4.31	118.85	110.05

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	1	NAG	C4-C3-C2	4.19	117.16	111.02
4	F	1	NAG	C8-C7-N2	-4.04	109.41	116.12
3	E	7	NAG	C2-N2-C7	4.00	128.25	122.90
3	E	8	NAG	C1-O5-C5	3.83	117.32	112.19
3	E	7	NAG	C1-O5-C5	3.79	117.26	112.19
5	G	1	NAG	C1-O5-C5	3.77	117.23	112.19
4	F	7	MAN	C1-O5-C5	3.72	117.17	112.19
5	G	2	NAG	C1-O5-C5	3.61	117.03	112.19
4	F	1	NAG	C1-O5-C5	3.61	117.03	112.19
3	E	3	BMA	O4-C4-C3	3.59	118.84	110.38
4	F	1	NAG	C3-C4-C5	-3.48	103.93	110.23
3	E	2	NAG	C2-N2-C7	3.38	127.43	122.90
4	F	1	NAG	O7-C7-N2	3.14	127.52	121.98
3	E	3	BMA	O4-C4-C5	-3.11	101.66	109.32
3	E	3	BMA	O3-C3-C4	2.97	117.38	110.38
4	F	4	MAN	C3-C4-C5	2.89	115.48	110.23
4	F	5	MAN	C1-O5-C5	2.85	116.01	112.19
3	E	4	MAN	C3-C4-C5	2.77	115.26	110.23
3	E	3	BMA	C1-O5-C5	2.70	115.81	112.19
4	F	3	BMA	O3-C3-C4	-2.68	104.07	110.38
3	E	2	NAG	C1-O5-C5	2.63	115.71	112.19
4	F	6	MAN	C1-O5-C5	-2.62	108.68	112.19
3	E	4	MAN	C1-C2-C3	-2.59	105.88	109.64
3	D	8	NAG	O7-C7-N2	2.56	126.51	121.98
5	G	2	NAG	C2-N2-C7	2.56	126.33	122.90
3	D	7	NAG	C1-O5-C5	2.50	115.54	112.19
3	D	1	NAG	C1-O5-C5	2.44	115.46	112.19
5	G	2	NAG	O5-C5-C6	2.38	112.30	107.66
3	E	2	NAG	O4-C4-C3	2.26	115.69	110.38
3	D	3	BMA	O4-C4-C3	-2.23	105.12	110.38
5	G	1	NAG	C6-C5-C4	-2.21	107.60	113.02
3	D	8	NAG	C1-O5-C5	2.20	115.14	112.19
3	E	5	NAG	O5-C1-C2	-2.20	107.89	111.29
3	D	2	NAG	O4-C4-C5	-2.15	104.04	109.32
3	E	3	BMA	O5-C5-C6	2.08	111.71	107.66
3	E	1	NAG	C3-C4-C5	-2.07	106.48	110.23
3	D	7	NAG	O7-C7-N2	2.02	125.56	121.98
4	F	3	BMA	C3-C4-C5	2.01	113.88	110.23

All (6) chirality outliers are listed below:

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom
-----	-------	-----	------	------

Mol	Chain	Res	Type	Atom
3	E	1	NAG	C1
3	E	4	MAN	C1
4	F	4	MAN	C1
4	F	6	MAN	C1
4	F	7	MAN	C1
4	F	8	MAN	C1

All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	8	MAN	O5-C5-C6-O6
3	E	7	NAG	O5-C5-C6-O6
5	G	2	NAG	O5-C5-C6-O6
3	D	7	NAG	O5-C5-C6-O6
3	E	1	NAG	O5-C5-C6-O6
4	F	8	MAN	C4-C5-C6-O6
5	G	2	NAG	C4-C5-C6-O6
3	E	8	NAG	O5-C5-C6-O6
4	F	4	MAN	O5-C5-C6-O6
4	F	7	MAN	O5-C5-C6-O6
3	E	7	NAG	C4-C5-C6-O6
3	E	8	NAG	C4-C5-C6-O6
5	G	1	NAG	C4-C5-C6-O6
4	F	3	BMA	O5-C5-C6-O6
3	E	1	NAG	C4-C5-C6-O6
5	G	1	NAG	O5-C5-C6-O6
3	D	7	NAG	C4-C5-C6-O6
4	F	6	MAN	O5-C5-C6-O6
3	D	5	NAG	C8-C7-N2-C2
3	D	5	NAG	O7-C7-N2-C2
3	E	8	NAG	C8-C7-N2-C2
3	E	8	NAG	O7-C7-N2-C2
3	D	8	NAG	O5-C5-C6-O6
3	E	5	NAG	C4-C5-C6-O6
3	E	5	NAG	O5-C5-C6-O6
4	F	4	MAN	C4-C5-C6-O6
4	F	3	BMA	C4-C5-C6-O6
4	F	5	MAN	C4-C5-C6-O6
4	F	6	MAN	C4-C5-C6-O6
3	D	8	NAG	C4-C5-C6-O6
3	D	4	MAN	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	F	7	MAN	C4-C5-C6-O6
3	E	7	NAG	C3-C2-N2-C7
4	F	5	MAN	O5-C5-C6-O6
3	E	4	MAN	O5-C5-C6-O6


There are no ring outliers.


5 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	8	NAG	1	0
4	F	6	MAN	1	0
4	F	3	BMA	1	0
3	E	5	NAG	1	0
4	F	1	NAG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	MLI	A	501	-	6,6,6	1.21	0	7,7,7	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MLI	A	501	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	501	MLI	C3-C1-C2-O7
6	A	501	MLI	C3-C1-C2-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	209/225 (92%)	-0.27	2 (0%) 79 76	35, 56, 101, 118	0
1	B	212/225 (94%)	0.13	4 (1%) 66 62	47, 74, 96, 108	0
2	C	165/204 (80%)	-0.23	4 (2%) 59 55	39, 53, 82, 102	0
All	All	586/654 (89%)	-0.12	10 (1%) 69 65	35, 61, 95, 118	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	5	LEU	4.4
2	C	32	ALA	3.5
2	C	38	GLN	3.5
1	B	232	PRO	3.2
1	B	443	LEU	3.1
1	B	253	ILE	2.9
1	A	236	GLY	2.8
1	B	422	VAL	2.6
1	A	444	SER	2.5
2	C	174	GLN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MAN	F	7	11/12	0.54	0.14	72,80,82,83	0
4	MAN	F	8	11/12	0.60	0.15	82,87,90,90	0
3	NAG	D	5	14/15	0.77	0.12	60,67,71,72	0
4	MAN	F	5	11/12	0.79	0.13	50,54,56,56	0
4	MAN	F	6	11/12	0.80	0.11	62,70,83,85	0
3	NAG	E	8	14/15	0.82	0.12	72,80,83,92	0
5	NAG	G	2	14/15	0.83	0.10	50,55,60,63	0
3	NAG	E	7	14/15	0.85	0.12	63,71,78,80	0
4	BMA	F	3	11/12	0.87	0.08	38,44,51,58	0
3	MAN	D	4	11/12	0.88	0.09	51,56,63,64	0
3	NAG	E	5	14/15	0.89	0.09	65,70,75,77	0
4	MAN	F	4	11/12	0.90	0.09	43,49,52,56	0
3	MAN	E	4	11/12	0.90	0.08	59,62,65,65	0
3	NAG	E	1	14/15	0.91	0.09	40,43,47,50	0
3	NAG	D	8	14/15	0.91	0.09	50,54,64,67	0
3	BMA	E	3	11/12	0.93	0.07	52,55,62,68	0
4	NAG	F	1	14/15	0.93	0.07	28,31,33,34	0
5	NAG	G	1	14/15	0.93	0.07	33,37,46,47	0
3	NAG	D	7	14/15	0.93	0.07	40,50,56,58	0
3	MAN	E	6	11/12	0.94	0.05	56,58,64,65	0
4	NAG	F	2	14/15	0.95	0.07	30,32,35,37	0
3	NAG	E	2	14/15	0.95	0.07	40,42,47,51	0
3	MAN	D	6	11/12	0.95	0.06	39,41,45,45	0
3	NAG	D	1	14/15	0.96	0.07	23,28,30,32	0
3	BMA	D	3	11/12	0.97	0.05	35,37,42,48	0
3	NAG	D	2	14/15	0.97	0.05	29,32,35,36	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	MLI	A	501	7/7	0.82	0.13	59,60,67,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.