



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:20 AM UTC

PDB ID : 5SGX / pdb_00005sgx
Title : CRYSTAL STRUCTURE OF HUMAN PHOSPHODIESTERASE 10 IN COMPLEX WITH N4(c1nn(c(n1)CCc3nn2c(ncc(C)c2n3)C)C)CCC[C@@H]4C(F)(F)F, micromolar IC50=0.027384
Authors : Joseph, C.; Benz, J.; Flohr, A.; Rudolph, M.G.
Deposited on : 2022-02-01
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	0% 82% 9% 9%
1	C	343	0% 85% 6% 8%
2	B	343	2% 80% 12% 8%
2	D	343	2% 84% 7% 8%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11221 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	0	6	0
			2581	1650	438	466	27			
1	C	316	Total	C	N	O	S	0	4	0
			2579	1648	438	469	24			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	447	GLY	-	expression tag	UNP Q9Y233
A	448	SER	-	expression tag	UNP Q9Y233
C	447	GLY	-	expression tag	UNP Q9Y233
C	448	SER	-	expression tag	UNP Q9Y233

- Molecule 2 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	315	Total	C	N	O	S	0	5	0
			2581	1649	434	470	28			
2	D	314	Total	C	N	O	S	0	5	0
			2572	1644	433	468	27			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	447	GLY	-	expression tag	UNP Q9Y233
B	448	SER	-	expression tag	UNP Q9Y233
D	447	GLY	-	expression tag	UNP Q9Y233
D	448	SER	-	expression tag	UNP Q9Y233


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		

- Molecule 5 is (4S)-5,8-dimethyl-2-(2-{1-methyl-3-[(2R)-2-(trifluoromethyl)pyrrolidin-1-yl]-1H-1,2,4-triazol-5-yl}ethyl)[1,2,4]triazolo[1,5-c]pyrimidine (CCD ID: IZH) (formula: C₁₇H₂₁F₃N₈) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	F	N	0	0
			28	17	3	8		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	F	N	0	0
			28	17	3	8		
5	C	1	Total	C	F	N	0	0
			28	17	3	8		
5	D	1	Total	C	F	N	0	0
			28	17	3	8		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	199	Total	O	0	0
			199	199		
6	B	223	Total	O	0	0
			223	223		
6	C	214	Total	O	0	0
			214	214		
6	D	152	Total	O	0	0
			152	152		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A




- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 2: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 2: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A





4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	135.60Å 135.60Å 235.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.72 – 1.93 43.72 – 1.93	Depositor EDS
% Data completeness (in resolution range)	97.0 (43.72-1.93) 97.0 (43.72-1.93)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.168 , 0.216 0.176 , 0.220	Depositor DCC
R_{free} test set	6085 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	31.9	Xtriage
Anisotropy	0.128	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11221	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CME, IZH, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.20	4/2640 (0.2%)	1.32	6/3569 (0.2%)
1	C	1.16	2/2642 (0.1%)	1.35	1/3575 (0.0%)
2	B	1.17	4/2626 (0.2%)	1.39	9/3551 (0.3%)
2	D	1.16	1/2617 (0.0%)	1.37	8/3540 (0.2%)
All	All	1.17	11/10525 (0.1%)	1.36	24/14235 (0.2%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	677	SER	C-O	6.66	1.32	1.24
1	A	730	TYR	C-O	6.51	1.31	1.24
2	D	535	HIS	CE1-NE2	6.33	1.38	1.32
2	B	567	HIS	CE1-NE2	5.66	1.38	1.32
1	C	710	PRO	C-O	-5.61	1.17	1.23
1	C	595	HIS	CE1-NE2	5.51	1.38	1.32
2	B	739	THR	C-O	-5.45	1.17	1.24
1	A	695	GLU	C-O	5.40	1.30	1.24
1	A	567	HIS	CE1-NE2	5.34	1.37	1.32
2	B	549	THR	C-O	5.11	1.30	1.23
1	A	584	ALA	C-O	-5.03	1.17	1.24

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	612	SER	CA-C-N	6.85	129.76	120.38
2	B	612	SER	C-N-CA	6.85	129.76	120.38
1	A	609	ASN	CA-CB-CG	-6.58	106.02	112.60
2	B	535	HIS	CA-CB-CG	-6.13	107.67	113.80
2	D	535	HIS	CA-CB-CG	-6.10	107.70	113.80
1	A	588	THR	CA-CB-OG1	-5.95	100.68	109.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	639	PHE	CA-C-N	5.91	126.54	119.98
2	B	639	PHE	C-N-CA	5.91	126.54	119.98
2	B	504	GLU	CB-CA-C	5.85	119.38	109.84
2	D	746	PRO	N-CA-C	5.80	117.77	110.70
1	C	722	VAL	N-CA-CB	5.62	114.04	110.50
1	A	553	ARG	CA-C-N	5.59	127.71	120.44
1	A	553	ARG	C-N-CA	5.59	127.71	120.44
2	D	466	VAL	CA-C-O	-5.56	115.49	121.27
2	D	458	GLY	CA-C-N	5.55	129.99	120.72
2	D	458	GLY	C-N-CA	5.55	129.99	120.72
2	B	735	ILE	N-CA-CB	5.47	114.72	110.45
1	A	609	ASN	N-CA-CB	5.44	118.55	110.49
2	D	457	GLN	CA-C-N	5.44	127.20	120.34
2	D	457	GLN	C-N-CA	5.44	127.20	120.34
2	B	466	VAL	CA-C-O	-5.22	115.84	121.27
2	D	593	GLN	N-CA-C	-5.15	105.74	111.36
1	A	463	THR	CA-CB-OG1	-5.09	101.97	109.60
2	B	541	LEU	CA-C-O	-5.04	115.53	120.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2581	0	2567	17	0
1	C	2579	0	2547	13	0
2	B	2581	0	2552	18	0
2	D	2572	0	2540	9	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	1	0	0	0	0
5	A	28	0	0	0	0
5	B	28	0	0	0	0
5	C	28	0	0	1	0
5	D	28	0	0	0	0
6	A	199	0	0	5	0
6	B	223	0	0	3	0
6	C	214	0	0	4	0
6	D	152	0	0	1	0
All	All	11221	0	10206	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:469:CYS:SG	6:C:1079:HOH:O	2.17	1.01
2:B:509:CME:HB2	6:B:1086:HOH:O	1.81	0.80
1:A:509[B]:CME:HB2	6:A:1053:HOH:O	1.86	0.75
1:C:624[B]:GLU:OE2	1:C:627:ARG:NH2	2.16	0.75
1:A:509[A]:CME:HB2	6:A:1053:HOH:O	1.87	0.74
1:A:461:GLN:HG2	6:A:1074:HOH:O	1.91	0.70
1:A:646:GLU:HG3	6:A:959:HOH:O	1.94	0.68
2:B:727:LEU:HD12	2:B:759:LEU:HD11	1.79	0.64
1:A:727:LEU:HD21	1:A:763:GLU:HG3	1.82	0.61
1:A:497:SER:OG	1:A:557:LEU:HD11	2.00	0.61
2:D:767:ARG:HE	2:D:767:ARG:HA	1.65	0.60
2:B:576:GLN:HG2	6:B:1059:HOH:O	2.06	0.56
2:B:727:LEU:HD12	2:B:759:LEU:CD1	2.36	0.56
2:D:492:TYR:CZ	2:D:496:ARG:HD2	2.40	0.55
2:B:497:SER:O	2:B:553:ARG:HD2	2.07	0.55
1:C:624[B]:GLU:HG2	6:C:982:HOH:O	2.07	0.54
1:A:497:SER:O	1:A:553:ARG:HD2	2.07	0.54
1:C:700:GLY:HA3	1:C:714:MET:O	2.08	0.53
2:B:492:TYR:CZ	2:B:496:ARG:HD2	2.43	0.53
2:B:498:CYS:SG	2:B:554:LYS:HG3	2.49	0.52
2:B:523:PRO:HD2	2:B:695:GLU:HG2	1.91	0.52
2:D:624[A]:GLU:OE2	2:D:627:ARG:NH2	2.29	0.51
2:D:740:THR:HA	2:D:743:GLN:HE21	1.75	0.50
1:A:469:CYS:SG	6:A:1057:HOH:O	1.98	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:642[A]:ARG:NE	1:A:740:THR:OG1	2.46	0.48
1:A:735:ILE:HB	1:A:736:PRO:HD3	1.96	0.48
2:B:700:GLY:HA3	2:B:714:MET:O	2.14	0.47
2:B:497:SER:OG	2:B:557:LEU:HD11	2.14	0.47
1:C:517:LYS:HE3	6:C:943:HOH:O	2.14	0.47
1:C:638:TYR:CD1	1:C:671:THR:HG21	2.50	0.47
1:A:642[A]:ARG:NH1	1:A:740:THR:OG1	2.49	0.46
2:B:727:LEU:HD11	6:B:1027:HOH:O	2.17	0.45
1:A:461:GLN:HG3	1:A:462:PHE:O	2.16	0.44
1:A:496:ARG:NH1	1:A:538:TYR:OH	2.47	0.44
1:C:498:CYS:SG	1:C:554:LYS:HG3	2.57	0.44
2:D:743:GLN:HG2	6:D:1034:HOH:O	2.17	0.44
1:C:693:TYR:OH	5:C:803:IZH:N4	2.51	0.44
1:C:716:ARG:O	1:C:719:LYS:HE2	2.18	0.43
1:C:592:GLU:HA	1:C:595:HIS:CD2	2.53	0.43
2:B:624[A]:GLU:OE2	2:B:627:ARG:NH2	2.34	0.43
1:A:722:VAL:N	1:A:723:PRO:CD	2.81	0.43
2:B:727:LEU:CD1	2:B:759:LEU:HD11	2.46	0.43
1:C:492:TYR:CZ	1:C:496:ARG:HD2	2.54	0.43
1:A:700:GLY:HA3	1:A:714:MET:O	2.19	0.42
1:A:497:SER:HA	1:A:542:GLN:NE2	2.34	0.42
1:A:681:LEU:HB3	1:A:683:PRO:HD2	2.02	0.42
1:C:697:TRP:CH2	1:C:719:LYS:HG2	2.55	0.42
2:D:533:VAL:HG13	2:D:673:CYS:HB3	2.01	0.42
2:D:767:ARG:HA	2:D:767:ARG:NE	2.32	0.42
2:B:551:LEU:HD23	2:B:551:LEU:HA	1.86	0.42
1:C:756[A]:ARG:HD2	6:C:1045:HOH:O	2.20	0.42
2:B:467:ARG:O	2:B:467:ARG:HG2	2.20	0.41
2:B:638:TYR:CD1	2:B:671:THR:HG21	2.56	0.41
2:B:696:PHE:CZ	2:B:713:MET:HE3	2.55	0.41
2:B:577:LYS:HD3	2:B:703:MET:HE1	2.03	0.41
2:D:497:SER:O	2:D:553:ARG:HD2	2.21	0.41
2:D:752:LEU:HD21	2:D:756:ARG:NH2	2.36	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/343 (92%)	309 (98%)	6 (2%)	0	100	100
1	C	317/343 (92%)	311 (98%)	6 (2%)	0	100	100
2	B	315/343 (92%)	308 (98%)	7 (2%)	0	100	100
2	D	314/343 (92%)	310 (99%)	4 (1%)	0	100	100
All	All	1261/1372 (92%)	1238 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/305 (94%)	282 (99%)	4 (1%)	59	52
1	C	285/305 (93%)	280 (98%)	5 (2%)	51	43
2	B	284/304 (93%)	277 (98%)	7 (2%)	42	29
2	D	282/304 (93%)	276 (98%)	6 (2%)	47	37
All	All	1137/1218 (93%)	1115 (98%)	22 (2%)	51	40

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	471[A]	GLU
1	A	471[B]	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	573	SER
1	A	681	LEU
2	B	613	THR
2	B	616	SER
2	B	642	ARG
2	B	709	GLN
2	B	720[A]	ASP
2	B	720[B]	ASP
2	B	749	GLU
1	C	506	GLU
1	C	613	THR
1	C	616	SER
1	C	719	LYS
1	C	722	VAL
2	D	459	LEU
2	D	500	THR
2	D	576	GLN
2	D	616	SER
2	D	621	GLN
2	D	767	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	461	GLN
1	A	542	GLN
1	A	598	GLN
1	A	604	GLN
1	A	709	GLN
1	A	726	GLN
1	A	761	GLN
2	B	484	ASN
2	B	542	GLN
2	B	545	HIS
2	B	604	GLN
2	B	650	GLN
2	B	724	GLN
2	B	726	GLN
2	B	761	GLN
1	C	495	HIS
1	C	542	GLN
1	C	604	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	659	GLN
1	C	709	GLN
1	C	726	GLN
1	C	743	GLN
2	D	484	ASN
2	D	545	HIS
2	D	604	GLN
2	D	659	GLN
2	D	726	GLN
2	D	743	GLN
2	D	761	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

9 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CME	C	509	1	8,9,10	0.74	0	6,9,11	0.83	0
2	CME	D	469[B]	-	8,9,10	0.39	0	6,9,11	1.00	0
2	CME	B	469[A]	-	8,9,10	0.56	0	6,9,11	0.87	0
2	CME	D	509	2	8,9,10	0.62	0	6,9,11	0.85	0
1	CME	A	509[A]	-	8,9,10	0.38	0	6,9,11	0.68	0
2	CME	B	469[B]	-	8,9,10	0.63	0	6,9,11	0.55	0
2	CME	B	509	2	8,9,10	0.64	0	6,9,11	1.19	1 (16%)
1	CME	A	509[B]	-	8,9,10	0.31	0	6,9,11	0.83	0
2	CME	D	469[A]	-	8,9,10	0.41	0	6,9,11	1.22	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	C	509	1	-	2/5/8/10	-
2	CME	D	469[B]	-	-	2/5/8/10	-
2	CME	B	469[A]	-	-	1/5/8/10	-
2	CME	D	509	2	-	1/5/8/10	-
1	CME	A	509[A]	-	-	2/5/8/10	-
2	CME	B	469[B]	-	-	3/5/8/10	-
2	CME	B	509	2	-	2/5/8/10	-
1	CME	A	509[B]	-	-	3/5/8/10	-
2	CME	D	469[A]	-	-	3/5/8/10	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	469[A]	CME	CB-SG-SD	-2.39	97.68	103.86
2	B	509	CME	CZ-CE-SD	-2.05	106.55	113.39

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	509[A]	CME	SD-CE-CZ-OH
2	B	509	CME	SD-CE-CZ-OH
1	C	509	CME	SD-CE-CZ-OH
2	D	469[A]	CME	SD-CE-CZ-OH
2	D	469[B]	CME	SD-CE-CZ-OH
2	B	469[B]	CME	CE-SD-SG-CB
2	B	469[A]	CME	SD-CE-CZ-OH
1	A	509[B]	CME	SD-CE-CZ-OH
2	B	469[B]	CME	N-CA-CB-SG
2	D	469[A]	CME	N-CA-CB-SG
1	A	509[A]	CME	CZ-CE-SD-SG
2	D	469[A]	CME	CZ-CE-SD-SG
1	A	509[B]	CME	CE-SD-SG-CB
1	A	509[B]	CME	CZ-CE-SD-SG
2	B	509	CME	CZ-CE-SD-SG
1	C	509	CME	CZ-CE-SD-SG
2	D	469[B]	CME	CZ-CE-SD-SG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	469[B]	CME	SD-CE-CZ-OH
2	D	509	CME	SD-CE-CZ-OH

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	509[A]	CME	1	0
2	B	509	CME	1	0
1	A	509[B]	CME	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	IZH	A	803	-	30,31,31	2.32	11 (36%)	30,47,47	1.71	5 (16%)
5	IZH	B	803	-	30,31,31	2.08	9 (30%)	30,47,47	2.22	7 (23%)
5	IZH	D	803	-	30,31,31	2.13	9 (30%)	30,47,47	2.12	9 (30%)
5	IZH	C	803	-	30,31,31	2.75	9 (30%)	30,47,47	1.23	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	IZH	A	803	-	-	1/12/25/25	0/4/4/4
5	IZH	B	803	-	-	1/12/25/25	0/4/4/4
5	IZH	D	803	-	-	0/12/25/25	0/4/4/4
5	IZH	C	803	-	-	0/12/25/25	0/4/4/4

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	803	IZH	C3-N7	9.97	1.43	1.34
5	A	803	IZH	C25-C21	7.68	1.67	1.52
5	B	803	IZH	C25-C21	6.68	1.65	1.52
5	D	803	IZH	C25-C21	6.45	1.64	1.52
5	C	803	IZH	C5-N6	5.67	1.40	1.32
5	C	803	IZH	C25-C21	5.60	1.63	1.52
5	D	803	IZH	C5-N6	4.86	1.39	1.32
5	A	803	IZH	C24-N1	-4.10	1.39	1.47
5	D	803	IZH	C14-N11	3.89	1.36	1.30
5	A	803	IZH	C3-N7	3.85	1.37	1.34
5	C	803	IZH	C3-N4	3.77	1.39	1.32
5	B	803	IZH	C12-C13	-3.56	1.29	1.36
5	B	803	IZH	C3-N4	3.49	1.38	1.32
5	C	803	IZH	C2-N17	3.19	1.42	1.37
5	A	803	IZH	C21-N1	-3.15	1.43	1.47
5	D	803	IZH	C3-N4	3.13	1.38	1.32
5	A	803	IZH	C3-N4	2.88	1.37	1.32
5	B	803	IZH	C8-N7	2.85	1.51	1.46
5	B	803	IZH	N15-N18	-2.82	1.33	1.38
5	A	803	IZH	F28-C25	-2.81	1.26	1.33
5	C	803	IZH	F28-C25	2.74	1.40	1.33
5	A	803	IZH	C14-N11	-2.74	1.25	1.30
5	A	803	IZH	C13-C16	-2.62	1.36	1.42
5	B	803	IZH	C16-N17	-2.61	1.27	1.33
5	D	803	IZH	F28-C25	2.45	1.39	1.33
5	B	803	IZH	C9-C3	2.43	1.52	1.49
5	D	803	IZH	C19-C13	-2.43	1.44	1.50
5	D	803	IZH	C3-N7	2.42	1.36	1.34
5	D	803	IZH	C16-N17	-2.41	1.28	1.33
5	C	803	IZH	C12-C13	-2.37	1.31	1.36
5	B	803	IZH	C14-N11	2.34	1.34	1.30
5	C	803	IZH	C2-N18	2.21	1.36	1.32
5	A	803	IZH	C12-C13	-2.21	1.32	1.36
5	A	803	IZH	C22-C21	-2.18	1.49	1.53
5	A	803	IZH	C12-N11	-2.17	1.31	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	803	IZH	C2-N18	-2.15	1.29	1.32
5	C	803	IZH	C19-C13	-2.10	1.45	1.50
5	B	803	IZH	C13-C16	2.05	1.47	1.42

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	803	IZH	C22-C21-C25	-6.44	106.68	113.33
5	A	803	IZH	C22-C21-C25	-5.95	107.18	113.33
5	B	803	IZH	N15-C16-N17	-5.94	104.69	110.13
5	D	803	IZH	C22-C21-C25	-5.84	107.30	113.33
5	D	803	IZH	C22-C21-N1	5.14	108.30	102.73
5	D	803	IZH	F27-C25-F26	-3.78	98.65	106.87
5	B	803	IZH	C22-C21-N1	3.63	106.67	102.73
5	A	803	IZH	N15-C16-N17	-3.62	106.83	110.13
5	B	803	IZH	C13-C16-N17	3.35	134.40	129.41
5	B	803	IZH	C10-C9-C3	-3.28	108.52	111.62
5	A	803	IZH	C13-C16-N17	3.27	134.27	129.41
5	D	803	IZH	F28-C25-F27	3.25	113.93	106.87
5	D	803	IZH	C13-C16-N17	2.92	133.75	129.41
5	D	803	IZH	N15-C16-N17	-2.71	107.65	110.13
5	C	803	IZH	C22-C21-N1	2.68	105.63	102.73
5	D	803	IZH	C10-C9-C3	-2.56	109.20	111.62
5	B	803	IZH	F28-C25-F26	2.49	112.29	106.87
5	C	803	IZH	C10-C9-C3	-2.40	109.35	111.62
5	C	803	IZH	C13-C16-N17	2.32	132.87	129.41
5	C	803	IZH	C21-N1-C5	-2.25	115.61	122.42
5	A	803	IZH	C22-C21-N1	2.17	105.08	102.73
5	D	803	IZH	C20-C14-N15	-2.16	114.85	120.05
5	D	803	IZH	C20-C14-N11	2.12	123.41	119.26
5	C	803	IZH	C8-N7-N6	2.08	122.40	119.29
5	B	803	IZH	C8-N7-N6	2.08	122.40	119.29
5	A	803	IZH	F26-C25-C21	-2.01	108.75	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	803	IZH	N7-C3-C9-C10
5	B	803	IZH	N7-C3-C9-C10

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	803	IZH	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand IZH B 803



Ligand IZH D 803





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/343 (90%)	-0.04	5 (1%) 70 77	18, 34, 61, 98	5 (1%)
1	C	315/343 (91%)	-0.06	5 (1%) 70 77	20, 34, 60, 104	4 (1%)
2	B	313/343 (91%)	-0.05	7 (2%) 62 69	20, 32, 59, 103	4 (1%)
2	D	312/343 (90%)	0.20	8 (2%) 57 64	26, 41, 67, 111	4 (1%)
All	All	1252/1372 (91%)	0.01	25 (1%) 65 72	18, 36, 63, 111	17 (1%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	456	TRP	9.5
1	C	456	TRP	6.9
2	B	457	GLN	5.5
2	D	459	LEU	5.2
2	B	459	LEU	5.0
2	B	771	THR	4.8
2	D	457	GLN	4.6
1	C	459	LEU	4.2
1	A	459	LEU	3.9
2	D	458	GLY	3.2
1	A	458	GLY	2.9
2	B	458	GLY	2.9
1	A	770	GLU	2.7
1	A	461	GLN	2.6
2	B	460	MET	2.4
1	C	458	GLY	2.4
2	D	500	THR	2.3
1	C	455	GLU	2.2
2	B	461	GLN	2.2
2	B	550	ASP	2.2
1	A	462	PHE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	462	PHE	2.1
2	D	769	GLU	2.1
2	D	460	MET	2.1
2	D	767	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CME	D	509	10/11	0.85	0.16	38,54,97,101	0
2	CME	B	469[B]	10/11	0.88	0.14	39,45,60,63	7
2	CME	B	469[A]	10/11	0.88	0.14	38,40,45,50	7
1	CME	A	509[B]	10/11	0.90	0.14	39,49,67,83	7
1	CME	C	509	10/11	0.90	0.13	37,41,76,78	0
1	CME	A	509[A]	10/11	0.90	0.14	39,48,62,73	7
2	CME	D	469[A]	10/11	0.92	0.15	40,54,63,71	7
2	CME	D	469[B]	10/11	0.92	0.15	40,42,44,45	7
2	CME	B	509	10/11	0.92	0.12	30,43,78,87	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	IZH	A	803	28/28	0.95	0.07	27,32,38,45	0
5	IZH	B	803	28/28	0.96	0.06	26,31,37,39	0
5	IZH	D	803	28/28	0.96	0.06	31,36,42,45	0
5	IZH	C	803	28/28	0.97	0.06	26,35,41,43	0
4	MG	A	802	1/1	0.99	0.02	26,26,26,26	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	B	802	1/1	0.99	0.02	23,23,23,23	0
4	MG	C	802	1/1	0.99	0.02	25,25,25,25	0
4	MG	D	802	1/1	0.99	0.02	34,34,34,34	0
3	ZN	A	801	1/1	1.00	0.01	29,29,29,29	0
3	ZN	B	801	1/1	1.00	0.01	27,27,27,27	0
3	ZN	C	801	1/1	1.00	0.02	28,28,28,28	0
3	ZN	D	801	1/1	1.00	0.03	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around IZH A 803:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IZH B 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around IZH D 803:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around IZH C 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.