



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 09:46 PM UTC

PDB ID : 7SGY / pdb_00007sgy
Title : Cannabis sativa bibenzyl synthase
Authors : Kimber, M.S.; Forrester, T.J.B.
Deposited on : 2021-10-07
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	427	
1	B	427	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6122 atoms, of which 8 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bibenzyl synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	383	Total	C	H	N	O	S	0	1	0
			2988	1894	4	512	562	16			
1	B	387	Total	C	H	N	O	S	0	0	0
			3011	1912	4	516	564	15			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	expression tag	UNP A0A803NJB5
A	-32	GLY	-	expression tag	UNP A0A803NJB5
A	-31	SER	-	expression tag	UNP A0A803NJB5
A	-30	SER	-	expression tag	UNP A0A803NJB5
A	-29	HIS	-	expression tag	UNP A0A803NJB5
A	-28	HIS	-	expression tag	UNP A0A803NJB5
A	-27	HIS	-	expression tag	UNP A0A803NJB5
A	-26	HIS	-	expression tag	UNP A0A803NJB5
A	-25	HIS	-	expression tag	UNP A0A803NJB5
A	-24	HIS	-	expression tag	UNP A0A803NJB5
A	-23	SER	-	expression tag	UNP A0A803NJB5
A	-22	SER	-	expression tag	UNP A0A803NJB5
A	-21	GLY	-	expression tag	UNP A0A803NJB5
A	-20	LEU	-	expression tag	UNP A0A803NJB5
A	-19	VAL	-	expression tag	UNP A0A803NJB5
A	-18	PRO	-	expression tag	UNP A0A803NJB5
A	-17	ARG	-	expression tag	UNP A0A803NJB5
A	-16	GLY	-	expression tag	UNP A0A803NJB5
A	-15	SER	-	expression tag	UNP A0A803NJB5
A	-14	HIS	-	expression tag	UNP A0A803NJB5
A	-13	MET	-	expression tag	UNP A0A803NJB5
A	-12	ALA	-	expression tag	UNP A0A803NJB5
A	-11	SER	-	expression tag	UNP A0A803NJB5
A	-10	MET	-	expression tag	UNP A0A803NJB5
A	-9	THR	-	expression tag	UNP A0A803NJB5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	GLY	-	expression tag	UNP A0A803NJB5
A	-7	GLY	-	expression tag	UNP A0A803NJB5
A	-6	GLN	-	expression tag	UNP A0A803NJB5
A	-5	GLN	-	expression tag	UNP A0A803NJB5
A	-4	MET	-	expression tag	UNP A0A803NJB5
A	-3	GLY	-	expression tag	UNP A0A803NJB5
A	-2	ARG	-	expression tag	UNP A0A803NJB5
A	-1	GLY	-	expression tag	UNP A0A803NJB5
A	0	SER	-	expression tag	UNP A0A803NJB5
A	191	VAL	ILE	conflict	UNP A0A803NJB5
A	254	ASP	HIS	conflict	UNP A0A803NJB5
A	389	ALA	SER	conflict	UNP A0A803NJB5
A	391	GLN	-	expression tag	UNP A0A803NJB5
A	392	THR	-	expression tag	UNP A0A803NJB5
A	393	SER	-	expression tag	UNP A0A803NJB5
B	-33	MET	-	expression tag	UNP A0A803NJB5
B	-32	GLY	-	expression tag	UNP A0A803NJB5
B	-31	SER	-	expression tag	UNP A0A803NJB5
B	-30	SER	-	expression tag	UNP A0A803NJB5
B	-29	HIS	-	expression tag	UNP A0A803NJB5
B	-28	HIS	-	expression tag	UNP A0A803NJB5
B	-27	HIS	-	expression tag	UNP A0A803NJB5
B	-26	HIS	-	expression tag	UNP A0A803NJB5
B	-25	HIS	-	expression tag	UNP A0A803NJB5
B	-24	HIS	-	expression tag	UNP A0A803NJB5
B	-23	SER	-	expression tag	UNP A0A803NJB5
B	-22	SER	-	expression tag	UNP A0A803NJB5
B	-21	GLY	-	expression tag	UNP A0A803NJB5
B	-20	LEU	-	expression tag	UNP A0A803NJB5
B	-19	VAL	-	expression tag	UNP A0A803NJB5
B	-18	PRO	-	expression tag	UNP A0A803NJB5
B	-17	ARG	-	expression tag	UNP A0A803NJB5
B	-16	GLY	-	expression tag	UNP A0A803NJB5
B	-15	SER	-	expression tag	UNP A0A803NJB5
B	-14	HIS	-	expression tag	UNP A0A803NJB5
B	-13	MET	-	expression tag	UNP A0A803NJB5
B	-12	ALA	-	expression tag	UNP A0A803NJB5
B	-11	SER	-	expression tag	UNP A0A803NJB5
B	-10	MET	-	expression tag	UNP A0A803NJB5
B	-9	THR	-	expression tag	UNP A0A803NJB5
B	-8	GLY	-	expression tag	UNP A0A803NJB5
B	-7	GLY	-	expression tag	UNP A0A803NJB5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	GLN	-	expression tag	UNP A0A803NJB5
B	-5	GLN	-	expression tag	UNP A0A803NJB5
B	-4	MET	-	expression tag	UNP A0A803NJB5
B	-3	GLY	-	expression tag	UNP A0A803NJB5
B	-2	ARG	-	expression tag	UNP A0A803NJB5
B	-1	GLY	-	expression tag	UNP A0A803NJB5
B	0	SER	-	expression tag	UNP A0A803NJB5
B	191	VAL	ILE	conflict	UNP A0A803NJB5
B	254	ASP	HIS	conflict	UNP A0A803NJB5
B	389	ALA	SER	conflict	UNP A0A803NJB5
B	391	GLN	-	expression tag	UNP A0A803NJB5
B	392	THR	-	expression tag	UNP A0A803NJB5
B	393	SER	-	expression tag	UNP A0A803NJB5

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	66	Total O 66 66	0	0
3	B	55	Total O 55 55	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.44Å 59.58Å 70.94Å 90.00° 107.70° 90.00°	Depositor
Resolution (Å)	44.69 – 2.50 44.69 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (44.69-2.50) 99.6 (44.69-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.184 , 0.228 0.184 , 0.227	Depositor DCC
R_{free} test set	2208 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	46.5	Xtriage
Anisotropy	0.681	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 68.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6122	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.34 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.8129e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/3038	0.28	0/4102
1	B	0.10	0/3063	0.28	0/4140
All	All	0.10	0/6101	0.28	0/8242

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2984	4	2980	88	0
1	B	3007	4	3012	92	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	66	0	0	3	0
3	B	55	0	0	7	0
All	All	6114	8	5992	170	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 170 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:THR:HG21	1:A:203:PRO:HD2	1.54	0.87
1:A:204:ASP:OD1	1:A:206:THR:N	2.09	0.86
1:B:40:ASP:OD2	1:B:55:LYS:NZ	2.08	0.85
1:A:286:LEU:HD21	1:A:374:PHE:HZ	1.39	0.85
1:B:291:ASN:HB3	1:B:292:PRO:CD	2.07	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/427 (89%)	365 (96%)	13 (3%)	1 (0%)	36	55
1	B	384/427 (90%)	367 (96%)	12 (3%)	5 (1%)	9	18
All	All	763/854 (89%)	732 (96%)	25 (3%)	6 (1%)	16	31

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	291	ASN
1	B	207	TYR
1	B	51	MET
1	B	292	PRO
1	A	203	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/354 (91%)	312 (97%)	9 (3%)	38	66
1	B	323/354 (91%)	311 (96%)	12 (4%)	30	57
All	All	644/708 (91%)	623 (97%)	21 (3%)	33	61

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	72	LEU
1	B	284	LYS
1	B	322	LEU
1	B	286	LEU
1	B	266	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	306	HIS
1	B	81	GLN
1	B	281	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSO	A	166	1	3,6,7	0.58	0	1,6,8	0.59	0
1	CSO	B	166	1	3,6,7	0.59	0	1,6,8	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	166	1	-	0/1/5/7	-
1	CSO	B	166	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	166	CSO	2	0
1	B	166	CSO	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	382/427 (89%)	-0.28	1 (0%) 90 87	24, 54, 106, 177	1 (0%)
1	B	386/427 (90%)	-0.30	0 100 100	35, 57, 97, 139	0
All	All	768/854 (89%)	-0.29	1 (0%) 92 90	24, 56, 99, 177	1 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	289	ALA	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	B	166	7/8	0.86	0.08	32,43,60,72	0
1	CSO	A	166	7/8	0.94	0.06	35,40,55,66	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	B	401	1/1	0.94	0.06	61,61,61,61	0
2	CL	A	401	1/1	0.98	0.06	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.