



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:57 PM UTC

PDB ID : 1SHN / pdb_00001shn
Title : Crystal structure of shrimp alkaline phosphatase with phosphate bound
Authors : de Backer, M.M.E.; McSweeney, S.; Lindley, P.F.; Hough, E.
Deposited on : 2004-02-26
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	% 84% 14% .
1	B	478	3% 81% 16% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	A	486	-	X	X	-
4	PO4	B	484	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7699 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alkaline phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	476	Total	C	N	O	S	0	0	0
			3731	2333	629	755	14			
1	B	476	Total	C	N	O	S	0	0	0
			3731	2333	629	755	14			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLU	-	cloning artifact	UNP Q9BHT8
A	2	GLU	-	cloning artifact	UNP Q9BHT8
A	3	ASP	-	cloning artifact	UNP Q9BHT8
A	184	ALA	ARG	conflict	UNP Q9BHT8
A	355	ALA	VAL	conflict	UNP Q9BHT8
B	1	GLU	-	cloning artifact	UNP Q9BHT8
B	2	GLU	-	cloning artifact	UNP Q9BHT8
B	3	ASP	-	cloning artifact	UNP Q9BHT8
B	184	ALA	ARG	conflict	UNP Q9BHT8
B	355	ALA	VAL	conflict	UNP Q9BHT8

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Zn	0	0
			3	3		
3	B	3	Total	Zn	0	0
			3	3		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is SULFATE ION (CCD ID: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	101	Total 101	O 101	0	0
6	B	82	Total 82	O 82	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alkaline phosphatase



- Molecule 1: alkaline phosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	170.68Å 170.68Å 83.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.64 – 2.15 39.64 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.3 (39.64-2.15) 96.2 (39.64-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.09 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.186 , 0.218 0.197 , 0.225	Depositor DCC
R_{free} test set	3357 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 28.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7699	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.78 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7362e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ZN, SO4, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.23	7/3811 (0.2%)	1.22	7/5174 (0.1%)
1	B	1.26	10/3811 (0.3%)	1.20	10/5174 (0.2%)
All	All	1.24	17/7622 (0.2%)	1.21	17/10348 (0.2%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	430	VAL	CA-CB	-9.36	1.44	1.54
1	B	117	TYR	CG-CD2	-6.71	1.25	1.39
1	A	271	THR	C-O	6.37	1.28	1.23
1	B	343	MET	SD-CE	-6.26	1.64	1.79
1	B	247	VAL	CA-C	-6.21	1.45	1.52
1	A	141	VAL	N-CA	-6.00	1.39	1.46
1	A	267	LEU	C-O	-5.85	1.17	1.24
1	B	113	THR	CA-CB	-5.85	1.46	1.54
1	B	117	TYR	CE1-CZ	-5.79	1.24	1.38
1	B	360	THR	CA-CB	-5.64	1.46	1.53
1	A	200	MET	SD-CE	-5.58	1.65	1.79
1	B	5	ALA	C-O	-5.36	1.17	1.24
1	A	339	MET	SD-CE	-5.34	1.66	1.79
1	A	271	THR	N-CA	5.33	1.49	1.46
1	B	246	TYR	N-CA	-5.27	1.39	1.46
1	B	290	MET	SD-CE	-5.18	1.66	1.79
1	A	360	THR	CA-CB	-5.11	1.46	1.52

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	22	LYS	CB-CG-CD	7.62	128.82	111.30
1	B	250	ARG	N-CA-C	7.33	118.91	111.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	76	THR	N-CA-C	7.23	122.27	113.38
1	A	215	ILE	N-CA-C	-6.35	101.39	109.58
1	A	476	ASP	CB-CA-C	-6.26	98.21	110.10
1	B	215	ILE	N-CA-C	-6.19	101.56	109.80
1	A	392	TYR	N-CA-C	5.77	118.61	109.50
1	A	104	ILE	N-CA-C	5.76	116.16	107.75
1	B	324	ARG	CG-CD-NE	-5.39	100.13	112.00
1	B	399	HIS	N-CA-C	5.37	116.94	107.61
1	B	22	LYS	CB-CA-C	-5.30	100.02	109.71
1	A	76	THR	N-CA-C	5.27	119.81	113.17
1	B	108	ALA	N-CA-C	5.26	119.44	113.19
1	A	357	HIS	CA-C-N	-5.09	117.90	122.43
1	A	357	HIS	C-N-CA	-5.09	117.90	122.43
1	B	22	LYS	CA-CB-CG	5.08	124.26	114.10
1	B	299	THR	OG1-CB-CG2	-5.05	99.21	109.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3731	0	3516	61	0
1	B	3731	0	3516	64	0
2	A	14	0	13	0	0
2	B	14	0	13	4	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
4	A	5	0	0	2	0
4	B	5	0	0	3	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	101	0	0	3	0
6	B	82	0	0	2	0
All	All	7699	0	7058	125	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (125) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:114:ASN:HD21	2:B:483:NAG:C1	1.27	1.45
1:A:339:MET:SD	1:A:343:MET:CE	2.40	1.09
1:A:339:MET:HG3	1:A:343:MET:HE3	1.32	1.06
1:A:339:MET:CG	1:A:343:MET:HE3	1.93	0.99
1:A:339:MET:SD	1:A:343:MET:HE2	2.04	0.97
1:B:313:ARG:HE	1:B:316:HIS:HD2	1.01	0.95
1:A:313:ARG:HE	1:A:316:HIS:HD2	1.06	0.95
1:B:114:ASN:HD21	2:B:483:NAG:C2	1.86	0.88
1:A:339:MET:CG	1:A:343:MET:CE	2.55	0.84
1:B:339:MET:HG3	1:B:343:MET:HE3	1.59	0.83
1:B:313:ARG:HE	1:B:316:HIS:CD2	1.93	0.83
1:A:339:MET:SD	1:A:343:MET:HE3	2.18	0.82
1:A:339:MET:HG3	1:A:343:MET:CE	2.09	0.82
1:A:86:SER:OG	4:A:486:PO4:O4	1.98	0.80
1:A:292:LYS:CB	1:A:343:MET:HE1	2.15	0.77
1:B:86:SER:OG	4:B:484:PO4:P	2.42	0.77
1:B:339:MET:SD	1:B:343:MET:CE	2.74	0.75
1:A:313:ARG:HE	1:A:316:HIS:CD2	1.99	0.73
1:B:339:MET:SD	1:B:343:MET:HE2	2.31	0.70
1:B:214:ASP:OD2	1:B:215:ILE:O	2.10	0.70
1:A:29:LYS:HE3	1:A:302:GLU:O	1.92	0.69
1:A:292:LYS:HB2	1:A:343:MET:HE1	1.75	0.69
4:B:484:PO4:O2	6:B:523:HOH:O	2.11	0.68
1:A:313:ARG:NE	1:A:316:HIS:HD2	1.88	0.66
1:A:215:ILE:O	1:A:216:GLU:HB2	1.98	0.64
1:B:146:ARG:H	1:B:272:HIS:HD2	1.46	0.64
1:B:369:ARG:O	1:B:370:ASN:HB2	1.97	0.64
1:B:359:HIS:HD2	6:B:505:HOH:O	1.83	0.61
1:A:359:HIS:HD2	6:A:488:HOH:O	1.84	0.61
1:B:339:MET:CG	1:B:343:MET:HE3	2.31	0.61
1:B:58:PHE:CD2	1:B:370:ASN:OD1	2.53	0.60
1:B:58:PHE:CG	1:B:370:ASN:OD1	2.54	0.60
1:B:339:MET:HG3	1:B:343:MET:CE	2.31	0.59
1:A:292:LYS:HG3	1:A:343:MET:CE	2.33	0.59
1:B:215:ILE:O	1:B:216:GLU:HB2	2.03	0.59
1:B:292:LYS:HA	1:B:343:MET:HE1	1.85	0.59
1:B:112:ARG:O	1:B:113:THR:HB	2.04	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:THR:OG1	1:B:318:HIS:HE1	1.87	0.57
1:A:203:GLY:N	1:A:271:THR:O	2.38	0.56
1:B:115:CYS:HB3	1:B:175:GLU:OE1	2.04	0.56
1:B:114:ASN:CG	2:B:483:NAG:C1	2.78	0.56
1:B:313:ARG:NE	1:B:316:HIS:HD2	1.87	0.56
1:B:292:LYS:CB	1:B:343:MET:HE1	2.37	0.55
1:A:208:PHE:CE1	1:A:221:GLY:CA	2.89	0.55
1:A:214:ASP:OD2	1:A:215:ILE:O	2.25	0.55
1:A:392:TYR:O	1:A:423:SER:HA	2.07	0.54
1:B:29:LYS:HE3	1:B:302:GLU:O	2.06	0.54
1:B:86:SER:OG	4:B:484:PO4:O4	2.26	0.54
1:A:146:ARG:H	1:A:272:HIS:HD2	1.56	0.53
1:A:112:ARG:O	1:A:113:THR:HB	2.08	0.53
1:B:146:ARG:H	1:B:272:HIS:CD2	2.27	0.53
1:A:215:ILE:HG13	6:A:511:HOH:O	2.08	0.53
1:A:119:LEU:HD11	1:A:179:ILE:HD11	1.92	0.52
1:A:473:THR:HG22	1:A:476:ASP:H	1.75	0.51
1:B:339:MET:CG	1:B:343:MET:CE	2.88	0.51
1:B:151:THR:HB	1:B:152:PRO:CD	2.41	0.51
1:A:315:ASP:OD1	1:A:359:HIS:HE1	1.94	0.50
1:B:315:ASP:OD1	1:B:359:HIS:HE1	1.94	0.50
1:A:194:LYS:O	1:A:237:LYS:NZ	2.44	0.50
1:A:86:SER:OG	4:A:486:PO4:P	2.69	0.50
1:B:112:ARG:O	1:B:113:THR:CB	2.60	0.49
1:A:292:LYS:HG3	1:A:343:MET:HE1	1.95	0.49
1:B:292:LYS:CA	1:B:343:MET:HE1	2.42	0.49
1:B:458:ILE:HB	1:B:459:PRO:HD3	1.95	0.49
1:A:292:LYS:CA	1:A:343:MET:HE1	2.42	0.49
1:A:26:LYS:NZ	1:A:347:GLU:OE1	2.35	0.48
1:A:280:ASP:OD1	1:A:282:GLU:HB2	2.13	0.48
1:A:292:LYS:CG	1:A:343:MET:HE1	2.42	0.48
1:B:392:TYR:O	1:B:423:SER:HA	2.14	0.48
1:A:370:ASN:N	1:A:370:ASN:HD22	2.12	0.48
1:A:271:THR:HG23	1:A:272:HIS:H	1.79	0.47
1:B:280:ASP:OD1	1:B:282:GLU:HB2	2.14	0.47
1:A:369:ARG:O	1:A:370:ASN:HB2	2.15	0.47
1:B:194:LYS:O	1:B:237:LYS:NZ	2.48	0.47
1:A:359:HIS:C	1:A:433:ASP:HB2	2.40	0.47
1:B:318:HIS:HB3	1:B:359:HIS:CD2	2.50	0.46
1:B:397:GLY:HA3	1:B:421:TYR:O	2.16	0.46
1:A:63:ILE:O	1:A:64:SER:C	2.59	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:LEU:C	1:A:352:LEU:HD23	2.40	0.46
1:A:297:MET:HE3	1:A:297:MET:HB3	1.71	0.46
1:A:458:ILE:HB	1:A:459:PRO:HD3	1.98	0.46
1:A:79:THR:OG1	1:A:99:THR:C	2.58	0.45
1:A:247:VAL:O	1:A:266:GLY:HA2	2.16	0.45
1:A:162:ARG:HG3	1:A:163:ASP:N	2.31	0.45
1:A:389:ILE:HD13	1:B:82:GLN:HE22	1.81	0.44
1:B:296:GLU:O	1:B:299:THR:HG22	2.17	0.44
1:A:112:ARG:O	1:A:113:THR:CB	2.65	0.44
1:B:90:ALA:HA	1:B:93:TYR:CZ	2.53	0.44
1:A:215:ILE:O	1:A:216:GLU:CB	2.66	0.44
1:B:297:MET:HB3	1:B:297:MET:HE3	1.71	0.44
1:A:93:TYR:CD1	1:A:94:LEU:HG	2.53	0.44
1:B:318:HIS:HB3	1:B:359:HIS:CG	2.53	0.44
1:B:114:ASN:ND2	2:B:483:NAG:C2	2.65	0.43
1:B:377:ALA:HB2	1:B:389:ILE:HA	2.01	0.43
1:B:416:ASP:HB3	1:B:419:PHE:HB2	2.01	0.43
1:B:119:LEU:HD11	1:B:179:ILE:HD11	2.00	0.43
1:B:292:LYS:HG3	1:B:343:MET:HE1	2.00	0.43
1:A:209:PRO:HA	1:A:223:ARG:HB2	2.01	0.43
1:B:292:LYS:HG3	1:B:343:MET:CE	2.48	0.43
1:A:287:LEU:N	1:A:288:PRO:CD	2.82	0.42
1:B:93:TYR:CD1	1:B:94:LEU:HG	2.54	0.42
1:B:261:THR:O	1:B:297:MET:HE1	2.20	0.42
1:A:318:HIS:HB3	1:A:359:HIS:CD2	2.55	0.42
1:B:209:PRO:HA	1:B:223:ARG:HB2	2.01	0.42
1:B:146:ARG:HA	1:B:202:GLY:C	2.44	0.42
1:A:90:ALA:HA	1:A:93:TYR:CZ	2.55	0.41
1:A:146:ARG:HA	1:A:202:GLY:C	2.45	0.41
1:A:149:HIS:CD2	1:A:152:PRO:HD2	2.55	0.41
1:A:271:THR:HG23	1:A:272:HIS:N	2.35	0.41
1:A:292:LYS:HA	1:A:343:MET:HE1	2.01	0.41
1:A:437:VAL:HB	6:A:507:HOH:O	2.19	0.41
1:B:287:LEU:N	1:B:288:PRO:CD	2.83	0.41
1:B:475:CYS:O	1:B:476:ASP:C	2.63	0.41
1:A:261:THR:O	1:A:297:MET:HE1	2.20	0.41
1:B:339:MET:SD	1:B:343:MET:HE3	2.58	0.41
1:B:473:THR:HG22	1:B:476:ASP:H	1.86	0.41
1:B:120:ASP:OD1	1:B:120:ASP:C	2.64	0.41
1:A:151:THR:HB	1:A:152:PRO:CD	2.51	0.41
1:B:96:GLY:C	1:B:456:ASN:HB2	2.46	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:208:PHE:CE2	1:B:221:GLY:CA	3.04	0.41
1:B:217:ASP:CB	1:B:219:ILE:HG13	2.50	0.41
1:B:41:LEU:HD23	1:B:41:LEU:HA	1.97	0.40
1:B:162:ARG:HG3	1:B:163:ASP:N	2.35	0.40
1:A:41:LEU:HD23	1:A:41:LEU:HA	1.98	0.40
1:B:287:LEU:HD23	1:B:333:MET:HA	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/478 (99%)	457 (96%)	16 (3%)	1 (0%)	43	44
1	B	474/478 (99%)	460 (97%)	13 (3%)	1 (0%)	43	44
All	All	948/956 (99%)	917 (97%)	29 (3%)	2 (0%)	43	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	216	GLU
1	B	216	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	394/398 (99%)	384 (98%)	10 (2%)	42	45
1	B	394/398 (99%)	386 (98%)	8 (2%)	48	54
All	All	788/796 (99%)	770 (98%)	18 (2%)	44	49

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LYS
1	A	178	GLU
1	A	179	ILE
1	A	265	MET
1	A	282	GLU
1	A	359	HIS
1	A	370	ASN
1	A	402	GLU
1	A	411	GLU
1	A	473	THR
1	B	22	LYS
1	B	67	GLU
1	B	179	ILE
1	B	245	SER
1	B	359	HIS
1	B	402	GLU
1	B	411	GLU
1	B	473	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	133	GLN
1	A	158	HIS
1	A	195	ASN
1	A	272	HIS
1	A	316	HIS
1	A	322	GLN
1	A	359	HIS
1	A	447	HIS
1	B	82	GLN
1	B	114	ASN
1	B	133	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	272	HIS
1	B	303	ASN
1	B	316	HIS
1	B	318	HIS
1	B	322	GLN
1	B	359	HIS
1	B	447	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	A	487	-	4,4,4	0.41	0	6,6,6	0.55	0
5	SO4	B	488	-	4,4,4	0.30	0	6,6,6	0.36	0
2	NAG	A	483	1	14,14,15	1.12	1 (7%)	17,19,21	3.02	8 (47%)
2	NAG	B	483	1	14,14,15	0.92	0	17,19,21	2.42	7 (41%)
4	PO4	A	486	3	4,4,4	1.48	1 (25%)	6,6,6	3.14	3 (50%)
4	PO4	B	484	3	4,4,4	1.36	1 (25%)	6,6,6	1.68	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	483	1	-	0/6/23/26	0/1/1/1
2	NAG	A	483	1	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	486	PO4	P-O4	-2.44	1.47	1.54
2	A	483	NAG	O5-C1	-2.27	1.39	1.43
4	B	484	PO4	P-O1	2.16	1.55	1.50

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	483	NAG	O6-C6-C5	5.54	130.21	111.33
2	A	483	NAG	C1-O5-C5	5.53	119.60	112.19
4	A	486	PO4	O2-P-O1	5.15	129.17	110.95
2	B	483	NAG	C1-O5-C5	4.90	118.76	112.19
2	A	483	NAG	O3-C3-C4	4.80	121.70	110.38
2	A	483	NAG	C1-C2-N2	-4.76	102.92	110.43
2	B	483	NAG	O6-C6-C5	4.53	126.76	111.33
4	A	486	PO4	O3-P-O1	-4.12	96.37	110.95
2	B	483	NAG	O5-C5-C6	4.05	115.55	107.66
4	A	486	PO4	O3-P-O2	-3.67	96.49	107.91
2	B	483	NAG	C3-C4-C5	3.46	116.50	110.23
2	A	483	NAG	O5-C1-C2	-3.33	106.14	111.29
2	A	483	NAG	O5-C5-C6	3.05	113.60	107.66
4	B	484	PO4	O3-P-O2	3.00	117.24	107.91
2	B	483	NAG	C4-C3-C2	-2.87	106.82	111.02
2	A	483	NAG	O7-C7-C8	-2.83	117.01	122.05
2	A	483	NAG	C6-C5-C4	2.78	119.85	113.02
2	B	483	NAG	O5-C1-C2	-2.53	107.38	111.29
2	B	483	NAG	C6-C5-C4	-2.22	107.56	113.02
4	B	484	PO4	O3-P-O1	-2.17	103.26	110.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	483	NAG	4	0
4	A	486	PO4	2	0
4	B	484	PO4	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/478 (99%)	0.04	7 (1%) 72 76	19, 27, 41, 59	0
1	B	476/478 (99%)	0.07	14 (2%) 53 57	19, 27, 41, 59	0
All	All	952/956 (99%)	0.05	21 (2%) 62 66	19, 27, 41, 59	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	215	ILE	3.3
1	B	471	GLY	3.1
1	B	476	ASP	3.0
1	A	297	MET	2.9
1	B	302	GLU	2.6
1	B	179	ILE	2.5
1	B	297	MET	2.4
1	B	215	ILE	2.4
1	B	402	GLU	2.4
1	B	473	THR	2.3
1	B	239	SER	2.3
1	B	299	THR	2.3
1	A	179	ILE	2.3
1	B	474	PHE	2.2
1	A	180	CYS	2.2
1	B	180	CYS	2.2
1	A	399	HIS	2.1
1	A	57	LYS	2.1
1	A	476	ASP	2.1
1	B	117	TYR	2.0
1	B	67	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	B	488	5/5	0.64	0.35	27,38,40,40	5
5	SO4	A	487	5/5	0.80	0.21	31,32,34,35	5
4	PO4	B	484	5/5	0.80	0.21	20,21,26,31	5
2	NAG	A	483	14/15	0.81	0.14	32,37,40,43	0
2	NAG	B	483	14/15	0.83	0.16	35,44,50,56	0
4	PO4	A	486	5/5	0.91	0.14	17,20,30,33	5
3	ZN	A	479	1/1	0.92	0.14	43,43,43,43	0
3	ZN	A	484	1/1	0.98	0.03	24,24,24,24	0
3	ZN	A	485	1/1	0.99	0.03	27,27,27,27	0
3	ZN	B	481	1/1	1.00	0.04	25,25,25,25	0
3	ZN	B	482	1/1	1.00	0.16	38,38,38,38	0
3	ZN	B	480	1/1	1.00	0.02	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.