



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 27, 2026 – 08:10 PM UTC

PDB ID : 3SH9 / pdb_00003sh9
Title : Crystal structure of fluorophore-labeled beta-lactamase PenP in complex with cefotaxime
Authors : Wong, W.-T.; Zhao, Y.-X.; Leung, Y.-C.
Deposited on : 2011-06-16
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	
1	B	266	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BB0	B	2	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4318 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	0	0
			1983	1243	344	392	4			
1	B	256	Total	C	N	O	S	0	0	0
			2000	1254	346	396	4			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	initiating methionine	UNP P00808
A	166	CYS	GLU	engineered mutation	UNP P00808
B	25	MET	-	initiating methionine	UNP P00808
B	166	CYS	GLU	engineered mutation	UNP P00808

- Molecule 2 is CEFOTAXIME, C3' cleaved, open, bound form (CCD ID: CEF) (formula: $C_{14}H_{15}N_5O_5S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			26	14	5	5	2		
2	B	1	Total	C	N	O	S	0	0
			26	14	5	5	2		

- Molecule 3 is 1-[6-(dimethylamino)naphthalen-2-yl]ethanone (CCD ID: BB0) (formula: C₁₄H₁₅NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			16	14	1	1		
3	B	1	Total	C	N	O	0	0
			16	14	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	116	Total	O	0	0
			116	116		
4	B	135	Total	O	0	0
			135	135		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-lactamase



• Molecule 1: Beta-lactamase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.39Å 91.63Å 66.01Å 90.00° 104.50° 90.00°	Depositor
Resolution (Å)	40.00 – 1.90 40.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.00-1.90) 99.9 (40.00-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.197 , 0.246 0.206 , 0.249	Depositor DCC
R_{free} test set	1983 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 32.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4318	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BB0, CEF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	6/2012 (0.3%)	1.11	7/2723 (0.3%)
1	B	0.95	4/2030 (0.2%)	1.35	14/2749 (0.5%)
All	All	0.96	10/4042 (0.2%)	1.24	21/5472 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	83	GLN	C-N	17.67	1.57	1.33
1	A	83	GLN	C-N	16.27	1.55	1.33
1	A	57	VAL	C-N	15.24	1.57	1.33
1	B	238	ALA	C-N	14.24	1.53	1.33
1	A	252	PRO	C-N	10.11	1.57	1.33
1	A	235	THR	C-N	-7.29	1.23	1.33
1	B	59	ALA	C-N	-6.68	1.25	1.33
1	B	57	VAL	C-O	-6.04	1.17	1.24
1	A	242	GLY	C-O	-5.37	1.17	1.24
1	A	59	ALA	C-N	5.33	1.40	1.33

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	252	PRO	O-C-N	-43.77	100.58	121.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	238	ALA	O-C-N	-21.62	102.06	123.46
1	A	83	GLN	O-C-N	-17.19	103.53	122.09
1	B	252	PRO	CA-C-N	14.59	135.96	119.32
1	B	252	PRO	C-N-CA	14.59	135.96	119.32
1	A	238	ALA	CA-C-N	11.32	141.10	120.95
1	A	238	ALA	C-N-CA	11.32	141.10	120.95
1	B	170	ASN	N-CA-C	8.85	125.10	111.56
1	B	59	ALA	O-C-N	-6.91	115.26	123.27
1	B	260	VAL	CB-CA-C	6.83	120.74	110.63
1	B	59	ALA	CA-C-N	5.87	130.32	121.40
1	B	59	ALA	C-N-CA	5.87	130.32	121.40
1	B	170	ASN	CA-C-N	5.86	132.25	121.70
1	B	170	ASN	C-N-CA	5.86	132.25	121.70
1	A	251	TRP	CA-C-N	-5.76	114.45	120.38
1	A	251	TRP	C-N-CA	-5.76	114.45	120.38
1	B	83	GLN	O-C-N	-5.56	115.82	122.27
1	B	238	ALA	O-C-N	5.54	128.80	123.04
1	B	56	THR	OG1-CB-CG2	5.38	120.07	109.30
1	A	86	LYS	O-C-N	-5.26	116.53	123.16
1	B	187	VAL	N-CA-CB	5.25	117.68	110.54

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	238	ALA	Mainchain
1	A	57	VAL	Mainchain
1	A	83	GLN	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1983	0	1996	15	0
1	B	2000	0	2014	38	0
2	A	26	0	0	1	0
2	B	26	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	16	0	15	4	1
3	B	16	0	15	8	1
4	A	116	0	0	3	0
4	B	135	0	0	10	0
All	All	4318	0	4040	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:CYS:SG	3:B:2:BB0:HAS	1.68	1.34
1:A:166:CYS:SG	3:A:2:BB0:CAS	2.28	1.21
1:A:166:CYS:SG	3:A:2:BB0:HASA	1.83	1.16
1:B:166:CYS:SG	3:B:2:BB0:CAS	2.34	1.15
1:A:166:CYS:SG	3:A:2:BB0:HAS	1.98	1.03
1:B:166:CYS:SG	3:B:2:BB0:CAF	2.63	0.86
1:B:166:CYS:SG	3:B:2:BB0:OAU	2.36	0.83
1:A:150:LYS:HD2	4:A:305:HOH:O	1.79	0.82
1:B:170:ASN:HB3	1:B:171:GLU:HB2	1.62	0.79
1:B:254:PRO:HD2	4:B:375:HOH:O	1.89	0.72
1:B:159:VAL:HG11	1:B:185:ALA:HB2	1.72	0.71
1:B:166:CYS:CB	3:B:2:BB0:HAS	2.20	0.71
1:A:178:GLN:NE2	4:A:341:HOH:O	2.24	0.70
1:A:166:CYS:HG	3:A:2:BB0:HAS	1.60	0.64
1:B:220:LEU:CD1	4:B:376:HOH:O	2.46	0.63
1:A:240:SER:O	1:A:241:TYR:HB2	2.01	0.61
1:B:220:LEU:HD13	4:B:376:HOH:O	1.99	0.61
1:B:159:VAL:CG1	1:B:185:ALA:HB2	2.33	0.58
1:B:187:VAL:HB	1:B:260:VAL:HG22	1.87	0.57
1:B:38:GLU:HG3	4:B:409:HOH:O	2.06	0.55
1:A:187:VAL:HG13	1:A:260:VAL:HG13	1.89	0.55
1:B:178:GLN:H	1:B:178:GLN:NE2	2.06	0.54
1:B:208:ILE:CD1	4:B:402:HOH:O	2.56	0.53
1:B:112:HIS:HD2	4:B:296:HOH:O	1.91	0.53
1:B:31:ASP:N	4:B:306:HOH:O	2.41	0.52
1:B:164:ARG:NH1	1:B:166:CYS:O	2.43	0.52
1:B:170:ASN:CB	1:B:171:GLU:HB2	2.39	0.51
2:A:3:CEF:O2	2:A:3:CEF:C3	2.58	0.51
1:B:208:ILE:HD11	4:B:402:HOH:O	2.09	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:31:ASP:HB3	4:B:24:HOH:O	2.10	0.51
1:B:220:LEU:HD12	1:B:220:LEU:N	2.26	0.50
1:B:220:LEU:HD22	1:B:235:THR:HG22	1.92	0.49
1:B:225:VAL:HG22	1:B:226:PRO:HD2	1.96	0.47
1:A:241:TYR:HA	1:A:269:LYS:O	2.14	0.47
1:B:104:ASN:HB3	3:B:2:BB0:CAI	2.44	0.46
1:B:166:CYS:HG	3:B:2:BB0:CAF	2.13	0.46
1:B:104:ASN:HB3	3:B:2:BB0:CAH	2.45	0.46
1:B:219:ALA:C	1:B:220:LEU:HD12	2.41	0.46
1:B:183:ALA:O	1:B:187:VAL:HG13	2.16	0.46
1:A:112:HIS:HD2	4:A:14:HOH:O	1.98	0.45
1:B:172:VAL:O	1:B:173:ASN:C	2.59	0.45
1:B:170:ASN:HB2	4:B:328:HOH:O	2.17	0.43
1:A:171:GLU:CD	1:A:270:LYS:HD3	2.43	0.43
1:A:48:ALA:HB2	1:A:261:LEU:HD13	2.01	0.43
1:B:190:LEU:HD21	1:B:211:MET:HE1	2.00	0.42
1:A:194:ALA:O	1:A:208:ILE:HD11	2.19	0.42
1:A:43:LYS:O	1:A:265:SER:HA	2.19	0.42
1:B:170:ASN:CB	1:B:171:GLU:CB	2.97	0.42
1:B:159:VAL:CG1	1:B:185:ALA:CB	2.97	0.42
1:B:220:LEU:CD1	1:B:220:LEU:N	2.83	0.42
1:A:164:ARG:HG3	1:A:179:ASP:OD2	2.20	0.41
1:B:212:LYS:HE2	1:B:230:GLU:HG2	2.02	0.41
1:B:159:VAL:HG13	1:B:185:ALA:CB	2.51	0.41
1:B:170:ASN:HB3	1:B:171:GLU:CB	2.41	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:2:BB0:OAU	3:B:2:BB0:CAI[2_656]	1.97	0.23

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/266 (94%)	240 (96%)	8 (3%)	2 (1%)	16	8
1	B	254/266 (96%)	245 (96%)	5 (2%)	4 (2%)	7	2
All	All	504/532 (95%)	485 (96%)	13 (3%)	6 (1%)	10	4

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	171	GLU
1	A	172	VAL
1	B	240	SER
1	A	103	VAL
1	B	173	ASN
1	B	103	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/225 (95%)	207 (97%)	7 (3%)	33	26
1	B	216/225 (96%)	200 (93%)	16 (7%)	13	6
All	All	430/450 (96%)	407 (95%)	23 (5%)	20	12

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ASP
1	A	159	VAL
1	A	161	ASN
1	A	164	ARG
1	A	178	GLN
1	A	225	VAL
1	A	233	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	44	LEU
1	B	56	THR
1	B	89	GLU
1	B	102	LEU
1	B	119	LEU
1	B	137	LEU
1	B	140	LYS
1	B	159	VAL
1	B	161	ASN
1	B	178	GLN
1	B	187	VAL
1	B	190	LEU
1	B	195	LEU
1	B	225	VAL
1	B	233	ASP
1	B	260	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	93	GLN
1	A	112	HIS
1	A	170	ASN
1	A	245	ASN
1	B	54	ASN
1	B	82	GLN
1	B	83	GLN
1	B	112	HIS
1	B	136	ASN
1	B	178	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BB0	B	2	-	17,17,17	0.87	0	24,24,24	0.62	0
2	CEF	B	301	1	24,27,27	3.49	9 (37%)	25,37,37	4.67	12 (48%)
2	CEF	A	3	1	24,27,27	3.32	8 (33%)	25,37,37	4.77	13 (52%)
3	BB0	A	2	-	17,17,17	0.90	0	24,24,24	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BB0	B	2	-	-	0/8/8/8	0/2/2/2
2	CEF	B	301	1	-	6/18/38/38	0/1/2/2
2	CEF	A	3	1	-	3/18/38/38	0/1/2/2
3	BB0	A	2	-	-	0/8/8/8	0/2/2/2

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	CEF	C1-C2	-10.30	1.37	1.50
2	A	3	CEF	C1-C2	-10.11	1.37	1.50
2	B	301	CEF	C1-S1	-7.78	1.66	1.82
2	A	3	CEF	C1-S1	-7.54	1.67	1.82
2	B	301	CEF	C4-N1	6.62	1.34	1.28
2	A	3	CEF	C4-C5	-6.07	1.36	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	CEF	C4-C5	-5.93	1.36	1.48
2	A	3	CEF	C4-N1	5.27	1.33	1.28
2	B	301	CEF	C4-C2	-3.48	1.35	1.46
2	A	3	CEF	C4-C2	-3.43	1.35	1.46
2	B	301	CEF	C14-N5	2.66	1.35	1.31
2	B	301	CEF	C13-C12	2.56	1.40	1.36
2	A	3	CEF	O2-C5	-2.30	1.24	1.30
2	A	3	CEF	C14-N5	2.16	1.34	1.31
2	A	3	CEF	C13-C12	2.07	1.40	1.36
2	B	301	CEF	O2-C5	-2.07	1.24	1.30
2	B	301	CEF	C10-N3	2.01	1.34	1.29

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3	CEF	C6-N1-C4	15.20	129.60	116.33
2	B	301	CEF	C6-N1-C4	13.76	128.34	116.33
2	B	301	CEF	C11-O5-N3	12.84	123.99	108.32
2	A	3	CEF	C13-S2-C14	10.45	93.35	88.84
2	A	3	CEF	C11-O5-N3	10.31	120.91	108.32
2	B	301	CEF	C13-S2-C14	9.21	92.81	88.84
2	B	301	CEF	C1-S1-C6	5.12	103.74	94.36
2	A	3	CEF	C1-S1-C6	4.51	102.62	94.36
2	A	3	CEF	C8-C7-N2	-4.40	102.86	109.80
2	A	3	CEF	C2-C1-S1	4.24	120.38	111.65
2	B	301	CEF	C2-C1-S1	3.71	119.30	111.65
2	A	3	CEF	C12-C13-S2	-3.37	106.35	110.24
2	B	301	CEF	C8-C7-N2	-3.28	104.62	109.80
2	A	3	CEF	S2-C14-N5	-3.08	112.11	114.54
2	A	3	CEF	O5-N3-C10	3.05	123.04	111.65
2	B	301	CEF	O5-N3-C10	3.04	122.99	111.65
2	B	301	CEF	C12-C13-S2	-3.02	106.75	110.24
2	B	301	CEF	S2-C14-N5	-3.01	112.17	114.54
2	A	3	CEF	C13-C12-C10	-2.80	120.68	125.69
2	B	301	CEF	O2-C5-O1	-2.71	117.46	123.90
2	B	301	CEF	C13-C12-C10	-2.50	121.21	125.69
2	B	301	CEF	S2-C14-N4	2.50	123.90	120.99
2	A	3	CEF	O2-C5-O1	-2.43	118.13	123.90
2	A	3	CEF	S2-C14-N4	2.14	123.48	120.99
2	A	3	CEF	O2-C5-C4	2.04	121.41	116.02

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	3	CEF	C2-C4-C5-O1
2	B	301	CEF	C2-C4-C5-O1
2	B	301	CEF	C12-C10-C9-O4
2	A	3	CEF	N3-C10-C9-N2
2	B	301	CEF	N3-C10-C9-O4
2	B	301	CEF	C12-C10-C9-N2
2	B	301	CEF	N3-C10-C9-N2
2	A	3	CEF	N3-C10-C9-O4
2	B	301	CEF	N1-C4-C5-O2


There are no ring outliers.

3 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	2	BB0	8	1
2	A	3	CEF	1	0
3	A	2	BB0	4	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand CEF B 301




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	254/266 (95%)	0.43	21 (8%) 17 18	10, 21, 42, 71	0
1	B	256/266 (96%)	0.13	14 (5%) 30 32	11, 17, 34, 54	0
All	All	510/532 (95%)	0.28	35 (6%) 23 24	10, 19, 40, 71	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	172	VAL	6.5
1	B	174	PRO	5.7
1	B	172	VAL	5.1
1	A	174	PRO	4.8
1	A	291	ASN	4.5
1	A	167	PRO	4.1
1	A	175	GLY	3.9
1	A	176	GLU	3.8
1	B	173	ASN	3.7
1	A	165	PHE	3.7
1	A	170	ASN	3.6
1	A	271	ASP	3.5
1	A	171	GLU	3.3
1	A	38	GLU	3.1
1	A	173	ASN	3.0
1	A	166	CYS	3.0
1	B	167	PRO	3.0
1	A	227	ASP	3.0
1	B	169	LEU	2.9
1	B	291	ASN	2.8
1	A	240	SER	2.8
1	B	176	GLU	2.6
1	B	290	LEU	2.6
1	B	175	GLY	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	171	GLU	2.5
1	A	268	ASP	2.5
1	A	255	LYS	2.5
1	A	177	THR	2.5
1	B	228	GLY	2.4
1	A	272	ALA	2.2
1	A	52	GLY	2.2
1	B	170	ASN	2.2
1	A	270	LYS	2.1
1	B	226	PRO	2.0
1	B	225	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BB0	B	2	16/16	0.60	0.33	85,85,85,85	0
3	BB0	A	2	16/16	0.68	0.38	126,126,126,126	0
2	CEF	A	3	26/26	0.92	0.10	14,25,34,34	0
2	CEF	B	301	26/26	0.93	0.09	15,22,24,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CEF A 3:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around CEF B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.