



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 07:45 PM UTC

PDB ID : 6SH7 / pdb_00006sh7
Title : Crystal structure of the human DEAH-helicase DHX15 in complex with the NKRF G-patch
Authors : Jonas, S.; Studer, M.K.; Ivanovic, L.
Deposited on : 2019-08-05
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	689	2% 87% 9% .
2	B	67	4% 67% 31% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5745 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor ATP-dependent RNA helicase DHX15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	665	Total	C	N	O	S	0	0	0
			5362	3406	927	992	37			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	GLY	-	expression tag	UNP O43143
A	108	PRO	-	expression tag	UNP O43143
A	109	HIS	-	expression tag	UNP O43143
A	110	MET	-	expression tag	UNP O43143
A	111	LEU	-	expression tag	UNP O43143
A	112	GLU	-	expression tag	UNP O43143

- Molecule 2 is a protein called NF-kappa-B-repressing factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	46	Total	C	N	O	S	0	0	0
			349	213	66	69	1			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	537	GLY	-	expression tag	UNP O15226
B	538	PRO	-	expression tag	UNP O15226
B	539	HIS	-	expression tag	UNP O15226
B	540	MET	-	expression tag	UNP O15226


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	34	Total 34	O 34	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor ATP-dependent RNA helicase DHX15



- Molecule 2: NF-kappa-B-repressing factor



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	82.24Å 90.02Å 213.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.23 – 2.21 46.23 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.23-2.21) 99.8 (46.23-2.21)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.00 (at 2.22Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.209 , 0.251 0.214 , 0.251	Depositor DCC
R_{free} test set	1983 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	61.8	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5745	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.10	0/5472	0.30	0/7413
2	B	0.11	0/352	0.29	0/467
All	All	0.10	0/5824	0.29	0/7880

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5362	0	5386	33	0
2	B	349	0	348	1	0
3	A	34	0	0	0	0
All	All	5745	0	5734	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:392:LEU:HD12	1:A:393:PRO:HD2	1.75	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:LYS:HG2	1:A:405:PRO:HD3	1.80	0.62
1:A:742:THR:OG1	1:A:743:THR:N	2.30	0.60
1:A:256:VAL:HG22	1:A:287:LYS:HB2	1.89	0.54
1:A:768:MET:O	1:A:777:LYS:NZ	2.31	0.54
1:A:193:PRO:HB3	1:A:266:THR:HG21	1.91	0.51
1:A:446:VAL:HG21	1:A:457:LEU:HD12	1.92	0.51
1:A:633:ALA:HA	1:A:636:GLN:HG2	1.93	0.50
1:A:389:TYR:CZ	1:A:392:LEU:HD13	2.46	0.50
1:A:564:GLU:HG3	1:A:671:PHE:HZ	1.77	0.50
1:A:384:LYS:HE2	1:A:405:PRO:HB3	1.94	0.49
1:A:343:HIS:HD2	1:A:415:ARG:HD2	1.76	0.49
1:A:356:LEU:HD13	1:A:361:GLU:HG2	1.94	0.49
1:A:130:ILE:HG21	1:A:208:GLU:HG2	1.94	0.48
1:A:771:PHE:HB3	1:A:777:LYS:HD2	1.94	0.48
2:B:554:ILE:H	2:B:554:ILE:HD12	1.79	0.47
1:A:385:ILE:HA	1:A:417:VAL:HG13	1.97	0.47
1:A:705:GLU:OE1	1:A:707:THR:OG1	2.31	0.46
1:A:349:GLU:O	1:A:477:ARG:NH1	2.30	0.46
1:A:192:GLN:O	1:A:237:THR:HA	2.16	0.46
1:A:514:LEU:HD22	1:A:519:ILE:HD12	1.99	0.45
1:A:178:MET:HE2	1:A:186:ARG:HB2	1.99	0.44
1:A:259:LEU:HD11	1:A:276:LEU:HD12	2.00	0.43
1:A:605:GLU:H	1:A:605:GLU:CD	2.26	0.43
1:A:389:TYR:O	1:A:397:GLN:NE2	2.51	0.42
1:A:399:ARG:O	1:A:402:GLU:HG2	2.18	0.42
1:A:704:LEU:HB3	1:A:732:GLU:HA	2.00	0.42
1:A:167:THR:HG22	1:A:201:VAL:HG21	2.01	0.42
1:A:564:GLU:HG3	1:A:671:PHE:CZ	2.55	0.42
1:A:585:ASN:OD1	1:A:675:ARG:HB3	2.18	0.42
1:A:719:VAL:HG12	1:A:745:ASN:HB3	2.02	0.41
1:A:466:ALA:HB2	1:A:494:MET:HE1	2.03	0.41
1:A:544:ASN:OD1	1:A:764:GLN:NE2	2.46	0.40
1:A:600:PHE:CE1	1:A:614:LYS:HG3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	657/689 (95%)	635 (97%)	22 (3%)	0	100	100
2	B	44/67 (66%)	42 (96%)	2 (4%)	0	100	100
All	All	701/756 (93%)	677 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	596/615 (97%)	588 (99%)	8 (1%)	61	75
2	B	36/53 (68%)	36 (100%)	0	100	100
All	All	632/668 (95%)	624 (99%)	8 (1%)	61	75

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	VAL
1	A	210	ASP
1	A	232	ILE
1	A	434	VAL
1	A	498	THR
1	A	510	VAL
1	A	517	LEU
1	A	742	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	113	GLN
1	A	123	HIS
1	A	136	GLN
1	A	341	GLN
1	A	343	HIS
1	A	395	GLN
1	A	495	GLN
1	A	513	GLN
1	A	709	HIS
1	A	729	HIS
1	A	788	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	665/689 (96%)	0.32	12 (1%) 67 65	46, 71, 132, 176	0
2	B	46/67 (68%)	0.77	3 (6%) 25 22	78, 106, 147, 167	0
All	All	711/756 (94%)	0.35	15 (2%) 63 61	46, 73, 134, 176	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	428	LEU	3.4
1	A	774	CYS	3.3
1	A	470	ALA	3.2
1	A	184	PRO	2.8
1	A	295	LEU	2.8
2	B	582	VAL	2.8
1	A	741	LEU	2.6
1	A	433	VAL	2.4
2	B	554	ILE	2.4
2	B	595	VAL	2.3
1	A	429	THR	2.3
1	A	498	THR	2.3
1	A	297	ALA	2.1
1	A	163	GLY	2.0
1	A	121	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.