



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:30 AM UTC

PDB ID : 3SJU / pdb_00003sju
Title : Hedamycin Polyketide Ketoreductase bound to NADPH
Authors : Javidpour, P.; Tsai, S.-C.
Deposited on : 2011-06-21
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	 2% 63% 24% 9%
1	B	279	 % 68% 18% 5% 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Keto reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	255	Total	C	N	O	S	0	0	0
			1848	1147	335	359	7			
1	B	255	Total	C	N	O	S	0	0	0
			1848	1147	335	359	7			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q67G28
A	-18	GLY	-	expression tag	UNP Q67G28
A	-17	SER	-	expression tag	UNP Q67G28
A	-16	SER	-	expression tag	UNP Q67G28
A	-15	HIS	-	expression tag	UNP Q67G28
A	-14	HIS	-	expression tag	UNP Q67G28
A	-13	HIS	-	expression tag	UNP Q67G28
A	-12	HIS	-	expression tag	UNP Q67G28
A	-11	HIS	-	expression tag	UNP Q67G28
A	-10	HIS	-	expression tag	UNP Q67G28
A	-9	SER	-	expression tag	UNP Q67G28
A	-8	SER	-	expression tag	UNP Q67G28
A	-7	GLY	-	expression tag	UNP Q67G28
A	-6	LEU	-	expression tag	UNP Q67G28
A	-5	VAL	-	expression tag	UNP Q67G28
A	-4	PRO	-	expression tag	UNP Q67G28
A	-3	ARG	-	expression tag	UNP Q67G28
A	-2	GLY	-	expression tag	UNP Q67G28
A	-1	SER	-	expression tag	UNP Q67G28
A	0	HIS	-	expression tag	UNP Q67G28
B	-19	MET	-	expression tag	UNP Q67G28
B	-18	GLY	-	expression tag	UNP Q67G28
B	-17	SER	-	expression tag	UNP Q67G28
B	-16	SER	-	expression tag	UNP Q67G28
B	-15	HIS	-	expression tag	UNP Q67G28

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP Q67G28
B	-13	HIS	-	expression tag	UNP Q67G28
B	-12	HIS	-	expression tag	UNP Q67G28
B	-11	HIS	-	expression tag	UNP Q67G28
B	-10	HIS	-	expression tag	UNP Q67G28
B	-9	SER	-	expression tag	UNP Q67G28
B	-8	SER	-	expression tag	UNP Q67G28
B	-7	GLY	-	expression tag	UNP Q67G28
B	-6	LEU	-	expression tag	UNP Q67G28
B	-5	VAL	-	expression tag	UNP Q67G28
B	-4	PRO	-	expression tag	UNP Q67G28
B	-3	ARG	-	expression tag	UNP Q67G28
B	-2	GLY	-	expression tag	UNP Q67G28
B	-1	SER	-	expression tag	UNP Q67G28
B	0	HIS	-	expression tag	UNP Q67G28

- # NDP


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	20	Total 20	O 20	0	0
3	B	18	Total 18	O 18	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Keto reductase



• Molecule 1: Keto reductase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	116.70Å 57.39Å 82.11Å 90.00° 131.61° 90.00°	Depositor
Resolution (Å)	50.00 – 2.40 50.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.0 (50.00-2.40) 99.0 (50.00-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.92 (at 2.39Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.182 , 0.248 0.184 , 0.245	Depositor DCC
R_{free} test set	791 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	33.9	Xtriage
Anisotropy	1.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3830	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	3/1878 (0.2%)	1.15	6/2553 (0.2%)
1	B	1.00	2/1878 (0.1%)	1.12	2/2553 (0.1%)
All	All	1.02	5/3756 (0.1%)	1.14	8/5106 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	130	ARG	C-N	-8.24	1.21	1.33
1	B	95	GLY	C-N	-6.46	1.25	1.33
1	B	149	VAL	CA-CB	5.74	1.61	1.54
1	A	243	ALA	CA-CB	5.20	1.62	1.53
1	A	202	ARG	C-N	5.10	1.40	1.33

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	145	GLY	N-CA-C	-5.93	107.56	115.32
1	A	190	THR	CA-C-N	5.67	124.87	118.97
1	A	190	THR	C-N-CA	5.67	124.87	118.97
1	A	239	THR	N-CA-CB	-5.27	101.58	110.49
1	A	108	VAL	N-CA-C	5.17	115.78	110.36
1	B	199	GLY	N-CA-C	-5.09	107.47	113.99
1	A	46	ASP	N-CA-C	5.05	116.60	111.14
1	B	95	GLY	N-CA-C	-5.03	106.52	111.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1848	0	1814	51	0
1	B	1848	0	1814	47	0
2	A	48	0	26	3	0
2	B	48	0	26	2	0
3	A	20	0	0	0	0
3	B	18	0	0	0	0
All	All	3830	0	3680	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:ARG:HG3	1:A:202:ARG:NH1	1.55	1.06
1:A:202:ARG:HH11	1:A:202:ARG:CG	1.70	1.00
1:A:114:THR:HB	1:B:105:TRP:HE1	1.26	0.98
1:A:202:ARG:HG3	1:A:202:ARG:HH11	0.80	0.94
1:A:114:THR:HB	1:B:105:TRP:NE1	1.89	0.88
1:B:129:MET:HE3	1:B:178:ILE:HD13	1.56	0.87
1:B:92:ASN:HD21	1:B:192:MET:HE3	1.40	0.86
1:B:39:LYS:HE2	1:B:40:ASN:HD21	1.43	0.83
1:A:170:GLY:HA2	1:A:180:VAL:HG13	1.61	0.82
1:B:129:MET:CE	1:B:178:ILE:HD13	2.12	0.80
1:A:236:TYR:O	1:A:239:THR:HB	1.85	0.76
1:A:97:THR:HA	1:A:100:LEU:HD22	1.68	0.76
1:A:110:ASP:HA	1:A:114:THR:HG23	1.68	0.73
1:B:13:SER:HB2	2:B:301:NDP:O3B	1.89	0.73
1:A:63:THR:HA	1:A:118:ARG:NH2	2.04	0.72
1:A:130:ARG:NH2	1:B:96:GLU:OE2	2.23	0.72
1:B:39:LYS:HE2	1:B:40:ASN:ND2	2.04	0.71
1:A:36:ARG:O	1:A:59:SER:HB2	1.91	0.70
1:A:56:ASP:OD2	1:A:78:ARG:NE	2.20	0.70
1:A:200:TYR:HB3	1:A:211:VAL:HG11	1.73	0.70
1:B:129:MET:HE3	1:B:178:ILE:CD1	2.21	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:251:ASN:ND2	1:A:257:GLY:HA3	2.10	0.67
1:A:239:THR:HG22	1:A:242:ALA:H	1.61	0.66
1:B:92:ASN:ND2	1:B:192:MET:HE3	2.11	0.65
1:B:91:ARG:HD3	1:B:92:ASN:O	1.97	0.65
1:B:14:SER:OG	1:B:191:PRO:HG2	1.96	0.65
1:B:189:GLU:CD	1:B:197:ARG:HH22	2.07	0.63
1:B:197:ARG:O	1:B:200:TYR:HB2	2.01	0.60
1:A:36:ARG:NH2	2:A:301:NDP:O3X	2.29	0.60
1:B:36:ARG:O	1:B:59:SER:OG	2.21	0.58
1:B:189:GLU:OE1	1:B:197:ARG:NH2	2.37	0.58
1:A:63:THR:HA	1:A:118:ARG:HH21	1.70	0.57
1:B:63:THR:HB	1:B:110:ASP:OD2	2.04	0.57
1:A:190:THR:HB	1:A:191:PRO:HD2	1.86	0.57
1:A:251:ASN:HD22	1:A:257:GLY:HA3	1.68	0.57
1:B:190:THR:HG22	1:B:227:PRO:HG3	1.87	0.56
1:A:197:ARG:O	1:A:200:TYR:HB2	2.07	0.55
1:B:206:VAL:O	1:B:210:GLU:HB2	2.06	0.55
1:A:197:ARG:HA	1:A:200:TYR:HD2	1.73	0.54
1:B:196:VAL:O	1:B:200:TYR:HD2	1.89	0.54
1:A:196:VAL:O	1:A:200:TYR:CD2	2.61	0.54
1:B:82:ILE:O	1:B:129:MET:HG3	2.07	0.54
1:B:92:ASN:HD21	1:B:192:MET:CE	2.16	0.54
2:A:301:NDP:H52A	2:A:301:NDP:H52N	1.89	0.53
1:A:124:LEU:O	1:A:130:ARG:HD2	2.09	0.53
1:A:194:GLU:O	1:A:198:GLU:HG2	2.09	0.52
1:B:136:ARG:HD3	1:B:237:LEU:O	2.08	0.52
1:A:239:THR:HG23	1:A:241:ALA:H	1.75	0.52
1:A:87:ASN:HD21	1:A:119:VAL:HG22	1.74	0.52
1:A:189:GLU:HA	1:A:189:GLU:OE2	2.10	0.51
1:A:170:GLY:CA	1:A:180:VAL:HG13	2.37	0.50
1:B:16:ILE:O	1:B:20:VAL:HG23	2.12	0.49
1:B:106:ALA:O	1:B:110:ASP:HB2	2.13	0.49
1:B:136:ARG:HG2	1:B:238:VAL:HA	1.94	0.49
1:B:191:PRO:HA	1:B:194:GLU:OE2	2.13	0.49
1:A:5:GLN:O	1:A:29:ILE:HG23	2.12	0.48
1:A:105:TRP:HZ2	1:B:113:LEU:HD22	1.79	0.48
1:A:172:GLU:HA	1:B:150:MET:HE2	1.94	0.48
1:B:91:ARG:HD2	1:B:108:VAL:HG22	1.96	0.48
1:A:202:ARG:NH1	1:A:202:ARG:CG	2.41	0.48
1:A:35:ALA:O	1:A:59:SER:HA	2.14	0.47
1:A:25:ALA:HB1	1:A:53:HIS:HB3	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:129:MET:HE1	1:B:178:ILE:HG21	1.98	0.46
1:A:204:TRP:HB3	1:A:206:VAL:HG13	1.97	0.46
1:A:136:ARG:HG3	1:A:238:VAL:HA	1.99	0.45
1:B:141:ALA:O	2:B:301:NDP:H6N	2.16	0.45
1:B:196:VAL:O	1:B:200:TYR:CD2	2.70	0.44
1:B:37:ASP:OD2	1:B:39:LYS:HG3	2.18	0.44
1:B:188:VAL:HG22	1:B:190:THR:HG23	1.99	0.43
1:B:203:HIS:O	1:B:204:TRP:C	2.61	0.43
1:A:97:THR:OG1	1:B:172:GLU:OE1	2.24	0.43
1:A:168:SER:OG	1:B:156:THR:HG21	2.18	0.43
1:B:85:LEU:HG	1:B:123:VAL:HG21	2.01	0.43
1:A:103:ALA:O	1:A:107:ASP:HB2	2.19	0.42
1:A:210:GLU:O	1:A:214:ARG:HG3	2.19	0.42
1:B:49:ARG:HH11	1:B:49:ARG:HG2	1.84	0.42
1:A:171:PHE:HB3	1:B:150:MET:HG2	2.01	0.42
1:B:153:ALA:N	1:B:154:PRO:CD	2.83	0.42
1:A:153:ALA:N	1:A:154:PRO:CD	2.83	0.42
1:B:221:LEU:O	1:B:221:LEU:HG	2.19	0.41
1:B:39:LYS:HG3	1:B:40:ASN:N	2.36	0.41
1:A:14:SER:OG	1:A:191:PRO:HG2	2.20	0.41
1:A:16:ILE:O	1:A:20:VAL:HG23	2.21	0.41
1:A:113:LEU:HD13	1:B:109:LEU:HD11	2.03	0.41
1:A:192:MET:HE3	2:A:301:NDP:O1A	2.20	0.41
1:A:188:VAL:C	1:A:224:TYR:HD1	2.28	0.41
1:B:139:ASN:O	1:B:182:ALA:HA	2.21	0.41
1:A:32:TYR:CD2	1:A:75:ALA:HB2	2.56	0.41
1:A:45:VAL:O	1:A:49:ARG:HG3	2.21	0.41
1:A:91:ARG:HD3	1:A:92:ASN:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/279 (91%)	231 (91%)	21 (8%)	1 (0%)	30	43
1	B	253/279 (91%)	232 (92%)	20 (8%)	1 (0%)	30	43
All	All	506/558 (91%)	463 (92%)	41 (8%)	2 (0%)	30	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	15	GLY
1	B	81	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/203 (90%)	159 (87%)	23 (13%)	4	6
1	B	182/203 (90%)	160 (88%)	22 (12%)	5	7
All	All	364/406 (90%)	319 (88%)	45 (12%)	4	6

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	LEU
1	A	31	VAL
1	A	48	LEU
1	A	77	GLU
1	A	100	LEU
1	A	113	LEU
1	A	114	THR
1	A	119	VAL
1	A	131	GLU
1	A	140	ILE
1	A	180	VAL
1	A	194	GLU
1	A	198	GLU
1	A	202	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	203	HIS
1	A	204	TRP
1	A	208	GLU
1	A	211	VAL
1	A	218	LYS
1	A	233	LEU
1	A	238	VAL
1	A	239	THR
1	A	244	SER
1	B	5	GLN
1	B	39	LYS
1	B	48	LEU
1	B	59	SER
1	B	85	LEU
1	B	91	ARG
1	B	101	ASP
1	B	113	LEU
1	B	119	VAL
1	B	129	MET
1	B	130	ARG
1	B	136	ARG
1	B	149	VAL
1	B	188	VAL
1	B	194	GLU
1	B	202	ARG
1	B	204	TRP
1	B	206	VAL
1	B	208	GLU
1	B	238	VAL
1	B	250	LEU
1	B	256	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	87	ASN
1	A	139	ASN
1	A	160	HIS
1	A	216	ASN
1	A	258	ASN
1	B	5	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	40	ASN
1	B	87	ASN
1	B	92	ASN
1	B	139	ASN
1	B	160	HIS
1	B	181	ASN
1	B	209	GLN
1	B	212	HIS
1	B	216	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	B	301	-	51,52,52	1.44	8 (15%)	71,80,80	1.60	16 (22%)
2	NDP	A	301	-	51,52,52	1.32	6 (11%)	71,80,80	1.98	17 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	B	301	-	-	6/34/77/77	0/5/5/5
2	NDP	A	301	-	-	3/34/77/77	0/5/5/5

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	NDP	C4N-C3N	-5.13	1.40	1.50
2	A	301	NDP	C4N-C3N	-4.65	1.41	1.50
2	B	301	NDP	PN-O3	3.61	1.63	1.59
2	A	301	NDP	C4N-C5N	-3.51	1.39	1.49
2	A	301	NDP	C8A-N7A	3.14	1.37	1.31
2	B	301	NDP	PA-O3	3.13	1.62	1.59
2	B	301	NDP	C6N-C5N	2.74	1.41	1.33
2	B	301	NDP	C4N-C5N	-2.64	1.42	1.49
2	B	301	NDP	C8A-N7A	2.59	1.36	1.31
2	B	301	NDP	C2N-C3N	2.40	1.41	1.35
2	A	301	NDP	C2N-C3N	2.32	1.41	1.35
2	A	301	NDP	C6N-C5N	2.28	1.40	1.33
2	A	301	NDP	C5A-N7A	-2.20	1.35	1.39
2	B	301	NDP	C4A-N9A	-2.02	1.33	1.37

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	NDP	N3A-C2A-N1A	-5.78	119.83	128.58
2	A	301	NDP	C5A-C4A-N3A	-5.41	119.27	126.72
2	A	301	NDP	O4D-C1D-N1N	5.37	118.33	108.08
2	A	301	NDP	O2N-PN-O3	4.68	119.92	107.27
2	B	301	NDP	C5A-C4A-N3A	-4.66	120.30	126.72
2	B	301	NDP	N3A-C2A-N1A	-4.13	122.32	128.58
2	A	301	NDP	C2A-N3A-C4A	4.07	121.78	111.83
2	B	301	NDP	O4D-C1D-N1N	3.81	115.36	108.08
2	B	301	NDP	N9A-C8A-N7A	-3.57	108.87	113.94
2	B	301	NDP	C4A-C5A-N7A	-3.42	106.67	110.58
2	A	301	NDP	C4A-C5A-N7A	-3.26	106.86	110.58
2	B	301	NDP	C2A-N3A-C4A	3.03	119.24	111.83
2	A	301	NDP	C1D-N1N-C6N	-2.92	114.61	120.77
2	A	301	NDP	C5A-C4A-N9A	2.90	108.98	105.81
2	A	301	NDP	C3N-C2N-N1N	-2.83	119.04	123.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	NDP	O2A-PA-O1A	2.83	125.61	112.44
2	A	301	NDP	N9A-C8A-N7A	-2.82	109.93	113.94
2	B	301	NDP	C5A-N7A-C8A	2.76	107.79	103.45
2	A	301	NDP	N3A-C4A-N9A	2.70	131.76	127.17
2	B	301	NDP	O2B-P2B-O1X	-2.68	99.80	109.33
2	A	301	NDP	C5A-N7A-C8A	2.54	107.44	103.45
2	B	301	NDP	N3A-C4A-N9A	2.48	131.38	127.17
2	B	301	NDP	C4A-N9A-C8A	2.48	108.34	105.74
2	A	301	NDP	C3N-C7N-N7N	2.41	121.95	117.67
2	B	301	NDP	O3-PA-O1A	-2.41	103.47	110.70
2	B	301	NDP	C5A-C4A-N9A	2.30	108.32	105.81
2	A	301	NDP	O3-PA-O1A	-2.27	103.88	110.70
2	A	301	NDP	O3-PN-O1N	-2.12	104.33	110.70
2	A	301	NDP	C5B-C4B-C3B	-2.11	107.63	115.21
2	B	301	NDP	O2X-P2B-O1X	2.08	118.93	110.83
2	B	301	NDP	C1D-N1N-C6N	-2.06	116.41	120.77
2	B	301	NDP	O2N-PN-O3	2.02	112.75	107.27
2	B	301	NDP	O2A-PA-O3	2.00	112.69	107.27

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	NDP	C5D-O5D-PN-O3
2	B	301	NDP	C5D-O5D-PN-O1N
2	B	301	NDP	C5D-O5D-PN-O2N
2	A	301	NDP	O4D-C4D-C5D-O5D
2	A	301	NDP	O4D-C1D-N1N-C6N
2	A	301	NDP	C3D-C4D-C5D-O5D
2	B	301	NDP	O4D-C1D-N1N-C6N
2	B	301	NDP	PA-O3-PN-O1N
2	B	301	NDP	PA-O3-PN-O2N


There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	NDP	2	0
2	A	301	NDP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/279 (91%)	-0.20	5 (1%) 65 60	30, 42, 71, 106	0
1	B	255/279 (91%)	-0.11	3 (1%) 76 73	30, 45, 89, 109	0
All	All	510/558 (91%)	-0.15	8 (1%) 70 66	30, 44, 83, 109	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	204	TRP	4.8
1	A	201	ALA	3.6
1	B	204	TRP	3.4
1	B	203	HIS	2.8
1	A	203	HIS	2.7
1	A	200	TYR	2.4
1	B	205	GLY	2.3
1	A	205	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NDP	A	301	48/48	0.92	0.08	38,52,57,58	0
2	NDP	B	301	48/48	0.93	0.07	39,53,58,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.