



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:38 AM UTC

PDB ID : 5SJG / pdb_00005sjg
Title : CRYSTAL STRUCTURE OF HUMAN PHOSPHODIESTERASE 10 IN
COMPLEX WITH C4CCN(c1cn(c(n1)CCc3nn2c(ncc(c2n3)C)C)C)C4=
O, micromolar IC50=0.039238
Authors : Joseph, C.; Benz, J.; Flohr, A.; Rudolph, M.G.
Deposited on : 2022-02-01
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	
2	B	343	
2	C	343	
2	D	343	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10883 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	312	Total	C	N	O	S	0	2	0
			2555	1633	434	463	25			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	447	GLY	-	expression tag	UNP Q9Y233
A	448	SER	-	expression tag	UNP Q9Y233

- Molecule 2 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	315	Total	C	N	O	S	0	3	0
			2562	1637	434	467	24			
2	C	314	Total	C	N	O	S	0	4	0
			2565	1639	436	466	24			
2	D	311	Total	C	N	O	S	0	1	0
			2528	1617	430	457	24			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	447	GLY	-	expression tag	UNP Q9Y233
B	448	SER	-	expression tag	UNP Q9Y233
C	447	GLY	-	expression tag	UNP Q9Y233
C	448	SER	-	expression tag	UNP Q9Y233
D	447	GLY	-	expression tag	UNP Q9Y233
D	448	SER	-	expression tag	UNP Q9Y233


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		

- Molecule 5 is 1-(2-{2-[(4S)-5,8-dimethyl[1,2,4]triazolo[1,5-c]pyrimidin-2-yl]ethyl}-1-methyl-1H-imidazol-4-yl)pyrrolidin-2-one (CCD ID: K2L) (formula: C₁₇H₂₁N₇O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			25	17	7	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			25	17	7	1		
5	C	1	Total	C	N	O	0	0
			25	17	7	1		
5	D	1	Total	C	N	O	0	0
			25	17	7	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	145	Total	O	0	0
			145	145		
6	B	166	Total	O	0	0
			166	166		
6	C	155	Total	O	0	0
			155	155		
6	D	99	Total	O	0	0
			99	99		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A




- Molecule 2: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A




- Molecule 2: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 2: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A





4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	135.48Å 135.48Å 235.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.58 – 1.97 43.58 – 1.97	Depositor EDS
% Data completeness (in resolution range)	86.7 (43.58-1.97) 78.0 (43.58-1.97)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.176 , 0.224 (Not available) , 0.210	Depositor DCC
R_{free} test set	5121 reflections (4.50%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10883	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, K2L, ZN, CME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	5/2601 (0.2%)	1.39	3/3517 (0.1%)
2	B	1.14	5/2622 (0.2%)	1.42	8/3548 (0.2%)
2	C	1.15	0/2628	1.41	1/3555 (0.0%)
2	D	1.15	3/2582 (0.1%)	1.44	9/3494 (0.3%)
All	All	1.16	13/10433 (0.1%)	1.42	21/14114 (0.1%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	567	HIS	CE1-NE2	7.10	1.39	1.32
2	D	535	HIS	CE1-NE2	7.01	1.39	1.32
2	B	677[A]	SER	C-O	5.80	1.30	1.24
2	B	677[B]	SER	C-O	5.80	1.30	1.24
1	A	748	THR	C-O	5.78	1.31	1.24
1	A	506	GLU	C-O	5.75	1.30	1.24
1	A	730	TYR	C-O	5.34	1.30	1.24
1	A	481	PRO	C-O	-5.31	1.17	1.24
1	A	740	THR	C-O	5.31	1.30	1.24
2	B	586	TYR	C-O	5.26	1.30	1.23
2	B	476	HIS	CE1-NE2	5.25	1.37	1.32
2	D	567	HIS	CE1-NE2	5.19	1.37	1.32
2	D	481	PRO	C-O	-5.04	1.17	1.24

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	609	ASN	CA-CB-CG	-7.81	104.79	112.60
1	A	609	ASN	N-CA-CB	6.25	119.34	110.46
2	D	535	HIS	CA-CB-CG	-5.99	107.81	113.80
2	B	619	TYR	N-CA-C	-5.99	104.67	111.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	609	ASN	N-CA-CB	5.85	118.76	110.46
2	D	746	PRO	N-CA-C	5.49	117.40	110.70
2	B	599	THR	N-CA-C	-5.34	105.54	111.36
2	B	504	GLU	CB-CA-C	5.32	119.02	110.29
2	B	612	SER	CA-C-N	5.24	127.56	120.38
2	B	612	SER	C-N-CA	5.24	127.56	120.38
2	D	639	PHE	CA-C-N	5.14	125.69	119.98
2	D	639	PHE	C-N-CA	5.14	125.69	119.98
2	B	698	ALA	N-CA-C	-5.11	105.71	111.28
2	D	554	LYS	CA-C-N	5.10	125.61	120.00
2	D	554	LYS	C-N-CA	5.10	125.61	120.00
2	D	549	THR	CA-C-N	5.09	127.11	120.28
2	D	549	THR	C-N-CA	5.09	127.11	120.28
2	C	724	GLN	N-CA-C	-5.04	105.97	111.82
2	B	554	LYS	CA-C-N	5.03	125.56	119.98
2	B	554	LYS	C-N-CA	5.03	125.56	119.98
1	A	602	ILE	CA-C-O	-5.00	115.54	120.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2555	0	2531	17	0
2	B	2562	0	2529	13	0
2	C	2565	0	2540	16	0
2	D	2528	0	2503	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	25	0	0	1	0
5	B	25	0	0	0	0
5	C	25	0	0	1	0
5	D	25	0	0	1	0
6	A	145	0	0	0	0
6	B	166	0	0	1	0
6	C	155	0	0	1	0
6	D	99	0	0	0	0
All	All	10883	0	10103	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:714:MET:HE2	6:C:1009:HOH:O	1.89	0.73
1:A:461:GLN:HE22	1:A:500:THR:HG21	1.60	0.67
2:B:497:SER:O	2:B:553:ARG:HD2	2.00	0.61
2:C:693:TYR:OH	5:C:803:K2L:N5	2.35	0.60
2:D:492:TYR:CZ	2:D:496:ARG:HD2	2.39	0.57
2:B:727:LEU:HD12	2:B:759:LEU:HD11	1.87	0.57
2:C:716:ARG:O	2:C:719:LYS:CG	2.54	0.55
1:A:497:SER:OG	1:A:557:LEU:HD11	2.09	0.52
1:A:735:ILE:HB	1:A:736:PRO:HD3	1.92	0.51
2:C:709:GLN:HE21	2:C:710:PRO:HD2	1.76	0.51
2:C:716:ARG:O	2:C:719:LYS:HG3	2.11	0.50
1:A:497:SER:O	1:A:553:ARG:HD2	2.11	0.50
1:A:496:ARG:NH1	1:A:538:TYR:OH	2.43	0.49
2:C:496:ARG:HD3	2:C:538:TYR:OH	2.13	0.48
2:C:749:GLU:HB3	2:C:750:PRO:HD3	1.95	0.48
2:D:497:SER:O	2:D:553:ARG:HD2	2.14	0.48
2:C:716:ARG:O	2:C:719:LYS:HG2	2.13	0.48
2:B:469:CYS:SG	6:B:1037:HOH:O	2.60	0.48
2:B:497:SER:HA	2:B:542:GLN:NE2	2.28	0.48
2:B:727:LEU:HD12	2:B:759:LEU:CD1	2.43	0.48
2:C:464:LEU:O	2:C:465:PRO:C	2.57	0.48
2:B:497:SER:OG	2:B:557:LEU:HD11	2.14	0.48
2:C:700:GLY:HA3	2:C:714:MET:O	2.14	0.47
1:A:700:GLY:HA3	1:A:714:MET:O	2.15	0.47
5:A:803:K2L:O25	5:A:803:K2L:C7	2.62	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:523:PRO:HD2	2:C:695:GLU:HG2	1.97	0.47
1:A:749:GLU:N	1:A:750:PRO:CD	2.78	0.46
2:B:638:TYR:CD1	2:B:671:THR:HG21	2.50	0.46
2:B:700:GLY:HA3	2:B:714:MET:O	2.15	0.46
2:B:677[A]:SER:HB2	2:B:688:THR:HG21	1.98	0.46
2:B:497:SER:HA	2:B:542:GLN:HE22	1.81	0.46
2:C:467:ARG:O	2:C:471:GLU:HB2	2.16	0.46
1:A:727:LEU:HD21	1:A:763:GLU:HG3	1.98	0.45
1:A:550:ASP:HA	1:A:553:ARG:NH1	2.32	0.44
1:A:492:TYR:CZ	1:A:496:ARG:HD2	2.53	0.44
2:B:550:ASP:OD1	2:B:550:ASP:N	2.44	0.43
2:C:638:TYR:CD1	2:C:671:THR:HG21	2.54	0.43
1:A:550:ASP:HA	1:A:553:ARG:HH12	1.84	0.43
1:A:660:SER:O	1:A:664:ARG:HG3	2.19	0.43
2:B:716:ARG:O	2:B:719:LYS:HB2	2.19	0.43
1:A:547:LEU:HD12	1:A:745:LEU:CD2	2.48	0.42
1:A:477:PHE:HB3	1:A:535:HIS:CE1	2.54	0.42
1:A:493:MET:SD	1:A:535:HIS:HA	2.59	0.42
1:A:461:GLN:NE2	1:A:500:THR:HG21	2.32	0.42
1:A:542:GLN:NE2	1:A:542:GLN:HA	2.35	0.41
2:D:693:TYR:OH	5:D:803:K2L:N5	2.53	0.41
2:C:592:GLU:HA	2:C:595:HIS:CD2	2.55	0.41
2:C:697:TRP:HB3	2:C:716:ARG:HB3	2.02	0.41
2:C:619:TYR:C	2:C:619:TYR:CD1	2.99	0.41
2:B:493:MET:SD	2:B:535:HIS:HA	2.60	0.40
2:D:497:SER:OG	2:D:557:LEU:HD11	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	262/343 (76%)	256 (98%)	6 (2%)	0	100	100
2	D	309/343 (90%)	300 (97%)	9 (3%)	0	100	100
All	All	571/686 (83%)	556 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	22/305 (7%)	22 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	D	761	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CME	B	509	2	8,9,10	0.48	0	6,9,11	0.60	0
1	CME	A	469	1	8,9,10	0.60	0	6,9,11	1.04	1 (16%)
2	CME	C	509	2	8,9,10	0.55	0	6,9,11	0.68	0
2	CME	D	509	2	8,9,10	0.57	0	6,9,11	0.82	0
1	CME	A	509	1	8,9,10	0.39	0	6,9,11	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CME	B	509	2	-	0/5/8/10	-
1	CME	A	469	1	-	0/5/8/10	-
2	CME	C	509	2	-	2/5/8/10	-
2	CME	D	509	2	-	1/5/8/10	-
1	CME	A	509	1	-	2/5/8/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	469	CME	CB-CA-C	2.05	116.36	110.80

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	509	CME	SD-CE-CZ-OH
2	C	509	CME	SD-CE-CZ-OH
2	D	509	CME	SD-CE-CZ-OH
1	A	509	CME	CZ-CE-SD-SG
2	C	509	CME	CZ-CE-SD-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	K2L	A	803	-	27,28,28	1.81	6 (22%)	27,41,41	1.72	4 (14%)
5	K2L	B	803	-	27,28,28	2.07	9 (33%)	27,41,41	1.78	4 (14%)
5	K2L	D	803	-	27,28,28	1.81	7 (25%)	27,41,41	1.77	8 (29%)
5	K2L	C	803	-	27,28,28	2.20	8 (29%)	27,41,41	1.54	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	K2L	A	803	-	-	2/5/19/19	0/4/4/4
5	K2L	B	803	-	-	2/5/19/19	0/4/4/4
5	K2L	D	803	-	-	0/5/19/19	0/4/4/4
5	K2L	C	803	-	-	1/5/19/19	0/4/4/4

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	803	K2L	C4-N8	5.49	1.41	1.36
5	A	803	K2L	C22-N2	-4.80	1.40	1.47
5	D	803	K2L	C15-N12	4.66	1.37	1.30
5	C	803	K2L	C4-N5	4.43	1.40	1.32
5	C	803	K2L	C3-N18	4.22	1.43	1.37
5	C	803	K2L	C15-N12	4.21	1.37	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	803	K2L	N16-N19	-4.19	1.30	1.38
5	C	803	K2L	C17-N16	3.75	1.44	1.38
5	B	803	K2L	C15-N12	3.62	1.36	1.30
5	A	803	K2L	C4-N8	3.59	1.39	1.36
5	D	803	K2L	O25-C24	-3.44	1.15	1.23
5	D	803	K2L	C4-N8	3.43	1.39	1.36
5	C	803	K2L	C10-C4	3.38	1.53	1.49
5	A	803	K2L	C24-N2	3.28	1.43	1.38
5	C	803	K2L	C24-N2	2.98	1.43	1.38
5	D	803	K2L	C20-C14	-2.89	1.43	1.50
5	A	803	K2L	C15-N12	2.87	1.35	1.30
5	B	803	K2L	C10-C4	2.87	1.53	1.49
5	A	803	K2L	C11-C3	-2.79	1.46	1.49
5	C	803	K2L	C4-N8	2.69	1.38	1.36
5	B	803	K2L	C13-N12	2.68	1.43	1.36
5	C	803	K2L	C21-C15	2.62	1.53	1.49
5	B	803	K2L	C4-N5	2.59	1.37	1.32
5	B	803	K2L	O25-C24	-2.46	1.17	1.23
5	D	803	K2L	C4-N5	2.28	1.36	1.32
5	B	803	K2L	C9-N8	2.20	1.51	1.46
5	D	803	K2L	C6-N2	-2.15	1.37	1.41
5	D	803	K2L	C6-N5	2.12	1.38	1.35
5	A	803	K2L	C11-C10	2.03	1.59	1.51
5	B	803	K2L	C17-N18	-2.00	1.29	1.33

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	803	K2L	C11-C10-C4	-5.74	106.19	111.62
5	A	803	K2L	C11-C10-C4	-4.79	107.09	111.62
5	B	803	K2L	N16-C17-N18	-4.27	106.23	110.13
5	A	803	K2L	N16-C17-N18	-4.20	106.29	110.13
5	D	803	K2L	C14-C17-N18	4.06	135.46	129.41
5	D	803	K2L	N16-C17-N18	-3.52	106.92	110.13
5	C	803	K2L	C6-N5-C4	-3.24	102.69	104.37
5	A	803	K2L	O25-C24-N2	-3.21	120.50	123.94
5	D	803	K2L	C1-C24-N2	3.12	110.74	108.03
5	C	803	K2L	C14-C17-N18	2.77	133.54	129.41
5	C	803	K2L	C11-C10-C4	-2.73	109.04	111.62
5	D	803	K2L	C6-N5-C4	-2.71	102.96	104.37
5	C	803	K2L	O25-C24-N2	2.63	126.76	123.94
5	C	803	K2L	C21-C15-N12	2.57	124.27	119.26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	803	K2L	C21-C15-N12	2.48	124.11	119.26
5	B	803	K2L	C14-C17-N16	2.44	121.87	118.28
5	D	803	K2L	N8-C4-N5	2.39	113.37	111.17
5	A	803	K2L	C14-C17-N18	2.34	132.89	129.41
5	B	803	K2L	C23-C22-N2	2.30	105.99	102.98
5	D	803	K2L	C11-C3-N18	2.15	125.11	120.27
5	C	803	K2L	N8-C4-N5	2.11	113.11	111.17
5	D	803	K2L	C21-C15-N16	-2.07	115.06	120.05
5	C	803	K2L	N16-C17-N18	-2.02	108.28	110.13

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	803	K2L	C10-C11-C3-N19
5	B	803	K2L	C10-C11-C3-N19
5	B	803	K2L	C10-C11-C3-N18
5	C	803	K2L	C10-C11-C3-N19
5	A	803	K2L	C10-C11-C3-N18


There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	803	K2L	1	0
5	D	803	K2L	1	0
5	C	803	K2L	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand K2L A 803




Ligand K2L B 803



Ligand K2L D 803





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	310/343 (90%)	0.14	7 (2%) 61 70	16, 33, 58, 89	2 (0%)
2	B	314/343 (91%)	0.11	8 (2%) 58 68	20, 31, 57, 94	3 (0%)
2	C	313/343 (91%)	0.11	7 (2%) 62 71	19, 33, 57, 98	4 (1%)
2	D	310/343 (90%)	0.36	8 (2%) 57 67	27, 39, 61, 75	1 (0%)
All	All	1247/1372 (90%)	0.18	30 (2%) 59 69	16, 34, 60, 98	10 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	768	GLY	5.3
2	B	457	GLN	5.2
1	A	459	LEU	5.2
2	B	771	THR	4.2
2	D	500	THR	3.7
2	D	652	GLY	3.6
2	B	458	GLY	3.6
2	C	459	LEU	3.6
2	D	459	LEU	3.4
2	C	462	PHE	2.9
2	B	459	LEU	2.8
2	B	460	MET	2.8
1	A	545	HIS	2.6
2	C	617	SER	2.6
2	C	458	GLY	2.6
2	D	458	GLY	2.6
1	A	770	GLU	2.5
1	A	460	MET	2.5
2	C	613	THR	2.4
2	B	461	GLN	2.4
2	C	457	GLN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	462	PHE	2.3
2	D	654	LEU	2.3
1	A	461	GLN	2.3
1	A	643	LYS	2.3
2	D	612	SER	2.2
2	C	614	LEU	2.1
2	B	719	LYS	2.1
2	B	550	ASP	2.1
2	D	462	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	A	509	10/11	0.85	0.16	37,46,87,92	0
2	CME	D	509	10/11	0.85	0.16	38,59,97,106	0
2	CME	B	509	10/11	0.88	0.14	32,41,83,89	0
2	CME	C	509	10/11	0.89	0.12	37,48,74,75	0
1	CME	A	469	10/11	0.95	0.11	29,36,58,58	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	K2L	B	803	25/25	0.95	0.06	24,28,31,35	0
5	K2L	A	803	25/25	0.96	0.06	24,28,35,39	0
5	K2L	C	803	25/25	0.96	0.06	26,30,37,38	0
5	K2L	D	803	25/25	0.96	0.06	29,34,40,46	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	D	802	1/1	0.98	0.03	34,34,34,34	0
4	MG	C	802	1/1	0.99	0.02	22,22,22,22	0
4	MG	A	802	1/1	0.99	0.01	24,24,24,24	0
4	MG	B	802	1/1	0.99	0.02	23,23,23,23	0
3	ZN	A	801	1/1	1.00	0.01	28,28,28,28	0
3	ZN	B	801	1/1	1.00	0.01	26,26,26,26	0
3	ZN	C	801	1/1	1.00	0.01	28,28,28,28	0
3	ZN	D	801	1/1	1.00	0.02	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around K2L B 803:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around K2L A 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around K2L C 803:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.