



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 07:48 AM UTC

PDB ID : 7SJ9 / pdb_00007sj9
EMDB ID : EMD-25159
Title : 13pf E254A microtubule from recombinant human tubulin decorated with EB3
Authors : LaFrance, B.J.; Greber, B.J.; Zhang, R.; McCollum, C.; Nogales, E.
Deposited on : 2021-10-16
Resolution : 3.80 Å(reported)
Based on initial model : 6DPU

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMD archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	10198 (3.30 - 4.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	11% 64% 31% 5%
1	C	457	13% 63% 32% 5%
1	E	457	11% 64% 31% 5%
1	J	457	41% 67% 28% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	457	
1	L	457	
2	B	456	
2	D	456	
2	F	456	
2	G	456	
2	H	456	
2	I	456	
3	M	281	
3	N	281	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 43240 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	435	Total	C	N	O	S	0	0
			3414	2163	584	645	22		
1	C	435	Total	C	N	O	S	0	0
			3414	2163	584	645	22		
1	E	435	Total	C	N	O	S	0	0
			3414	2163	584	645	22		
1	J	435	Total	C	N	O	S	0	0
			3414	2163	584	645	22		
1	K	435	Total	C	N	O	S	0	0
			3414	2163	584	645	22		
1	L	435	Total	C	N	O	S	0	0
			3414	2163	584	645	22		

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	37F	HIS	-	insertion	UNP P68363
A	37G	HIS	-	insertion	UNP P68363
A	37H	HIS	-	insertion	UNP P68363
A	40	HIS	-	insertion	UNP P68363
A	41	HIS	-	insertion	UNP P68363
A	42	HIS	-	insertion	UNP P68363
A	254	ALA	GLU	engineered mutation	UNP P68363
C	37F	HIS	-	insertion	UNP P68363
C	37G	HIS	-	insertion	UNP P68363
C	37H	HIS	-	insertion	UNP P68363
C	40	HIS	-	insertion	UNP P68363
C	41	HIS	-	insertion	UNP P68363
C	42	HIS	-	insertion	UNP P68363
C	254	ALA	GLU	engineered mutation	UNP P68363
E	37F	HIS	-	insertion	UNP P68363
E	37G	HIS	-	insertion	UNP P68363
E	37H	HIS	-	insertion	UNP P68363
E	40	HIS	-	insertion	UNP P68363

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	41	HIS	-	insertion	UNP P68363
E	42	HIS	-	insertion	UNP P68363
E	254	ALA	GLU	engineered mutation	UNP P68363
J	37F	HIS	-	insertion	UNP P68363
J	37G	HIS	-	insertion	UNP P68363
J	37H	HIS	-	insertion	UNP P68363
J	40	HIS	-	insertion	UNP P68363
J	41	HIS	-	insertion	UNP P68363
J	42	HIS	-	insertion	UNP P68363
J	254	ALA	GLU	engineered mutation	UNP P68363
K	37F	HIS	-	insertion	UNP P68363
K	37G	HIS	-	insertion	UNP P68363
K	37H	HIS	-	insertion	UNP P68363
K	40	HIS	-	insertion	UNP P68363
K	41	HIS	-	insertion	UNP P68363
K	42	HIS	-	insertion	UNP P68363
K	254	ALA	GLU	engineered mutation	UNP P68363
L	37F	HIS	-	insertion	UNP P68363
L	37G	HIS	-	insertion	UNP P68363
L	37H	HIS	-	insertion	UNP P68363
L	40	HIS	-	insertion	UNP P68363
L	41	HIS	-	insertion	UNP P68363
L	42	HIS	-	insertion	UNP P68363
L	254	ALA	GLU	engineered mutation	UNP P68363

- Molecule 2 is a protein called Tubulin beta-3 chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	429	Total 3367	C 2117	N 577	O 648	S 25	0	0
2	D	429	Total 3367	C 2117	N 577	O 648	S 25	0	0
2	F	429	Total 3367	C 2117	N 577	O 648	S 25	0	0
2	G	429	Total 3367	C 2117	N 577	O 648	S 25	0	0
2	H	429	Total 3367	C 2117	N 577	O 648	S 25	0	0
2	I	429	Total 3367	C 2117	N 577	O 648	S 25	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	451	GLU	-	expression tag	UNP Q13509
B	452	ASN	-	expression tag	UNP Q13509
B	453	LEU	-	expression tag	UNP Q13509
B	454	TYR	-	expression tag	UNP Q13509
B	455	PHE	-	expression tag	UNP Q13509
B	456	GLN	-	expression tag	UNP Q13509
D	451	GLU	-	expression tag	UNP Q13509
D	452	ASN	-	expression tag	UNP Q13509
D	453	LEU	-	expression tag	UNP Q13509
D	454	TYR	-	expression tag	UNP Q13509
D	455	PHE	-	expression tag	UNP Q13509
D	456	GLN	-	expression tag	UNP Q13509
F	451	GLU	-	expression tag	UNP Q13509
F	452	ASN	-	expression tag	UNP Q13509
F	453	LEU	-	expression tag	UNP Q13509
F	454	TYR	-	expression tag	UNP Q13509
F	455	PHE	-	expression tag	UNP Q13509
F	456	GLN	-	expression tag	UNP Q13509
G	451	GLU	-	expression tag	UNP Q13509
G	452	ASN	-	expression tag	UNP Q13509
G	453	LEU	-	expression tag	UNP Q13509
G	454	TYR	-	expression tag	UNP Q13509
G	455	PHE	-	expression tag	UNP Q13509
G	456	GLN	-	expression tag	UNP Q13509
H	451	GLU	-	expression tag	UNP Q13509
H	452	ASN	-	expression tag	UNP Q13509
H	453	LEU	-	expression tag	UNP Q13509
H	454	TYR	-	expression tag	UNP Q13509
H	455	PHE	-	expression tag	UNP Q13509
H	456	GLN	-	expression tag	UNP Q13509
I	451	GLU	-	expression tag	UNP Q13509
I	452	ASN	-	expression tag	UNP Q13509
I	453	LEU	-	expression tag	UNP Q13509
I	454	TYR	-	expression tag	UNP Q13509
I	455	PHE	-	expression tag	UNP Q13509
I	456	GLN	-	expression tag	UNP Q13509

- Molecule 3 is a protein called Microtubule-associated protein RP/EB family member 3.


Mol	Chain	Residues	Atoms					AltConf	Trace
3	M	131	Total	C	N	O	S	0	0
			1079	702	181	189	7		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
3	N	131	Total	C	N	O	S	0	0
			1079	702	181	189	7		

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
4	A	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	B	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	C	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	D	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	E	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	F	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	G	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	H	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	I	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	J	1	Total	C	N	O	P	0
			32	10	5	14	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
4	K	1	Total	C	N	O	P	0
			32	10	5	14	3	
4	L	1	Total	C	N	O	P	0
			32	10	5	14	3	


- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
5	A	1	Total	Mg	0
			1	1	
5	B	1	Total	Mg	0
			1	1	
5	C	1	Total	Mg	0
			1	1	
5	D	1	Total	Mg	0
			1	1	
5	E	1	Total	Mg	0
			1	1	
5	F	1	Total	Mg	0
			1	1	
5	G	1	Total	Mg	0
			1	1	
5	H	1	Total	Mg	0
			1	1	
5	I	1	Total	Mg	0
			1	1	
5	J	1	Total	Mg	0
			1	1	
5	K	1	Total	Mg	0
			1	1	
5	L	1	Total	Mg	0
			1	1	


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain







• Molecule 1: Tubulin alpha-1B chain




• Molecule 1: Tubulin alpha-1B chain







• Molecule 1: Tubulin alpha-1B chain




• Molecule 1: Tubulin alpha-1B chain








• Molecule 2: Tubulin beta-3 chain




• Molecule 2: Tubulin beta-3 chain










• Molecule 2: Tubulin beta-3 chain





• Molecule 2: Tubulin beta-3 chain



- Molecule 3: Microtubule-associated protein RP/EB family member 3



- Molecule 3: Microtubule-associated protein RP/EB family member 3



4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=-0.19°, rise=81.64 Å, axial sym=C1	Depositor
Number of segments used	10008	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; CTFFind4 and CTFRefine were used within the RELION3.1 framework.	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{Å}^2$)	40	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	64000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	15.034	Depositor
Minimum map value	-7.169	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	3.8	Depositor
Map size (Å)	471.04, 471.04, 471.04	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.92, 0.92, 0.92	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/3493	0.36	0/4741
1	C	0.14	0/3493	0.36	0/4741
1	E	0.16	0/3493	0.37	0/4741
1	J	0.14	0/3493	0.35	0/4741
1	K	0.13	0/3493	0.32	0/4741
1	L	0.14	0/3493	0.35	0/4741
2	B	0.15	0/3442	0.38	0/4663
2	D	0.17	0/3442	0.44	2/4663 (0.0%)
2	F	0.15	0/3442	0.40	0/4663
2	G	0.16	0/3442	0.39	1/4663 (0.0%)
2	H	0.16	0/3442	0.42	4/4663 (0.1%)
2	I	0.16	0/3442	0.41	1/4663 (0.0%)
3	M	0.13	0/1107	0.36	0/1490
3	N	0.12	0/1107	0.36	0/1490
All	All	0.15	0/43824	0.38	8/59404 (0.0%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	359	ARG	N-CA-C	10.94	126.49	111.39
2	G	359	ARG	N-CA-C	7.34	121.03	110.10
2	H	358	PRO	CA-C-O	-7.06	115.94	121.38
2	D	358	PRO	CA-C-O	-7.02	115.97	121.38
2	D	358	PRO	N-CA-C	5.73	122.57	113.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3414	0	3320	112	0
1	C	3414	0	3320	103	0
1	E	3414	0	3320	118	0
1	J	3414	0	3320	87	0
1	K	3414	0	3320	92	0
1	L	3414	0	3320	90	0
2	B	3367	0	3251	123	0
2	D	3367	0	3251	134	0
2	F	3367	0	3251	120	0
2	G	3367	0	3251	104	0
2	H	3367	0	3251	114	0
2	I	3367	0	3251	103	0
3	M	1079	0	1065	28	0
3	N	1079	0	1065	20	0
4	A	32	0	12	4	0
4	B	32	0	12	2	0
4	C	32	0	12	4	0
4	D	32	0	12	1	0
4	E	32	0	12	4	0
4	F	32	0	12	0	0
4	G	32	0	12	2	0
4	H	32	0	12	2	0
4	I	32	0	12	2	0
4	J	32	0	12	5	0
4	K	32	0	12	5	0
4	L	32	0	12	3	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	G	1	0	0	0	0
5	H	1	0	0	0	0
5	I	1	0	0	0	0
5	J	1	0	0	0	0
5	K	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	L	1	0	0	0	0
All	All	43240	0	41700	1233	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 1233 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:358:PRO:O	2:D:359:ARG:HG2	1.55	1.05
2:H:358:PRO:O	2:H:359:ARG:HG2	1.54	1.05
1:E:352:LYS:HE2	2:F:99:ASN:HD21	1.35	0.92
2:I:26:ASP:HB3	2:I:359:ARG:NH2	1.87	0.90
1:E:222:PRO:HD2	2:G:324:LYS:HE3	1.58	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	429/457 (94%)	415 (97%)	14 (3%)	0	100	100
1	C	429/457 (94%)	415 (97%)	14 (3%)	0	100	100
1	E	429/457 (94%)	415 (97%)	14 (3%)	0	100	100
1	J	429/457 (94%)	421 (98%)	8 (2%)	0	100	100
1	K	429/457 (94%)	419 (98%)	10 (2%)	0	100	100
1	L	429/457 (94%)	416 (97%)	13 (3%)	0	100	100
2	B	427/456 (94%)	413 (97%)	14 (3%)	0	100	100
2	D	427/456 (94%)	410 (96%)	16 (4%)	1 (0%)	43	73

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	427/456 (94%)	410 (96%)	17 (4%)	0	100	100
2	G	427/456 (94%)	412 (96%)	15 (4%)	0	100	100
2	H	427/456 (94%)	411 (96%)	15 (4%)	1 (0%)	43	73
2	I	427/456 (94%)	410 (96%)	17 (4%)	0	100	100
3	M	129/281 (46%)	119 (92%)	10 (8%)	0	100	100
3	N	129/281 (46%)	121 (94%)	8 (6%)	0	100	100
All	All	5394/6040 (89%)	5207 (96%)	185 (3%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	360	GLY
2	H	360	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/384 (96%)	368 (100%)	0	100	100
1	C	368/384 (96%)	368 (100%)	0	100	100
1	E	368/384 (96%)	368 (100%)	0	100	100
1	J	368/384 (96%)	368 (100%)	0	100	100
1	K	368/384 (96%)	368 (100%)	0	100	100
1	L	368/384 (96%)	368 (100%)	0	100	100
2	B	368/392 (94%)	365 (99%)	3 (1%)	73	76
2	D	368/392 (94%)	367 (100%)	1 (0%)	86	84
2	F	368/392 (94%)	367 (100%)	1 (0%)	86	84
2	G	368/392 (94%)	368 (100%)	0	100	100
2	H	368/392 (94%)	368 (100%)	0	100	100
2	I	368/392 (94%)	368 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	M	116/249 (47%)	115 (99%)	1 (1%)	70	74
3	N	116/249 (47%)	116 (100%)	0	100	100
All	All	4648/5154 (90%)	4642 (100%)	6 (0%)	87	89

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	298	ASN
2	F	359	ARG
3	M	118	ASN
2	B	359	ARG
2	B	298	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 95 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	247	ASN
1	J	197	HIS
2	H	370	ASN
2	I	256	ASN
1	K	15	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 12 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GTP	B	501	5	33,34,34	0.93	1 (3%)	50,54,54	1.63	9 (18%)
4	GTP	C	501	5	33,34,34	0.92	1 (3%)	50,54,54	1.59	9 (18%)
4	GTP	L	501	5	33,34,34	0.96	1 (3%)	50,54,54	1.59	9 (18%)
4	GTP	A	501	5	33,34,34	0.92	1 (3%)	50,54,54	1.59	9 (18%)
4	GTP	J	501	5	33,34,34	0.94	1 (3%)	50,54,54	1.60	9 (18%)
4	GTP	D	501	5	33,34,34	0.94	1 (3%)	50,54,54	1.63	9 (18%)
4	GTP	F	501	5	33,34,34	0.93	1 (3%)	50,54,54	1.61	9 (18%)
4	GTP	K	501	5	33,34,34	0.93	1 (3%)	50,54,54	1.59	9 (18%)
4	GTP	E	501	5	33,34,34	0.92	1 (3%)	50,54,54	1.59	9 (18%)
4	GTP	G	501	5	33,34,34	0.94	1 (3%)	50,54,54	1.62	9 (18%)
4	GTP	I	501	5	33,34,34	0.94	1 (3%)	50,54,54	1.62	9 (18%)
4	GTP	H	501	5	33,34,34	0.92	1 (3%)	50,54,54	1.61	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	B	501	5	-	6/22/38/38	0/3/3/3
4	GTP	C	501	5	-	5/22/38/38	0/3/3/3
4	GTP	L	501	5	-	4/22/38/38	0/3/3/3
4	GTP	A	501	5	-	5/22/38/38	0/3/3/3
4	GTP	J	501	5	-	5/22/38/38	0/3/3/3
4	GTP	D	501	5	-	5/22/38/38	0/3/3/3
4	GTP	F	501	5	-	6/22/38/38	0/3/3/3
4	GTP	K	501	5	-	5/22/38/38	0/3/3/3
4	GTP	E	501	5	-	5/22/38/38	0/3/3/3
4	GTP	G	501	5	-	5/22/38/38	0/3/3/3
4	GTP	I	501	5	-	6/22/38/38	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	H	501	5	-	6/22/38/38	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	501	GTP	C2-N3	2.21	1.38	1.33
4	B	501	GTP	C2-N3	2.21	1.38	1.33
4	F	501	GTP	C2-N3	2.21	1.38	1.33
4	J	501	GTP	C2-N3	2.20	1.38	1.33
4	L	501	GTP	C2-N3	2.20	1.38	1.33

The worst 5 of 108 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	501	GTP	C5-C4-N3	-5.15	120.20	128.39
4	F	501	GTP	C5-C4-N3	-5.12	120.24	128.39
4	B	501	GTP	C5-C4-N3	-5.12	120.25	128.39
4	K	501	GTP	C5-C4-N3	-5.09	120.28	128.39
4	A	501	GTP	C5-C4-N3	-5.07	120.32	128.39

There are no chirality outliers.

5 of 63 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	GTP	C5'-O5'-PA-O3A
4	A	501	GTP	C5'-O5'-PA-O1A
4	A	501	GTP	C5'-O5'-PA-O2A
4	B	501	GTP	C5'-O5'-PA-O3A
4	B	501	GTP	C5'-O5'-PA-O1A

There are no ring outliers.

11 monomers are involved in 34 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	501	GTP	2	0
4	C	501	GTP	4	0
4	L	501	GTP	3	0
4	A	501	GTP	4	0
4	J	501	GTP	5	0
4	D	501	GTP	1	0
4	K	501	GTP	5	0



Continued on next page...



Continued from previous page...



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	501	GTP	4	0
4	G	501	GTP	2	0
4	I	501	GTP	2	0
4	H	501	GTP	2	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-25159. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256



Y Index: 256




Z Index: 256


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 361



Y Index: 358




Z Index: 257


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.8. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1330 nm^3 ; this corresponds to an approximate mass of 1201 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.263 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-25159 and PDB model 7SJ9. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 3.8 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3.8).

9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 56% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (3.8) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5620	 0.4410
A	 0.6430	 0.4630
B	 0.6090	 0.4460
C	 0.6210	 0.4480
D	 0.5860	 0.4370
E	 0.6200	 0.4490
F	 0.5850	 0.4320
G	 0.6450	 0.4530
H	 0.6650	 0.4640
I	 0.6440	 0.4520
J	 0.4410	 0.4180
K	 0.4760	 0.4290
L	 0.4510	 0.4170
M	 0.1570	 0.4130
N	 0.1950	 0.4210

