



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 03:42 PM UTC

PDB ID : 8SJS / pdb\_00008sjs  
Title : [3T18] Self-assembling right-handed tensegrity triangle with 18 interjunction base pairs and P63 symmetry  
Authors : Janowski, J.; Vecchioni, S.; Sha, R.; Ohayon, Y.P.  
Deposited on : 2023-04-18  
Resolution : 6.31 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 6.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1154 (8.50-4.00)
Clashscore	190562	1016 (8.60-4.02)
RSRZ outliers	180081	1147 (8.50-4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	19	100%
2	B	18	6% 94%
3	C	13	23% 77%
4	D	12	8% 83% 8%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 1265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*CP\*AP\*GP\*AP\*GP\*CP\*CP\*TP\*GP\*AP\*CP\*AP\*TP\*AP\*CP\*CP\*GP\*CP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	19	384	183	75	108	18	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(P\*TP\*GP\*CP\*GP\*GP\*TP\*AP\*TP\*GP\*TP\*CP\*AP\*CP\*CP\*AP\*CP\*GP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	18	369	175	68	108	18	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(\*TP\*GP\*CP\*GP\*CP\*TP\*GP\*TP\*GP\*GP\*CP\*TP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	13	263	126	45	80	12	0	0	0


- Molecule 4 is a DNA chain called DNA (5'-D(P\*TP\*CP\*GP\*TP\*GP\*GP\*AP\*CP\*AP\*GP\*CP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	12	249	117	48	72	12	0	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(\*CP\*AP\*GP\*AP\*GP\*CP\*CP\*TP\*GP\*AP\*CP\*AP\*TP\*AP\*CP\*CP\*GP\*CP\*A)-3')

Chain A:  100%


T101  
A102  
G103  
A104  
G105  
C106  
G107  
T108  
G109  
A110  
C111  
A112  
T113  
A114  
C115  
C116  
G117  
C118  
A119

- Molecule 2: DNA (5'-D(P\*TP\*GP\*CP\*GP\*GP\*TP\*AP\*TP\*GP\*TP\*CP\*AP\*CP\*CP\*AP\*CP\*GP\*A)-3')

Chain B:  6% 94%


T101  
G102  
C103  
G104  
G105  
T106  
A107  
T108  
G109  
T110  
C111  
A112  
C113  
C114  
A115  
C116  
G117  
A118

- Molecule 3: DNA (5'-D(\*TP\*GP\*CP\*GP\*CP\*TP\*GP\*TP\*GP\*GP\*CP\*TP\*C)-3')

Chain C:  23% 77%

T101  
G102  
G103  
G104  
G105  
T106  
G107  
T108  
G109  
G110  
C111  
T112  
C113

- Molecule 4: DNA (5'-D(P\*TP\*CP\*GP\*TP\*GP\*GP\*AP\*CP\*AP\*GP\*CP\*G)-3')

Chain D:  8% 83% 8%

T101  
C102  
G103  
T104  
G105  
G106  
A107  
C108  
A109  
G110  
C111  
G112

## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	194.20Å 194.20Å 55.43Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.78 – 6.31 41.78 – 6.31	Depositor EDS
% Data completeness (in resolution range)	73.8 (41.78-6.31) 66.4 (41.78-6.31)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.73 (at 6.19Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.110 , 0.119 0.112 , 0.121	Depositor DCC
$R_{free}$ test set	99 reflections (3.62%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	303.5	Xtrriage
Anisotropy	1.152	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.02 , 0.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.087 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.99	EDS
Total number of atoms	1265	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	547.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.87	0/431	0.98	0/662
2	B	0.63	0/413	0.80	0/635
3	C	0.65	0/293	0.69	0/451
4	D	0.60	0/279	0.95	1/429 (0.2%)
All	All	0.71	0/1416	0.87	1/2177 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	109	DA	C3'-C2'-C1'	-5.09	93.97	101.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	384	0	213	32	2
2	B	369	0	203	45	2
3	C	263	0	149	23	0
4	D	249	0	135	9	2
All	All	1265	0	700	96	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 50.

The worst 5 of 96 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:DC:N3	3:C:110:DG:N2	2.06	1.04
1:A:118:DC:H2'	1:A:119:DA:C5	2.06	0.89
3:C:109:DG:N3	3:C:109:DG:H2'	1.96	0.79
1:A:116:DC:H1'	1:A:117:DG:C8	2.19	0.78
1:A:108:DT:H2'	1:A:109:DG:C8	2.22	0.74

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:DA:O3'	4:D:101:DT:OP2[2_565]	1.18	1.02
2:B:101:DT:P	2:B:118:DA:O3'[2_565]	1.60	0.60
1:A:119:DA:O3'	4:D:101:DT:P[2_565]	1.63	0.57
2:B:101:DT:OP1	2:B:118:DA:O3'[2_565]	1.68	0.52

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

#### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

#### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	19/19 (100%)	-0.15	0 100 100	483, 556, 608, 611	0
2	B	18/18 (100%)	0.05	0 100 100	482, 550, 591, 602	0
3	C	13/13 (100%)	-0.04	0 100 100	442, 517, 570, 575	0
4	D	12/12 (100%)	0.06	0 100 100	456, 581, 672, 679	0
All	All	62/62 (100%)	-0.03	0 100 100	442, 549, 611, 679	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.