



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:06 AM UTC

PDB ID : 3SKH / pdb_00003skh
Title : I. Novel HCV NS5B Polymerase Inhibitors: Discovery of Indole 2- Carboxylic Acids with C3-Heterocycles
Authors : Lesburg, C.A.; Anilkumar, G.N.
Deposited on : 2011-06-22
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	576	% 86% 11% ..
1	B	576	6% 84% 12% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9204 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HCV NS5B RNA_DEPENDENT RNA POLYMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	558	Total	C	N	O	S	0	0	0
			4347	2739	769	807	32			
1	B	558	Total	C	N	O	S	0	2	0
			4353	2742	769	809	33			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	440	GLY	GLU	conflict	UNP O92972
A	520	ILE	THR	conflict	UNP O92972
A	571	GLU	-	expression tag	UNP O92972
A	572	ASN	-	expression tag	UNP O92972
A	573	LEU	-	expression tag	UNP O92972
A	574	TYR	-	expression tag	UNP O92972
A	575	PHE	-	expression tag	UNP O92972
A	576	GLN	-	expression tag	UNP O92972
B	440	GLY	GLU	conflict	UNP O92972
B	520	ILE	THR	conflict	UNP O92972
B	571	GLU	-	expression tag	UNP O92972
B	572	ASN	-	expression tag	UNP O92972
B	573	LEU	-	expression tag	UNP O92972
B	574	TYR	-	expression tag	UNP O92972
B	575	PHE	-	expression tag	UNP O92972
B	576	GLN	-	expression tag	UNP O92972

- Molecule 2 is 1-benzyl-5-chloro-3-(2-fluorophenyl)-1H-indole-2-carboxylic acid (CCD ID: 058) (formula: C₂₂H₁₅ClFNO₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Cl	F	N	O	0	0
			27	22	1	1	1	2		
2	B	1	Total	C	Cl	F	N	O	0	0
			27	22	1	1	1	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	230	Total	O	0	0
			230	230		
3	B	220	Total	O	0	0
			220	220		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HCV NS5B RNA_DEPENDENT RNA POLYMERASE



• Molecule 1: HCV NS5B RNA_DEPENDENT RNA POLYMERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.44Å 107.63Å 136.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.04 – 2.50 50.04 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.7 (50.04-2.50) 98.7 (50.04-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.51Å)	Xtriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.199 , 0.265 0.209 , 0.277	Depositor DCC
R_{free} test set	2286 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	40.8	Xtriage
Anisotropy	0.685	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9204	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 058

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	2/4441 (0.0%)	1.37	17/6026 (0.3%)
1	B	0.87	0/4455	1.37	19/6045 (0.3%)
All	All	0.87	2/8896 (0.0%)	1.37	36/12071 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	494	VAL	C-N	6.19	1.39	1.33
1	A	348	ALA	C-N	5.24	1.39	1.33

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	261	TYR	N-CA-C	6.55	119.42	111.82
1	A	201	VAL	N-CA-CB	6.45	118.10	110.55
1	B	33	HIS	CA-C-N	6.35	128.79	120.28
1	B	33	HIS	C-N-CA	6.35	128.79	120.28
1	B	329	THR	N-CA-C	6.11	117.61	111.07
1	A	507	VAL	N-CA-CB	5.94	117.50	110.55
1	B	351	GLY	N-CA-C	-5.92	104.82	110.21
1	B	328	GLY	N-CA-C	-5.78	105.82	112.29
1	B	196	SER	N-CA-C	-5.66	101.90	110.39
1	B	244	ASP	N-CA-C	-5.58	100.71	109.25
1	A	165	LEU	CA-C-N	5.56	126.27	119.99
1	A	165	LEU	C-N-CA	5.56	126.27	119.99
1	A	479	PRO	CA-C-N	5.54	126.25	119.99
1	A	479	PRO	C-N-CA	5.54	126.25	119.99
1	B	105	ALA	N-CA-C	5.51	118.00	111.33
1	B	444	ASP	CA-CB-CG	5.50	118.10	112.60
1	A	63	HIS	CA-C-N	5.49	127.58	120.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63	HIS	C-N-CA	5.49	127.58	120.44
1	A	559	ASP	CA-CB-CG	5.46	118.06	112.60
1	B	261	TYR	N-CA-C	5.43	117.20	111.28
1	A	129	ASP	CA-CB-CG	5.38	117.98	112.60
1	A	175	LEU	N-CA-C	5.36	120.83	113.97
1	B	543	SER	CA-C-N	5.30	127.91	120.28
1	B	543	SER	C-N-CA	5.30	127.91	120.28
1	A	239	ILE	N-CA-C	-5.23	105.30	111.00
1	B	104	GLY	CA-C-N	5.22	127.53	120.38
1	B	104	GLY	C-N-CA	5.22	127.53	120.38
1	B	500	TRP	CA-C-N	5.14	127.12	120.44
1	B	500	TRP	C-N-CA	5.14	127.12	120.44
1	B	330	GLN	CA-C-N	5.11	127.55	120.29
1	B	330	GLN	C-N-CA	5.11	127.55	120.29
1	A	230	GLU	CA-C-N	5.10	127.07	120.44
1	A	230	GLU	C-N-CA	5.10	127.07	120.44
1	A	351	GLY	N-CA-C	-5.08	101.14	113.18
1	B	309	GLN	N-CA-C	5.08	118.08	109.76
1	A	244	ASP	N-CA-C	-5.04	102.18	109.59

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4347	0	4365	18	0
1	B	4353	0	4366	24	0
2	A	27	0	14	0	0
2	B	27	0	14	0	0
3	A	230	0	0	1	0
3	B	220	0	0	0	0
All	All	9204	0	8759	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:321:VAL:HG21	1:B:365:SER:HB2	1.66	0.76
1:B:321:VAL:HG21	1:B:365:SER:CB	2.21	0.69
1:A:527:ASN:HD21	1:A:534:LEU:H	1.43	0.66
1:B:389:THR:HG23	1:B:492:LEU:HD21	1.83	0.61
1:B:466:LEU:HD21	1:B:547:LEU:HD11	1.82	0.61
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.84	0.59
1:B:46:SER:HA	1:B:49:GLN:HE21	1.68	0.58
1:A:201:VAL:HG13	1:A:384:LEU:HD13	1.85	0.58
1:A:389:THR:HG23	1:A:492:LEU:HD21	1.86	0.57
1:A:405:ILE:HG13	1:A:446:GLN:HE21	1.70	0.55
1:A:501:ARG:O	1:A:505:ARG:HG2	2.08	0.54
1:B:434:LEU:HD11	1:B:511:LEU:HG	1.90	0.53
1:B:309:GLN:O	1:B:324:CYS:HB2	2.09	0.52
1:B:234:ARG:HG3	1:B:262:ILE:HD11	1.92	0.52
1:A:82:LEU:HD13	1:A:249:ALA:HB2	1.92	0.51
1:A:105:ALA:O	1:A:109:ARG:HG3	2.12	0.48
1:A:17:GLU:HB3	1:A:41:THR:HG22	1.95	0.48
1:A:7:THR:HG21	1:A:273:ASN:ND2	2.29	0.47
1:B:144:VAL:HB	1:B:394:ARG:HG2	1.95	0.47
1:B:447:ILE:HB	1:B:452:TYR:CE1	2.50	0.47
1:A:517:ARG:HA	1:A:520:ILE:HD12	1.99	0.45
1:B:192:GLY:HA3	1:B:316:ASN:OD1	2.16	0.45
1:A:24:ASN:HB3	1:A:27:SER:OG	2.16	0.45
1:B:485:VAL:O	1:B:489:LEU:HG	2.17	0.45
1:B:234:ARG:CG	1:B:262:ILE:HD11	2.47	0.44
1:B:495:PRO:HG2	1:B:503:ARG:HH12	1.82	0.44
1:A:264:GLY:HA3	3:A:603:HOH:O	2.17	0.44
1:B:22:PRO:HG2	1:B:400:ALA:HB1	1.98	0.44
1:B:313:MET:HG2	1:B:322:VAL:HG22	1.99	0.44
1:B:359:ASP:HB3	1:B:362:LEU:HD12	2.00	0.43
1:B:461:GLN:HB3	1:B:542:ALA:HA	2.00	0.43
1:B:48:ARG:NH1	1:B:51:LYS:HD2	2.34	0.43
1:A:542:ALA:HA	1:A:545:LEU:HD12	2.00	0.42
1:B:141:LYS:HD2	1:B:160:ILE:HD11	2.02	0.42
1:A:48:ARG:HG2	1:A:159:LEU:HD13	2.00	0.42
1:A:134:ILE:HG13	1:A:259:ARG:HB3	2.02	0.42
1:A:488:CYS:SG	1:A:492:LEU:HD23	2.60	0.41
1:A:219:TYR:HB3	1:A:320:LEU:HD23	2.02	0.41
1:B:257:THR:HA	1:B:261:TYR:HB2	2.02	0.41
1:B:434:LEU:HG	1:B:439:LEU:HD11	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:ILE:HD13	1:A:169:VAL:HG11	2.02	0.40
1:B:458:ASP:HA	1:B:461:GLN:HE22	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	554/576 (96%)	531 (96%)	23 (4%)	0	100	100
1	B	556/576 (96%)	529 (95%)	25 (4%)	2 (0%)	30	49
All	All	1110/1152 (96%)	1060 (96%)	48 (4%)	2 (0%)	43	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	538	PRO
1	B	532	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	475/490 (97%)	454 (96%)	21 (4%)	25	50
1	B	477/490 (97%)	453 (95%)	24 (5%)	22	44

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	952/980 (97%)	907 (95%)	45 (5%)	23	47

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLU
1	A	57	LEU
1	A	100	LYS
1	A	111	LEU
1	A	121	SER
1	A	184	GLN
1	A	201	VAL
1	A	233	ILE
1	A	234	ARG
1	A	274	CYS
1	A	315	VAL
1	A	336	LEU
1	A	355	GLN
1	A	379	LYS
1	A	384	LEU
1	A	466	LEU
1	A	507	VAL
1	A	535	LYS
1	A	544	GLN
1	A	545	LEU
1	A	556	SER
1	B	23	ILE
1	B	30	LEU
1	B	31	LEU
1	B	35	ASN
1	B	57	LEU
1	B	81	LYS
1	B	83	LEU
1	B	86	GLU
1	B	172	LYS
1	B	184	GLN
1	B	201	VAL
1	B	212	LYS
1	B	262	ILE
1	B	315	VAL
1	B	336	LEU
1	B	364	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	384	LEU
1	B	405	ILE
1	B	434	LEU
1	B	461	GLN
1	B	469	LEU
1	B	484	ARG
1	B	547	LEU
1	B	556	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	ASN
1	A	35	ASN
1	A	184	GLN
1	A	273	ASN
1	A	309	GLN
1	A	446	GLN
1	A	514	GLN
1	A	527	ASN
1	A	544	GLN
1	B	24	ASN
1	B	35	ASN
1	B	49	GLN
1	B	273	ASN
1	B	411	ASN
1	B	428	HIS
1	B	436	GLN
1	B	461	GLN
1	B	514	GLN
1	B	544	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	058	A	577	-	30,30,30	1.27	4 (13%)	40,43,43	1.08	3 (7%)
2	058	B	577	-	30,30,30	1.28	2 (6%)	40,43,43	1.02	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	058	A	577	-	-	6/12/12/12	0/4/4/4
2	058	B	577	-	-	6/12/12/12	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	577	058	C11-C6	-2.74	1.46	1.49
2	A	577	058	C11-C6	-2.33	1.46	1.49
2	B	577	058	C7-C6	-2.29	1.42	1.47
2	A	577	058	C11-C16	2.20	1.41	1.38
2	A	577	058	O1-C10	2.19	1.36	1.30
2	A	577	058	C7-C6	-2.11	1.42	1.47

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	577	058	C8-N1-C5	2.87	111.02	108.20
2	B	577	058	C8-N1-C5	2.57	110.73	108.20
2	A	577	058	C10-C5-N1	2.50	125.70	122.59
2	B	577	058	C10-C5-N1	2.44	125.62	122.59
2	B	577	058	C15-C16-C11	-2.07	120.83	123.07
2	A	577	058	O2-C10-C5	2.07	126.29	120.58
2	B	577	058	C12-C11-C16	2.06	118.99	116.70

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	577	058	O2-C10-C5-N1
2	B	577	058	O1-C10-C5-N1
2	B	577	058	O2-C10-C5-N1
2	B	577	058	O1-C10-C5-C6
2	B	577	058	O2-C10-C5-C6
2	A	577	058	O1-C10-C5-C6
2	A	577	058	O2-C10-C5-C6
2	A	577	058	O1-C10-C5-N1
2	A	577	058	C22-C17-C9-N1
2	B	577	058	C22-C17-C9-N1
2	A	577	058	C18-C17-C9-N1
2	B	577	058	C18-C17-C9-N1

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand 058 A 577




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	558/576 (96%)	0.05	4 (0%) 84 81	29, 43, 65, 93	0
1	B	558/576 (96%)	0.46	32 (5%) 29 26	14, 50, 87, 122	3 (0%)
All	All	1116/1152 (96%)	0.25	36 (3%) 50 46	14, 46, 77, 122	3 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	242	CYS	8.7
1	B	23	ILE	5.2
1	B	26	LEU	5.0
1	B	542	ALA	4.9
1	B	548	SER	4.5
1	B	545	LEU	4.3
1	B	563	SER	3.9
1	B	540	PRO	3.8
1	B	534	LEU	3.8
1	B	451[A]	CYS	3.5
1	B	544	GLN	3.4
1	B	543	SER	3.4
1	B	541	ALA	3.1
1	B	24	ASN	3.0
1	B	547	LEU	2.9
1	B	319[A]	ASP	2.9
1	B	562	HIS	2.8
1	B	529	ALA	2.8
1	B	536	LEU	2.7
1	A	535	LYS	2.7
1	B	14	CYS	2.6
1	B	538	PRO	2.5
1	A	563	SER	2.5
1	B	532	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	501	ARG	2.3
1	B	21	LEU	2.3
1	B	22	PRO	2.3
1	A	402	HIS	2.2
1	B	530	VAL	2.2
1	B	531	ARG	2.2
1	B	25	PRO	2.2
1	B	533	LYS	2.2
1	B	474	LEU	2.1
1	A	547	LEU	2.1
1	B	537	THR	2.1
1	B	561	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	058	A	577	27/27	0.92	0.10	40,48,51,53	0
2	058	B	577	27/27	0.92	0.09	44,54,57,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 058 A 577:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 058 B 577:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.