



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:48 PM UTC

PDB ID : 6SKH / pdb_00006skh
Title : X-ray structure of human glutamate carboxypeptidase II (GCPII) - the E424M inactive mutant, in complex with a inhibitor sulfamide inhibitor GluAsp
Authors : Motlova, L.; Novakova, Z.; Barinka, C.
Deposited on : 2019-08-15
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	6% 89% 9% ..
2	B	3	33% 67%
3	C	2	50% 50%
3	D	2	50% 50%
4	E	4	25% 75%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	5	 60% 40%

2 Entry composition [i](#)

There are 15 unique types of molecules in this entry. The entry contains 7027 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	696	Total	C	N	O	S	0	83	0
			6136	3924	1038	1152	22			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	424	MET	GLU	engineered mutation	UNP Q04609

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	2	Total	Zn	0	0
			2	2		


- Molecule 9 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Ca	0	0
			1	1		

- Molecule 10 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	1	Total	Cl	0	0
			1	1		

- Molecule 11 is beta-D-mannopyranose (CCD ID: BMA) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 12 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 13 is (2 {S})-2-[[[(2 {S})-1,4-bis(oxidanyl)-1,4-bis(oxidanylidene)butan-2-yl]sulfamoylamino]pentanedioic acid (CCD ID: LHK) (formula: C₉H₁₄N₂O₁₀S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
13	A	1	Total	C	N	O	S	0	0
			22	9	2	10	1		

- Molecule 14 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
14	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	591	Total	O	0	0
			591	591		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate carboxypeptidase 2



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 4: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.63Å 130.53Å 159.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.21 – 1.58 49.21 – 1.58	Depositor EDS
% Data completeness (in resolution range)	99.1 (49.21-1.58) 99.2 (49.21-1.58)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.58Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.171 , 0.193 0.178 , 0.205	Depositor DCC
R_{free} test set	4263 reflections (2.95%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.562	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7027	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, CA, CL, BMA, LHK, EDO, NAG, PEG, GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/6376	0.84	4/8623 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	124	HIS	CA-C-N	6.25	126.21	119.78
1	A	124	HIS	C-N-CA	6.25	126.21	119.78
1	A	147	PRO	O-C-N	5.18	123.59	121.15
1	A	368	PRO	N-CA-CB	5.06	108.54	103.48

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	414	ARG	Sidechain
1	A	662[A]	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6136	0	5983	48	0
2	B	39	0	34	0	0
3	C	28	0	25	0	0
3	D	28	0	25	0	0
4	E	50	0	43	2	0
5	F	61	0	52	0	0
6	A	28	0	26	1	0
7	A	16	0	24	0	0
8	A	2	0	0	0	0
9	A	1	0	0	0	0
10	A	1	0	0	0	0
11	A	11	0	10	0	0
12	A	7	0	10	0	0
13	A	22	0	0	0	0
14	A	6	0	8	0	0
15	A	591	0	0	3	0
All	All	7027	0	6240	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:MET:HE1	1:A:586[A]:GLU:HG2	1.44	0.97
1:A:651[B]:GLN:O	1:A:652[B]:ASP:OD1	1.95	0.85
1:A:58:MET:CE	1:A:586[A]:GLU:HG2	2.07	0.84
1:A:188[B]:LEU:HD11	1:A:329:VAL:HG11	1.60	0.80
1:A:655[A]:LYS:HD3	15:A:1386:HOH:O	1.85	0.75
1:A:58:MET:HE1	1:A:586[A]:GLU:CG	2.16	0.74
1:A:400:ARG:O	1:A:404[B]:THR:HG23	1.90	0.72
1:A:131[A]:ILE:HG22	1:A:137:GLU:HG2	1.72	0.70
1:A:395[B]:VAL:HG21	15:A:1375:HOH:O	1.93	0.67
1:A:656[B]:SER:O	1:A:658[B]:PRO:HD3	1.98	0.63
1:A:591[A]:ILE:HG22	1:A:661[A]:LEU:HD21	1.82	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:659[B]:ILE:O	1:A:663[B]:MET:HG3	2.00	0.61
1:A:359[B]:ILE:HD12	1:A:359[B]:ILE:N	2.15	0.60
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.40	0.57
1:A:369:ASP:OD2	1:A:662[B]:ARG:NH2	2.37	0.56
1:A:655[A]:LYS:HE3	4:E:4:MAN:H4	1.87	0.56
1:A:655[A]:LYS:HD2	4:E:4:MAN:O2	2.06	0.55
1:A:486[A]:GLU:H	1:A:486[A]:GLU:CD	2.15	0.54
1:A:517:SER:HB2	1:A:694:PRO:HG3	1.89	0.54
1:A:591[A]:ILE:CG2	1:A:661[A]:LEU:HD21	2.38	0.54
1:A:131[A]:ILE:CG2	1:A:137:GLU:HG2	2.40	0.52
1:A:188[B]:LEU:HD11	1:A:329:VAL:CG1	2.35	0.52
1:A:586[A]:GLU:HA	1:A:586[A]:GLU:OE1	2.10	0.51
1:A:651[B]:GLN:O	1:A:651[B]:GLN:NE2	2.44	0.51
1:A:539:LYS:NZ	1:A:546:PHE:O	2.28	0.50
1:A:653[B]:PHE:O	1:A:655[B]:LYS:HD3	2.11	0.50
1:A:645[B]:LYS:HE3	15:A:924:HOH:O	2.13	0.49
1:A:169:MET:HA	1:A:344:MET:O	2.13	0.48
1:A:198:GLY:O	1:A:223:LYS:HE2	2.12	0.48
1:A:246:TRP:CD1	6:A:809:NAG:H83	2.49	0.47
1:A:58:MET:CE	1:A:586[A]:GLU:CG	2.86	0.47
1:A:92:LEU:O	1:A:96[A]:ILE:HG12	2.15	0.46
1:A:625:TYR:CE1	1:A:722[B]:SER:HA	2.50	0.46
1:A:651[B]:GLN:HE21	1:A:651[B]:GLN:C	2.24	0.45
1:A:171:GLU:HG2	1:A:343:LYS:HG3	1.99	0.45
1:A:551:LEU:HD22	1:A:556:TYR:HB2	1.99	0.44
1:A:645[B]:LYS:HD3	1:A:645[B]:LYS:HA	1.85	0.44
1:A:242:TYR:CG	1:A:243:PRO:HA	2.54	0.42
1:A:325:VAL:HB	1:A:326:PRO:HD2	2.01	0.41
1:A:489[B]:GLU:H	1:A:489[B]:GLU:HG2	1.50	0.41
1:A:70:ILE:HG21	1:A:573:HIS:HB3	2.03	0.41
1:A:392:ALA:HA	1:A:395[B]:VAL:HG22	2.01	0.41
1:A:494:TYR:CE1	1:A:498[B]:THR:HG21	2.56	0.41
1:A:663[A]:MET:HE3	1:A:663[A]:MET:HB2	1.78	0.41
1:A:169:MET:HG2	1:A:345:HIS:HA	2.04	0.40
1:A:444:GLU:OE1	1:A:444:GLU:HA	2.22	0.40
1:A:653[B]:PHE:CG	1:A:654[B]:ASP:N	2.89	0.40
1:A:498[B]:THR:HG22	1:A:510:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	777/707 (110%)	744 (96%)	29 (4%)	4 (0%)	24	9

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	654[A]	ASP
1	A	654[B]	ASP
1	A	382	VAL
1	A	544	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	676/603 (112%)	666 (98%)	10 (2%)	57	31

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	189[A]	GLU
1	A	189[B]	GLU
1	A	303	GLN
1	A	351	GLU
1	A	486[A]	GLU
1	A	486[B]	GLU
1	A	537	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	539	LYS
1	A	557	GLU
1	A	600	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	A	97	GLN
1	A	303	GLN
1	A	451	ASN
1	A	667	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	0.31	0	17,19,21	0.87	0
2	NAG	B	2	2	14,14,15	0.45	0	17,19,21	1.21	2 (11%)
2	BMA	B	3	2	11,11,12	0.61	0	15,15,17	1.24	3 (20%)
3	NAG	C	1	1,3	14,14,15	0.48	0	17,19,21	1.19	2 (11%)
3	NAG	C	2	3	14,14,15	0.35	0	17,19,21	0.60	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	1	1,3	14,14,15	0.48	0	17,19,21	0.74	0
3	NAG	D	2	3	14,14,15	0.35	0	17,19,21	0.85	1 (5%)
4	NAG	E	1	1,4	14,14,15	0.49	0	17,19,21	0.98	2 (11%)
4	NAG	E	2	4	14,14,15	0.35	0	17,19,21	0.80	0
4	BMA	E	3	4	11,11,12	0.33	0	15,15,17	1.12	1 (6%)
4	MAN	E	4	4	11,11,12	0.49	0	15,15,17	0.66	0
5	NAG	F	1	1,5	14,14,15	0.40	0	17,19,21	1.00	1 (5%)
5	NAG	F	2	5	14,14,15	0.41	0	17,19,21	1.19	2 (11%)
5	BMA	F	3	5	11,11,12	0.37	0	15,15,17	0.84	0
5	MAN	F	4	5	11,11,12	0.36	0	15,15,17	0.75	0
5	MAN	F	5	5	11,11,12	0.48	0	15,15,17	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
4	NAG	E	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	E	2	4	-	2/6/23/26	0/1/1/1
4	BMA	E	3	4	-	1/2/19/22	0/1/1/1
4	MAN	E	4	4	-	0/2/19/22	0/1/1/1
5	NAG	F	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	F	2	5	-	2/6/23/26	0/1/1/1
5	BMA	F	3	5	-	0/2/19/22	0/1/1/1
5	MAN	F	4	5	-	0/2/19/22	0/1/1/1
5	MAN	F	5	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	3	BMA	C1-O5-C5	3.33	116.65	112.19
2	B	3	BMA	C1-C2-C3	2.93	113.91	109.64
3	C	1	NAG	C4-C3-C2	2.86	115.20	111.02
2	B	2	NAG	C2-N2-C7	2.74	126.57	122.90
5	F	2	NAG	C1-O5-C5	2.73	115.85	112.19
2	B	2	NAG	C1-O5-C5	2.66	115.76	112.19
3	D	2	NAG	C1-O5-C5	2.65	115.74	112.19
3	C	1	NAG	C1-C2-N2	-2.49	106.50	110.43
2	B	3	BMA	C2-C3-C4	2.30	114.91	110.86
5	F	1	NAG	O5-C1-C2	-2.28	107.76	111.29
2	B	3	BMA	C3-C4-C5	2.25	114.31	110.23
5	F	2	NAG	O4-C4-C5	-2.24	103.80	109.32
4	E	1	NAG	O5-C5-C4	-2.16	105.56	110.83
4	E	1	NAG	O5-C1-C2	-2.12	108.01	111.29

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C8-C7-N2-C2
3	C	1	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
2	B	3	BMA	O5-C5-C6-O6
2	B	3	BMA	C4-C5-C6-O6
4	E	1	NAG	C8-C7-N2-C2
4	E	1	NAG	O7-C7-N2-C2
4	E	2	NAG	C8-C7-N2-C2
4	E	2	NAG	O7-C7-N2-C2
5	F	2	NAG	C8-C7-N2-C2
5	F	2	NAG	O7-C7-N2-C2
5	F	1	NAG	C4-C5-C6-O6
4	E	3	BMA	C4-C5-C6-O6
5	F	1	NAG	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	4	MAN	2	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
13	LHK	A	829	-	21,21,21	2.84	5 (23%)	29,29,29	1.75	7 (24%)
7	EDO	A	827	-	3,3,3	0.47	0	2,2,2	0.22	0
7	EDO	A	828	-	3,3,3	0.40	0	2,2,2	0.45	0
6	NAG	A	809	1	14,14,15	0.63	0	17,19,21	1.20	1 (5%)
7	EDO	A	819	-	3,3,3	0.44	0	2,2,2	0.40	0
14	GOL	A	830	-	5,5,5	0.35	0	5,5,5	0.41	0
6	NAG	A	808	1	14,14,15	0.37	0	17,19,21	1.11	1 (5%)
7	EDO	A	826	-	3,3,3	0.44	0	2,2,2	0.33	0
12	PEG	A	825	-	6,6,6	0.48	0	5,5,5	0.37	0
11	BMA	A	824	-	11,11,12	0.44	0	15,15,17	0.92	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
13	LHK	A	829	-	-	12/27/27/27	-
7	EDO	A	827	-	-	0/1/1/1	-
7	EDO	A	828	-	-	1/1/1/1	-
6	NAG	A	809	1	-	2/6/23/26	0/1/1/1
7	EDO	A	819	-	-	0/1/1/1	-
14	GOL	A	830	-	-	0/4/4/4	-
6	NAG	A	808	1	-	2/6/23/26	0/1/1/1
7	EDO	A	826	-	-	1/1/1/1	-
12	PEG	A	825	-	-	1/4/4/4	-
11	BMA	A	824	-	-	0/2/19/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	A	829	LHK	OAE-SAV	8.94	1.55	1.43
13	A	829	LHK	OAF-SAV	6.17	1.51	1.43
13	A	829	LHK	SAV-N	4.57	1.67	1.62
13	A	829	LHK	CAU-CAS	2.02	1.57	1.52
13	A	829	LHK	OE1-CD	-2.00	1.24	1.30

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
13	A	829	LHK	OAF-SAV-N	4.83	115.26	106.49
6	A	809	NAG	C1-O5-C5	4.02	117.58	112.19
13	A	829	LHK	OAF-SAV-OAE	-3.46	115.24	120.34
6	A	808	NAG	C1-O5-C5	3.24	116.53	112.19
13	A	829	LHK	CG-CB-CA	-2.79	108.02	113.16
13	A	829	LHK	OAE-SAV-NAO	2.78	111.54	106.49
11	A	824	BMA	C1-C2-C3	2.54	113.35	109.64
13	A	829	LHK	CAS-CAU-NAO	2.40	116.96	111.61
13	A	829	LHK	OAE-SAV-N	-2.25	102.39	106.49
13	A	829	LHK	OAB-CAQ-CAM	-2.24	115.95	122.84

There are no chirality outliers.

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	808	NAG	C8-C7-N2-C2
6	A	808	NAG	O7-C7-N2-C2
13	A	829	LHK	CAM-CAU-NAO-SAV
13	A	829	LHK	CAU-NAO-SAV-OAE
13	A	829	LHK	CAU-NAO-SAV-N
13	A	829	LHK	C-CA-N-SAV
13	A	829	LHK	CAQ-CAM-CAU-NAO
7	A	826	EDO	O1-C1-C2-O2
13	A	829	LHK	CA-N-SAV-OAF
13	A	829	LHK	CB-CA-N-SAV
13	A	829	LHK	CAQ-CAM-CAU-CAS
13	A	829	LHK	OXT-C-CA-N
12	A	825	PEG	C1-C2-O2-C3
6	A	809	NAG	O5-C5-C6-O6
6	A	809	NAG	C4-C5-C6-O6
13	A	829	LHK	O-C-CA-N
7	A	828	EDO	O1-C1-C2-O2
13	A	829	LHK	OAJ-CAS-CAU-CAM
13	A	829	LHK	OAD-CAS-CAU-CAM

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	809	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	696/707 (98%)	0.41	40 (5%)	29 40	9, 25, 45, 102	83 (11%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	541	TRP	7.1
1	A	546	PHE	5.9
1	A	544	ASN	5.8
1	A	548	GLY	5.7
1	A	543	THR	5.3
1	A	547	SER	4.1
1	A	154	VAL	4.1
1	A	719[A]	VAL	3.8
1	A	155	SER	3.8
1	A	656[A]	SER	3.8
1	A	335	GLY	3.4
1	A	655[A]	LYS	3.3
1	A	337	PHE	3.1
1	A	279[A]	TYR	3.1
1	A	310	GLY	3.1
1	A	334	THR	3.0
1	A	506	PHE	3.0
1	A	153	ASN	2.9
1	A	131[A]	ILE	2.8
1	A	733	TYR	2.6
1	A	138	ILE	2.6
1	A	55	LYS	2.6
1	A	652[A]	ASP	2.5
1	A	219	LEU	2.5
1	A	545	LYS	2.5
1	A	56	HIS	2.3
1	A	304	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	341	LYS	2.2
1	A	136[A]	ASN	2.2
1	A	651[A]	GLN	2.2
1	A	321	GLY	2.2
1	A	194	ILE	2.2
1	A	313	ALA	2.2
1	A	750	ALA	2.2
1	A	124	HIS	2.2
1	A	132	ASN	2.2
1	A	152	GLU	2.1
1	A	122	LYS	2.1
1	A	336	ASN	2.1
1	A	174[A]	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BMA	B	3	11/12	0.56	0.15	78,89,96,97	0
3	NAG	C	2	14/15	0.58	0.18	66,68,72,78	0
3	NAG	C	1	14/15	0.72	0.15	46,53,60,61	0
3	NAG	D	2	14/15	0.75	0.14	55,60,68,72	0
5	MAN	F	5	11/12	0.77	0.15	50,55,59,64	0
4	MAN	E	4	11/12	0.79	0.16	45,50,52,52	0
4	BMA	E	3	11/12	0.80	0.13	46,50,54,64	0
2	NAG	B	2	14/15	0.81	0.15	40,48,56,67	0
5	NAG	F	2	14/15	0.85	0.13	37,38,47,56	0
3	NAG	D	1	14/15	0.86	0.12	39,43,48,52	0
5	MAN	F	4	11/12	0.88	0.12	42,43,47,48	0
4	NAG	E	2	14/15	0.90	0.10	31,38,43,48	0
5	BMA	F	3	11/12	0.91	0.10	39,40,42,46	0
5	NAG	F	1	14/15	0.92	0.10	23,29,37,49	0
4	NAG	E	1	14/15	0.93	0.09	26,29,32,38	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	B	1	14/15	0.94	0.09	33,38,44,48	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NAG	A	808	14/15	0.54	0.19	68,77,84,87	0
11	BMA	A	824	11/12	0.62	0.23	117,124,128,128	0
7	EDO	A	819	4/4	0.68	0.37	95,95,96,96	0
14	GOL	A	830	6/6	0.81	0.17	35,40,41,48	0
7	EDO	A	826	4/4	0.82	0.26	79,80,82,83	0
12	PEG	A	825	7/7	0.82	0.20	52,53,57,58	0
7	EDO	A	827	4/4	0.82	0.23	69,71,72,76	0
7	EDO	A	828	4/4	0.83	0.13	77,78,78,78	0
6	NAG	A	809	14/15	0.89	0.11	30,44,52,53	0
13	LHK	A	829	22/22	0.96	0.09	22,25,35,38	0
10	CL	A	823	1/1	0.99	0.04	22,22,22,22	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	CA	A	822	1/1	1.00	0.02	18,18,18,18	0
8	ZN	A	820	1/1	1.00	0.02	20,20,20,20	0
8	ZN	A	821	1/1	1.00	0.02	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.