



wwPDB EM Validation Summary Report ⓘ

Mar 7, 2026 – 11:30 PM UTC

PDB ID : 6SKL / pdb_00006skl
EMDB ID : EMD-10227
Title : Cryo-EM structure of the CMG Fork Protection Complex at a replication fork
- Conformation 1
Authors : Yeeles, J.; Baretic, D.; Jenkyn-Bedford, M.
Deposited on : 2019-08-16
Resolution : 3.70 Å(reported)
Based on initial models : 5MQI, 4C8H, 5U8S

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMD archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11569 (3.20 - 4.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	2	868	27% 75% 23%
2	3	971	12% 62% 37%
3	4	933	46% 70% 27%
4	5	775	16% 77% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	6	1017	
6	7	845	
7	A	208	
8	B	213	
9	C	217	
10	D	294	
11	E	650	
12	F	927	
12	G	927	
12	H	927	
13	I	85	
14	J	61	
15	X	1238	
16	Y	317	

2 Entry composition [i](#)

There are 19 unique types of molecules in this entry. The entry contains 118220 atoms, of which 59036 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA replication licensing factor MCM2.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	2	669	Total	C	H	N	O	S	0	0
			10572	3310	5301	948	995	18		

- Molecule 2 is a protein called DNA replication licensing factor MCM3.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	3	616	Total	C	H	N	O	S	0	0
			9707	3041	4886	862	905	13		

- Molecule 3 is a protein called DNA replication licensing factor MCM4.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	4	679	Total	C	H	N	O	S	0	0
			10882	3389	5484	939	1039	31		

- Molecule 4 is a protein called Minichromosome maintenance protein 5.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	5	611	Total	C	H	N	O	S	0	0
			9676	3018	4874	826	934	24		

- Molecule 5 is a protein called DNA replication licensing factor MCM6.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	6	637	Total	C	H	N	O	S	0	0
			10044	3162	5031	881	945	25		

- Molecule 6 is a protein called DNA replication licensing factor MCM7.

Mol	Chain	Residues	Atoms						AltConf	Trace
6	7	635	Total	C	H	N	O	S	0	0
			9980	3142	5018	861	933	26		

- Molecule 7 is a protein called DNA replication complex GINS protein PSF1.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	A	197	Total	C	H	N	O	S	0	0
			3227	1012	1616	277	313	9		

- Molecule 8 is a protein called DNA replication complex GINS protein PSF2.

Mol	Chain	Residues	Atoms						AltConf	Trace
8	B	193	Total	C	H	N	O	S	0	0
			3292	1039	1675	286	287	5		

- Molecule 9 is a protein called DNA replication complex GINS protein PSF3.

Mol	Chain	Residues	Atoms						AltConf	Trace
9	C	177	Total	C	H	N	O	S	0	0
			2858	924	1436	230	260	8		

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-22	MET	-	initiating methionine	UNP Q12146
C	-21	GLY	-	expression tag	UNP Q12146
C	-20	SER	-	expression tag	UNP Q12146
C	-19	SER	-	expression tag	UNP Q12146
C	-18	HIS	-	expression tag	UNP Q12146
C	-17	HIS	-	expression tag	UNP Q12146
C	-16	HIS	-	expression tag	UNP Q12146
C	-15	HIS	-	expression tag	UNP Q12146
C	-14	HIS	-	expression tag	UNP Q12146
C	-13	HIS	-	expression tag	UNP Q12146
C	-12	SER	-	expression tag	UNP Q12146
C	-11	SER	-	expression tag	UNP Q12146
C	-10	GLY	-	expression tag	UNP Q12146
C	-9	LEU	-	expression tag	UNP Q12146
C	-8	VAL	-	expression tag	UNP Q12146
C	-7	PRO	-	expression tag	UNP Q12146
C	-6	ARG	-	expression tag	UNP Q12146
C	-5	GLY	-	expression tag	UNP Q12146
C	-4	SER	-	expression tag	UNP Q12146
C	-3	HIS	-	expression tag	UNP Q12146
C	-2	MET	-	expression tag	UNP Q12146
C	-1	ALA	-	expression tag	UNP Q12146
C	0	SER	-	expression tag	UNP Q12146

- Molecule 10 is a protein called DNA replication complex GINS protein SLD5.

Mol	Chain	Residues	Atoms						AltConf	Trace
10	D	243	Total	C	H	N	O	S	0	0
			4008	1276	2004	327	389	12		

- Molecule 11 is a protein called Cell division control protein 45.

Mol	Chain	Residues	Atoms						AltConf	Trace
11	E	564	Total	C	H	N	O	S	0	0
			9129	2916	4560	772	867	14		

- Molecule 12 is a protein called DNA polymerase alpha-binding protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
12	F	424	Total	C	H	N	O	S	0	0
			6759	2188	3355	564	637	15		
12	G	422	Total	C	H	N	O	S	0	0
			6693	2172	3313	557	636	15		
12	H	425	Total	C	H	N	O	S	0	0
			6769	2193	3358	565	638	15		

- Molecule 13 is a DNA chain called DNA fork, leading-strand template.

Mol	Chain	Residues	Atoms						AltConf	Trace
13	I	37	Total	C	H	N	O	P	0	0
			1192	370	430	116	239	37		

- Molecule 14 is a DNA chain called DNA fork, lagging-strand template.

Mol	Chain	Residues	Atoms						AltConf	Trace
14	J	22	Total	C	H	N	O	P	0	0
			686	211	248	74	131	22		

- Molecule 15 is a protein called Topoisomerase 1-associated factor 1.

Mol	Chain	Residues	Atoms						AltConf	Trace
15	X	665	Total	C	H	N	O	S	0	0
			10990	3505	5580	912	974	19		


- Molecule 16 is a protein called Chromosome segregation in meiosis protein 3.

Mol	Chain	Residues	Atoms						AltConf	Trace
16	Y	94	Total	C	H	N	O	S	0	0
			1616	507	828	144	133	4		

- Molecule 17 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
17	2	1	Total	Zn	0
			1	1	
17	4	1	Total	Zn	0
			1	1	
17	5	1	Total	Zn	0
			1	1	
17	6	1	Total	Zn	0
			1	1	
17	7	1	Total	Zn	0
			1	1	

- Molecule 18 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						AltConf
18	2	1	Total	C	H	N	O	P	0
			44	10	13	6	12	3	
18	3	1	Total	C	H	N	O	P	0
			44	10	13	6	12	3	
18	5	1	Total	C	H	N	O	P	0
			44	10	13	6	12	3	


- Molecule 19 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
19	2	1	Total 1	Mg 1	0
19	3	1	Total 1	Mg 1	0
19	5	1	Total 1	Mg 1	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA replication licensing factor MCM2



Chain 3:



Chain 4:








Chain 5:











- Molecule 8: DNA replication complex GINS protein PSF2





- Molecule 9: DNA replication complex GINS protein PSF3





- Molecule 10: DNA replication complex GINS protein SLD5





- Molecule 11: Cell division control protein 45




- Molecule 12: DNA polymerase alpha-binding protein









• Molecule 12: DNA polymerase alpha-binding protein






- Molecule 13: DNA fork, leading-strand template




- Molecule 14: DNA fork, lagging-strand template



- Molecule 15: Topoisomerase 1-associated factor 1





- Molecule 16: Chromosome segregation in meiosis protein 3



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	35000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	37	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	130000	Depositor
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.068	Depositor
Minimum map value	-0.033	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.018	Depositor
Map size ($\text{\AA}$)	377.64, 377.64, 377.64	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.049, 1.049, 1.049	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	2	0.81	0/5361	1.41	15/7244 (0.2%)
2	3	0.79	0/4907	1.36	10/6655 (0.2%)
3	4	0.77	0/5468	1.37	17/7376 (0.2%)
4	5	0.77	0/4869	1.35	11/6581 (0.2%)
5	6	0.78	0/5095	1.37	16/6875 (0.2%)
6	7	0.77	0/5037	1.34	19/6814 (0.3%)
7	A	0.76	0/1631	1.40	5/2194 (0.2%)
8	B	0.84	0/1650	1.38	7/2231 (0.3%)
9	C	0.77	0/1456	1.34	4/1966 (0.2%)
10	D	0.78	0/2040	1.31	3/2755 (0.1%)
11	E	0.80	1/4653 (0.0%)	1.35	15/6297 (0.2%)
12	F	0.80	0/3489	1.30	4/4724 (0.1%)
12	G	0.77	0/3465	1.34	6/4696 (0.1%)
12	H	0.77	0/3496	1.31	6/4735 (0.1%)
13	I	0.78	0/849	1.51	7/1312 (0.5%)
14	J	0.81	0/488	1.47	2/747 (0.3%)
15	X	0.73	0/5512	1.31	17/7426 (0.2%)
16	Y	0.76	0/804	1.25	0/1074
All	All	0.78	1/60270 (0.0%)	1.35	164/81702 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	2	0	3
2	3	0	3
3	4	0	1
4	5	0	3
5	6	0	3

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
7	A	0	1
8	B	0	3
11	E	0	6
12	F	0	3
13	I	0	8
14	J	0	5
15	X	0	5
16	Y	0	1
All	All	0	45

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	E	461	ASN	N-CA	-5.99	1.34	1.46

The worst 5 of 164 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	2	587	LYS	CB-CA-C	8.62	123.42	111.86
8	B	171	ASP	N-CA-C	7.87	120.55	111.11
4	5	41	ASP	CA-CB-CG	7.66	120.26	112.60
5	6	176	ARG	NE-CZ-NH2	7.01	125.51	119.20
9	C	169	LEU	N-CA-C	6.88	120.15	109.07

There are no chirality outliers.

5 of 45 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	2	247	ARG	Sidechain
1	2	327	ARG	Sidechain
1	2	336	TYR	Sidechain
2	3	247	TYR	Sidechain
2	3	400	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	2	5271	5301	5294	5	0
2	3	4821	4886	4882	7	0
3	4	5398	5484	5472	5	0
4	5	4802	4874	4865	3	0
5	6	5013	5031	5022	2	0
6	7	4962	5018	5006	6	0
7	A	1611	1616	1615	2	0
8	B	1617	1675	1674	1	0
9	C	1422	1436	1436	1	0
10	D	2004	2004	2001	2	0
11	E	4569	4560	4556	1	0
12	F	3404	3355	3352	0	0
12	G	3380	3313	3310	0	0
12	H	3411	3358	3355	2	0
13	I	762	430	427	0	0
14	J	438	248	248	1	0
15	X	5410	5580	5573	1	0
16	Y	788	828	827	0	0
17	2	1	0	0	0	0
17	4	1	0	0	0	0
17	5	1	0	0	0	0
17	6	1	0	0	0	0
17	7	1	0	0	0	0
18	2	31	13	13	3	0
18	3	31	13	13	3	0
18	5	31	13	13	2	0
19	2	1	0	0	0	0
19	3	1	0	0	0	0
19	5	1	0	0	0	0
All	All	59184	59036	58954	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

The worst 5 of 33 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:3:412:SER:H	18:3:1500:ANP:HNB1	1.15	0.92
1:2:546:GLY:H	18:2:902:ANP:HNB1	1.20	0.84
2:3:412:SER:N	18:3:1500:ANP:HNB1	1.90	0.67
6:7:206:PRO:HD3	6:7:352:THR:HG21	1.76	0.66
4:5:419:GLY:H	18:5:801:ANP:HNB1	1.42	0.66

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	2	665/868 (77%)	642 (96%)	23 (4%)	0	100	100
2	3	606/971 (62%)	591 (98%)	15 (2%)	0	100	100
3	4	663/933 (71%)	631 (95%)	32 (5%)	0	100	100
4	5	601/775 (78%)	580 (96%)	21 (4%)	0	100	100
5	6	625/1017 (62%)	601 (96%)	24 (4%)	0	100	100
6	7	619/845 (73%)	593 (96%)	26 (4%)	0	100	100
7	A	193/208 (93%)	189 (98%)	4 (2%)	0	100	100
8	B	189/213 (89%)	185 (98%)	4 (2%)	0	100	100
9	C	171/217 (79%)	170 (99%)	1 (1%)	0	100	100
10	D	237/294 (81%)	231 (98%)	6 (2%)	0	100	100
11	E	554/650 (85%)	544 (98%)	10 (2%)	0	100	100
12	F	418/927 (45%)	395 (94%)	23 (6%)	0	100	100
12	G	416/927 (45%)	406 (98%)	10 (2%)	0	100	100
12	H	419/927 (45%)	403 (96%)	16 (4%)	0	100	100
15	X	649/1238 (52%)	630 (97%)	19 (3%)	0	100	100
16	Y	92/317 (29%)	92 (100%)	0	0	100	100
All	All	7117/11327 (63%)	6883 (97%)	234 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	2	577/770 (75%)	577 (100%)	0	100	100
2	3	530/835 (64%)	529 (100%)	1 (0%)	87	86
3	4	614/848 (72%)	614 (100%)	0	100	100
4	5	548/688 (80%)	548 (100%)	0	100	100
5	6	550/886 (62%)	549 (100%)	1 (0%)	87	86
6	7	547/753 (73%)	546 (100%)	1 (0%)	87	86
7	A	182/193 (94%)	182 (100%)	0	100	100
8	B	183/198 (92%)	183 (100%)	0	100	100
9	C	159/192 (83%)	159 (100%)	0	100	100
10	D	234/279 (84%)	234 (100%)	0	100	100
11	E	507/586 (86%)	506 (100%)	1 (0%)	87	86
12	F	375/825 (46%)	375 (100%)	0	100	100
12	G	372/825 (45%)	371 (100%)	1 (0%)	86	83
12	H	375/825 (46%)	374 (100%)	1 (0%)	86	83
15	X	607/1125 (54%)	607 (100%)	0	100	100
16	Y	87/285 (30%)	87 (100%)	0	100	100
All	All	6447/10113 (64%)	6441 (100%)	6 (0%)	87	88

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
11	E	246	THR
12	G	671	ASP
12	H	837	LEU
5	6	690	ASN
2	3	274	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 74 such sidechains are listed below:

Mol	Chain	Res	Type
12	H	510	GLN
15	X	604	GLN
12	H	565	ASN
15	X	416	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	5	579	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 8 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
18	ANP	2	902	19	33,33,33	1.62	8 (24%)	45,52,52	1.48	9 (20%)
18	ANP	5	801	19	33,33,33	1.48	7 (21%)	45,52,52	1.50	6 (13%)
18	ANP	3	1500	19	33,33,33	1.69	9 (27%)	45,52,52	1.46	8 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
18	ANP	2	902	19	-	5/18/38/38	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
18	ANP	5	801	19	-	3/18/38/38	0/3/3/3
18	ANP	3	1500	19	-	3/18/38/38	0/3/3/3

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
18	3	1500	ANP	PB-O3A	-4.19	1.53	1.59
18	3	1500	ANP	PA-O3A	-4.05	1.55	1.59
18	5	801	ANP	PB-O1B	3.76	1.51	1.46
18	2	902	ANP	PB-O3A	-3.56	1.54	1.59
18	2	902	ANP	PG-O1G	3.51	1.51	1.46

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
18	3	1500	ANP	O3A-PB-N3B	-4.76	93.38	106.59
18	5	801	ANP	O2A-PA-O3A	3.95	117.94	107.27
18	2	902	ANP	O3A-PB-N3B	-3.44	97.04	106.59
18	5	801	ANP	O3A-PB-N3B	-3.39	97.19	106.59
18	5	801	ANP	C4-C5-N7	3.23	114.28	110.58

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
18	2	902	ANP	PB-N3B-PG-O1G
18	3	1500	ANP	PB-N3B-PG-O1G
18	5	801	ANP	PB-N3B-PG-O1G
18	5	801	ANP	PA-O3A-PB-O1B
18	5	801	ANP	PA-O3A-PB-O2B


There are no ring outliers.

3 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
18	2	902	ANP	3	0
18	5	801	ANP	2	0
18	3	1500	ANP	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand ANP 5 801





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-10227. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 180



Y Index: 148




Z Index: 166


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views

This section was not generated.


6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 222 nm³; this corresponds to an approximate mass of 201 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.270 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.270 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.70	-	-
Author-provided FSC curve	3.64	4.31	3.73
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-10227 and PDB model 6SKL. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.018).

9.4 Atom inclusion [i](#)



At the recommended contour level, 45% of all backbone atoms, 45% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.018) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.4500	 0.3520
2	 0.4770	 0.3730
3	 0.5650	 0.4100
4	 0.2950	 0.2520
5	 0.5550	 0.4200
6	 0.4760	 0.3570
7	 0.4340	 0.3250
A	 0.5090	 0.3620
B	 0.6430	 0.4560
C	 0.5790	 0.4180
D	 0.5830	 0.3980
E	 0.5380	 0.3970
F	 0.4960	 0.3800
G	 0.2430	 0.2560
H	 0.2690	 0.2680
I	 0.5510	 0.3190
J	 0.6370	 0.3460
X	 0.4200	 0.3340
Y	 0.4650	 0.3070

