



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:58 PM UTC

PDB ID : 6SLM / pdb_00006slm
Title : Crystal structure of full-length HPV31 E6 oncoprotein in complex with LXXLL peptide of ubiquitin ligase E6AP
Authors : Conrady, M.; Gogl, G.; Cousido-Siah, A.; Mitschler, A.; Trave, G.; Simon, C.
Deposited on : 2019-08-20
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	 4% 81% 16% ..
2	B	2	 100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4281 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose/maltodextrin-binding periplasmic protein,Protein E6,Ubiquitin-protein ligase E3A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	539	Total	C	N	O	S	0	2	0
			4213	2695	709	794	15			

There are 29 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	630	MET	ALA	conflict	UNP P0AEX9
A	712	ALA	ASP	conflict	UNP P0AEX9
A	713	ALA	LYS	conflict	UNP P0AEX9
A	869	ALA	LYS	conflict	UNP P0AEX9
A	989	ALA	GLU	conflict	UNP P0AEX9
A	992	ALA	LYS	conflict	UNP P0AEX9
A	993	ALA	ASP	conflict	UNP P0AEX9
A	997	ASN	ARG	conflict	UNP P0AEX9
A	998	ALA	ILE	conflict	UNP P0AEX9
A	999	ALA	THR	conflict	UNP P0AEX9
A	1000	ALA	LYS	conflict	UNP P0AEX9
A	1097	SER	CYS	conflict	UNP P17386
A	1111	SER	CYS	conflict	UNP P17386
A	1150	GLY	-	linker	UNP P17386
A	1151	SER	-	linker	UNP P17386
A	1152	SER	-	linker	UNP P17386
A	1153	GLY	-	linker	UNP P17386
A	1154	SER	-	linker	UNP P17386
A	1155	GLY	-	linker	UNP P17386
A	1156	SER	-	linker	UNP P17386
A	1157	GLY	-	linker	UNP P17386
A	1158	SER	-	linker	UNP P17386
A	1159	GLY	-	linker	UNP P17386
A	1160	SER	-	linker	UNP P17386
A	1161	GLY	-	linker	UNP P17386
A	1162	SER	-	linker	UNP P17386

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1163	ALA	-	linker	UNP P17386
A	1164	ALA	-	linker	UNP P17386
A	1165	ALA	-	linker	UNP P17386

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	31	Total	O	0	0
			31	31		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose/maltodextrin-binding periplasmic protein, Protein E6, Ubiquitin-protein ligase E3A



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	113.64Å 113.64Å 185.97Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.21 – 2.80 49.21 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.21-2.80) 92.6 (49.21-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.230 , 0.268 0.238 , 0.275	Depositor DCC
R_{free} test set	894 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	56.6	Xtriage
Anisotropy	0.274	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 67.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4281	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.15	0/4312	0.31	0/5850

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4213	0	4141	57	0
2	B	23	0	21	2	0
3	A	2	0	0	0	0
4	A	12	0	16	2	0
5	A	31	0	0	3	0
All	All	4281	0	4178	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1144:ARG:HG2	1:A:1144:ARG:HH21	1.43	0.84
1:A:637:LEU:HB2	1:A:665:VAL:HG12	1.62	0.79
2:B:1:GLC:O3	2:B:2:GLC:O2	1.60	0.78
1:A:775:SER:HB2	1:A:852:THR:HG22	1.74	0.68
1:A:752:LEU:HD21	1:A:756:PRO:HD3	1.77	0.67
1:A:860:TRP:HB3	5:A:1303:HOH:O	1.95	0.67
1:A:689:ILE:HD11	1:A:906:ALA:HB1	1.77	0.65
1:A:711:PRO:HG2	1:A:716:GLN:OE1	1.98	0.64
1:A:814:ASP:O	1:A:819:LYS:NZ	2.32	0.63
1:A:902:ASN:HB3	1:A:905:LEU:HB3	1.80	0.63
1:A:749:LYS:NZ	5:A:1305:HOH:O	2.31	0.63
1:A:1005:PRO:O	1:A:1010:ARG:NH2	2.32	0.63
1:A:748:ASN:HD22	1:A:870:VAL:HG13	1.65	0.61
1:A:916:THR:HG23	1:A:919:GLY:H	1.66	0.60
1:A:1072:LYS:NZ	1:A:1148:GLN:CG	2.65	0.59
1:A:1144:ARG:HH21	1:A:1144:ARG:CG	2.11	0.59
1:A:756:PRO:HG2	1:A:854:MET:HE1	1.84	0.59
1:A:778:MET:HB2	1:A:852:THR:HG21	1.85	0.59
1:A:900:SER:O	1:A:903:LYS:NZ	2.36	0.58
1:A:1008:ARG:HG2	1:A:1054:TYR:CZ	2.41	0.56
1:A:774:LYS:HD2	1:A:851:GLU:HA	1.87	0.56
1:A:1102:ARG:HD3	1:A:1107:GLN:HG3	1.90	0.54
1:A:752:LEU:HD11	1:A:765:LEU:HD11	1.90	0.53
1:A:1011:LYS:HE2	1:A:1051:THR:HG22	1.92	0.52
1:A:1072:LYS:HZ1	1:A:1148:GLN:CG	2.21	0.52
1:A:779:PHE:HB2	5:A:1306:HOH:O	2.10	0.52
1:A:1144:ARG:NH1	4:A:1204:GOL:O2	2.43	0.51
1:A:642:ASN:ND2	1:A:644:ASP:OD1	2.45	0.50
1:A:739:ALA:HA	1:A:932:VAL:HA	1.94	0.50
1:A:797:TYR:HB2	4:A:1203:GOL:H11	1.94	0.50
1:A:794:ASP:HB3	1:A:817:GLY:HA2	1.95	0.48
1:A:944:ASP:HB3	1:A:947:ILE:HB	1.94	0.48
1:A:698:GLY:HA3	1:A:962:ASN:O	2.13	0.48
1:A:884:PRO:HB3	1:A:956:LYS:HD2	1.96	0.47
1:A:673:LEU:H	1:A:673:LEU:HD23	1.80	0.47
1:A:689:ILE:HD12	1:A:910:LEU:HD11	1.96	0.47
1:A:695:ASP:OD2	2:B:2:GLC:O3	2.27	0.45
1:A:859:PRO:HA	1:A:862:TRP:CE2	2.51	0.45
1:A:1001:MET:HE1	1:A:1018:ALA:HB1	1.98	0.45
1:A:1021:ILE:HG13	1:A:1022:PRO:HD2	1.99	0.45
1:A:709:ILE:HG13	1:A:711:PRO:HD3	2.00	0.44
1:A:727:VAL:HG23	1:A:733:LEU:HD12	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:634:GLU:HA	1:A:902:ASN:HD21	1.83	0.43
1:A:1144:ARG:HG2	1:A:1144:ARG:NH2	2.21	0.43
1:A:921:GLU:HG2	1:A:925:LYS:HE3	2.00	0.42
1:A:959:ILE:H	1:A:959:ILE:HD12	1.85	0.42
1:A:1144:ARG:CG	1:A:1144:ARG:NH2	2.73	0.42
1:A:663:ILE:HG12	1:A:905:LEU:HD21	2.01	0.42
1:A:1114:GLU:HG2	1:A:1137:ILE:HG13	2.01	0.42
1:A:789:PRO:HG3	1:A:887:PRO:HB3	2.02	0.42
1:A:862:TRP:HB2	1:A:928:PRO:HG2	2.01	0.42
1:A:976:ALA:HB2	1:A:994:ALA:HB2	2.00	0.42
1:A:979:ASN:HB3	1:A:985:GLN:HG2	2.00	0.42
1:A:736:TYR:HB2	1:A:894:ALA:HB3	2.01	0.42
1:A:1142:ARG:O	1:A:1142:ARG:HG3	2.20	0.42
1:A:1012:LEU:HD11	1:A:1042:VAL:HG13	2.03	0.41
1:A:1144:ARG:HG2	1:A:1144:ARG:H	1.46	0.40
1:A:719:LEU:HB2	1:A:724:TRP:HE1	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	536/551 (97%)	515 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	435/448 (97%)	424 (98%)	11 (2%)	42 76

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	645	LYS
1	A	716	GLN
1	A	757	LYS
1	A	888	PHE
1	A	903	LYS
1	A	1021	ILE
1	A	1083	VAL
1	A	1105	THR
1	A	1144	ARG
1	A	1145	THR
1	A	1147	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	642	ASN
1	A	679	GLN
1	A	782	GLN
1	A	871	ASN
1	A	883	GLN
1	A	995	GLN
1	A	1004	ASN
1	A	1013	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLC	B	1	2	12,12,12	1.48	3 (25%)	17,17,17	2.04	3 (17%)
2	GLC	B	2	2	11,11,12	1.60	3 (27%)	15,15,17	1.57	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	2/2/22/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	GLC	O5-C5	-2.83	1.37	1.43
2	B	1	GLC	O5-C5	-2.59	1.38	1.44
2	B	2	GLC	O5-C1	-2.58	1.39	1.43
2	B	2	GLC	C1-C2	-2.04	1.47	1.52
2	B	1	GLC	C3-C2	-2.01	1.47	1.52
2	B	1	GLC	C4-C5	-2.01	1.48	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	C4-C3-C2	-6.94	98.64	110.83
2	B	2	GLC	O3-C3-C2	-3.73	102.44	110.05
2	B	1	GLC	O1-C1-O5	-2.30	103.58	110.41
2	B	2	GLC	O2-C2-C1	-2.08	104.47	109.22
2	B	2	GLC	C6-C5-C4	2.02	117.98	113.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	C6-C5-C4	-2.02	108.06	113.02

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	GLC	O5-C5-C6-O6
2	B	1	GLC	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	GLC	2	0
2	B	1	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	1204	-	5,5,5	0.09	0	5,5,5	0.49	0
4	GOL	A	1203	-	5,5,5	0.96	0	5,5,5	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1204	-	-	4/4/4/4	-
4	GOL	A	1203	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1204	GOL	C1-C2-C3-O3
4	A	1203	GOL	C1-C2-C3-O3
4	A	1204	GOL	O1-C1-C2-C3
4	A	1203	GOL	O2-C2-C3-O3
4	A	1204	GOL	O2-C2-C3-O3
4	A	1204	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1204	GOL	1	0
4	A	1203	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	539/551 (97%)	0.47	23 (4%)	40 31	29, 83, 138, 185	1 (0%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1151	SER	4.5
1	A	751	LEU	3.8
1	A	1143	PRO	3.5
1	A	1149	VAL	3.4
1	A	941	LEU	3.2
1	A	1178	GLU	2.9
1	A	713	ALA	2.9
1	A	869	ALA	2.9
1	A	1142	ARG	2.9
1	A	1145	THR	2.9
1	A	1148	GLN	2.8
1	A	634	GLU	2.4
1	A	1165	ALA	2.3
1	A	870	VAL	2.3
1	A	774	LYS	2.3
1	A	910	LEU	2.3
1	A	1152	SER	2.2
1	A	872	TYR	2.1
1	A	712	ALA	2.1
1	A	1076[A]	PHE	2.0
1	A	719	LEU	2.0
1	A	646	GLY	2.0
1	A	715	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	B	1	12/12	0.92	0.12	61,66,70,76	0
2	GLC	B	2	11/12	0.93	0.08	42,53,61,63	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	1203	6/6	0.58	0.23	92,94,98,99	0
4	GOL	A	1204	6/6	0.86	0.18	55,69,71,76	0
3	ZN	A	1201	1/1	0.99	0.02	42,42,42,42	0
3	ZN	A	1202	1/1	0.99	0.03	53,53,53,53	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around ZN A 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.