



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:09 AM UTC

PDB ID : 8SLG / pdb_00008slg
Title : Crystal Structure of Glycine tRNA ligase from Mycobacterium thermoresistibile (glycyl adenylate bound)
Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2023-04-21
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	12% 83% 6% 11%
1	B	482	9% 87% 5% 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7437 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycine-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	0	0
			3472	2194	612	653	13			
1	B	447	Total	C	N	O	S	0	0	0
			3600	2271	631	684	14			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP G7CIG9
A	-19	ALA	-	expression tag	UNP G7CIG9
A	-18	HIS	-	expression tag	UNP G7CIG9
A	-17	HIS	-	expression tag	UNP G7CIG9
A	-16	HIS	-	expression tag	UNP G7CIG9
A	-15	HIS	-	expression tag	UNP G7CIG9
A	-14	HIS	-	expression tag	UNP G7CIG9
A	-13	HIS	-	expression tag	UNP G7CIG9
A	-12	MET	-	expression tag	UNP G7CIG9
A	-11	GLY	-	expression tag	UNP G7CIG9
A	-10	THR	-	expression tag	UNP G7CIG9
A	-9	LEU	-	expression tag	UNP G7CIG9
A	-8	GLU	-	expression tag	UNP G7CIG9
A	-7	ALA	-	expression tag	UNP G7CIG9
A	-6	GLN	-	expression tag	UNP G7CIG9
A	-5	THR	-	expression tag	UNP G7CIG9
A	-4	GLN	-	expression tag	UNP G7CIG9
A	-3	GLY	-	expression tag	UNP G7CIG9
A	-2	PRO	-	expression tag	UNP G7CIG9
A	-1	GLY	-	expression tag	UNP G7CIG9
A	0	SER	-	expression tag	UNP G7CIG9
B	-20	MET	-	initiating methionine	UNP G7CIG9
B	-19	ALA	-	expression tag	UNP G7CIG9
B	-18	HIS	-	expression tag	UNP G7CIG9
B	-17	HIS	-	expression tag	UNP G7CIG9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP G7CIG9
B	-15	HIS	-	expression tag	UNP G7CIG9
B	-14	HIS	-	expression tag	UNP G7CIG9
B	-13	HIS	-	expression tag	UNP G7CIG9
B	-12	MET	-	expression tag	UNP G7CIG9
B	-11	GLY	-	expression tag	UNP G7CIG9
B	-10	THR	-	expression tag	UNP G7CIG9
B	-9	LEU	-	expression tag	UNP G7CIG9
B	-8	GLU	-	expression tag	UNP G7CIG9
B	-7	ALA	-	expression tag	UNP G7CIG9
B	-6	GLN	-	expression tag	UNP G7CIG9
B	-5	THR	-	expression tag	UNP G7CIG9
B	-4	GLN	-	expression tag	UNP G7CIG9
B	-3	GLY	-	expression tag	UNP G7CIG9
B	-2	PRO	-	expression tag	UNP G7CIG9
B	-1	GLY	-	expression tag	UNP G7CIG9
B	0	SER	-	expression tag	UNP G7CIG9

- Molecule 2 is 5'-O-(glycylsulfamoyl)adenosine (CCD ID: G5A) (formula: C₁₂H₁₇N₇O₇S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			27	12	7	7	1		
2	B	1	Total	C	N	O	S	0	0
			27	12	7	7	1		


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	159	Total 159	O 159	0	0
6	B	130	Total 130	O 130	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	170.25Å 87.11Å 99.36Å 90.00° 104.21° 90.00°	Depositor
Resolution (Å)	48.16 – 1.95 48.16 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.16-1.95) 99.9 (48.16-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.95Å)	Xtriage
Refinement program	PHENIX (1.21rc1_4932: ???)	Depositor
R, R_{free}	0.174 , 0.201 0.180 , 0.205	Depositor DCC
R_{free} test set	5109 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	38.1	Xtriage
Anisotropy	0.561	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7437	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, G5A, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/3561	0.52	0/4833
1	B	0.28	0/3691	0.51	0/5014
All	All	0.29	0/7252	0.52	0/9847

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3472	0	3301	16	0
1	B	3600	0	3434	16	0
2	A	27	0	17	0	0
2	B	27	0	17	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	B	18	0	24	2	0
6	A	159	0	0	1	0
6	B	130	0	0	0	0
All	All	7437	0	6793	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:GLU:OE2	1:B:384:ARG:NH1	2.32	0.62
1:A:84:HIS:ND1	1:A:139:GLU:OE1	2.33	0.60
1:B:43:GLU:HG2	5:B:501:GOL:H12	1.85	0.58
1:A:98:ARG:HG2	1:A:198:ILE:HB	1.90	0.53
1:A:78:GLY:O	1:A:82:VAL:HG22	2.10	0.52
1:B:142:MET:HE1	1:B:197:GLU:HA	1.93	0.51
1:A:232:ILE:HA	1:A:286:ILE:HD13	1.93	0.50
1:A:135:THR:HG22	1:A:136:GLU:N	2.26	0.50
1:A:14:ARG:NH2	1:A:413:GLU:OE2	2.32	0.50
1:B:11:LEU:HD11	1:B:413:GLU:HG2	1.94	0.49
1:B:50:TRP:CD2	5:B:502:GOL:H32	2.47	0.49
1:B:74:TRP:CH2	1:B:162:PRO:HD2	2.48	0.49
1:B:123:VAL:HG22	1:B:124:VAL:H	1.78	0.48
1:B:5:ILE:HG21	1:B:202:ASN:OD1	2.13	0.48
1:A:397:GLU:OE1	6:A:601:HOH:O	2.20	0.48
1:A:308:TYR:CE2	1:A:310:GLN:HG2	2.50	0.47
1:A:130:THR:HG22	1:A:130:THR:O	2.14	0.46
1:B:354:ARG:CZ	1:B:356:VAL:HG21	2.44	0.46
1:A:184:PRO:HA	1:A:218:PHE:O	2.16	0.46
1:B:197:GLU:HG2	1:B:206:ARG:HG2	1.98	0.46
1:A:282:GLU:O	1:A:329:ARG:HD3	2.16	0.45
1:B:346:ASN:HB3	1:B:352:ASP:OD2	2.17	0.45
1:B:184:PRO:HA	1:B:218:PHE:O	2.17	0.45
1:A:82:VAL:HG21	1:A:261:HIS:CG	2.52	0.45
1:B:124:VAL:HA	1:B:131:LYS:HA	1.99	0.44
1:B:82:VAL:HG21	1:B:261:HIS:CG	2.52	0.44
1:A:391:ARG:NH2	1:B:58:GLU:OE1	2.51	0.43
1:A:443:GLU:HG2	1:A:445:ILE:HD11	2.00	0.43
1:A:69:LEU:HB3	1:A:70:PRO:HD2	2.02	0.41
1:B:100:ASP:OD1	1:B:100:ASP:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	419/482 (87%)	412 (98%)	6 (1%)	1 (0%)	43	36
1	B	443/482 (92%)	437 (99%)	4 (1%)	2 (0%)	24	16
All	All	862/964 (89%)	849 (98%)	10 (1%)	3 (0%)	36	28

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	290	THR
1	A	290	THR
1	B	131	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/412 (89%)	366 (100%)	1 (0%)	86	87
1	B	382/412 (93%)	379 (99%)	3 (1%)	73	74
All	All	749/824 (91%)	745 (100%)	4 (0%)	81	82

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	409	ARG
1	B	82	VAL
1	B	87	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	197	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	171	ASN
1	A	174	ASN
1	A	392	GLN
1	B	97	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	G5A	B	504	-	28,29,29	0.75	1 (3%)	38,43,43	0.59	1 (2%)
5	GOL	B	502	-	5,5,5	0.24	0	5,5,5	0.33	0
2	G5A	A	501	-	28,29,29	0.77	1 (3%)	38,43,43	0.62	1 (2%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	501	-	5,5,5	0.11	0	5,5,5	0.73	0
5	GOL	B	503	-	5,5,5	0.32	0	5,5,5	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G5A	B	504	-	-	0/16/33/33	0/3/3/3
5	GOL	B	502	-	-	4/4/4/4	-
2	G5A	A	501	-	-	0/16/33/33	0/3/3/3
5	GOL	B	501	-	-	0/4/4/4	-
5	GOL	B	503	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	G5A	C-N3S	-2.50	1.32	1.39
2	B	504	G5A	C-N3S	-2.43	1.33	1.39

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	G5A	O5'-S-O1S	2.52	113.09	105.48
2	B	504	G5A	O5'-S-O1S	2.45	112.91	105.48

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	502	GOL	O1-C1-C2-C3
5	B	502	GOL	C1-C2-C3-O3
5	B	502	GOL	O2-C2-C3-O3
5	B	502	GOL	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	502	GOL	1	0
5	B	501	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/482 (89%)	0.72	59 (13%) 6 7	31, 44, 107, 168	0
1	B	447/482 (92%)	0.69	42 (9%) 14 16	31, 49, 94, 142	0
All	All	876/964 (90%)	0.71	101 (11%) 9 11	31, 47, 99, 168	0

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	124	VAL	7.7
1	A	134	TRP	7.5
1	A	-15	HIS	7.5
1	A	91	LEU	7.3
1	A	130	THR	6.7
1	B	118	VAL	6.5
1	B	151	ILE	6.4
1	A	-19	ALA	6.2
1	B	119	PRO	5.9
1	A	2	ALA	5.5
1	A	97	HIS	5.1
1	A	135	THR	5.1
1	B	347	ALA	4.8
1	A	198	ILE	4.3
1	B	120	MET	4.2
1	A	353	LYS	4.2
1	A	-16	HIS	4.1
1	A	128	CYS	4.0
1	B	4	ILE	4.0
1	B	348	LYS	4.0
1	A	376	HIS	3.8
1	A	98	ARG	3.8
1	A	99	GLN	3.7
1	B	124	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	134	TRP	3.7
1	B	461	CYS	3.6
1	A	137	PRO	3.6
1	A	94	HIS	3.6
1	B	123	VAL	3.6
1	A	87	LEU	3.6
1	B	2	ALA	3.6
1	A	151	ILE	3.6
1	A	-18	HIS	3.6
1	B	101	HIS	3.5
1	A	375	ARG	3.5
1	A	92	ASN	3.5
1	B	102	LEU	3.5
1	B	92	ASN	3.4
1	A	95	ARG	3.4
1	A	88	VAL	3.3
1	A	343	GLU	3.3
1	B	126	PRO	3.3
1	B	104	GLU	3.3
1	A	129	GLY	3.3
1	A	199	THR	3.1
1	B	133	ARG	3.1
1	A	378	ASP	3.0
1	A	82	VAL	2.9
1	B	130	THR	2.9
1	A	277	GLY	2.9
1	A	138	ARG	2.9
1	A	126	PRO	2.8
1	A	152	GLU	2.8
1	A	140	PHE	2.8
1	A	90	CYS	2.8
1	A	461	CYS	2.8
1	B	381	PRO	2.8
1	A	125	CYS	2.8
1	A	409	ARG	2.7
1	B	82	VAL	2.7
1	A	93	CYS	2.7
1	B	199	THR	2.7
1	A	153	SER	2.7
1	A	136	GLU	2.7
1	B	352	ASP	2.7
1	B	380	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	91	LEU	2.6
1	A	404	ILE	2.6
1	A	155	GLU	2.6
1	B	131	LYS	2.6
1	B	153	SER	2.6
1	A	86	PRO	2.5
1	B	353	LYS	2.5
1	B	376	HIS	2.5
1	B	150	PRO	2.5
1	B	433	THR	2.5
1	A	-17	HIS	2.4
1	A	200	PRO	2.4
1	A	150	PRO	2.3
1	B	122	GLU	2.3
1	A	5	ILE	2.2
1	A	342	ASP	2.2
1	A	311	ALA	2.2
1	B	430	ASN	2.2
1	A	89	GLU	2.2
1	A	4	ILE	2.1
1	B	3	SER	2.1
1	B	345	PRO	2.1
1	B	72	GLU	2.1
1	A	377	ALA	2.1
1	A	310	GLN	2.1
1	A	147	TYR	2.1
1	A	447	ILE	2.1
1	B	342	ASP	2.1
1	A	81	ASP	2.1
1	B	121	ASP	2.1
1	B	142	MET	2.1
1	B	75	VAL	2.1
1	A	58	GLU	2.0
1	B	129	GLY	2.0
1	B	406	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	B	503	6/6	0.62	0.23	58,73,79,79	0
5	GOL	B	502	6/6	0.84	0.18	47,51,55,56	0
5	GOL	B	501	6/6	0.84	0.17	47,51,57,57	0
3	ZN	A	502	1/1	0.94	0.09	150,150,150,150	0
2	G5A	A	501	27/27	0.95	0.08	33,37,39,40	0
4	MG	B	506	1/1	0.96	0.09	37,37,37,37	0
2	G5A	B	504	27/27	0.96	0.08	33,41,43,46	0
3	ZN	B	505	1/1	0.98	0.08	77,77,77,77	0
4	MG	A	503	1/1	0.99	0.10	38,38,38,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around G5A A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around G5A B 504:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.