



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:36 AM UTC

PDB ID : 1SM3 / pdb_00001sm3
Title : CRYSTAL STRUCTURE OF THE TUMOR SPECIFIC ANTIBODY SM3
COMPLEX WITH ITS PEPTIDE EPITOPE
Authors : Dokurno, P.; Bates, P.A.; Band, H.A.; Stewart, L.M.D.; Lally, J.M.; Burchell,
J.M.; Taylor-Papadimitriou, J.; Sternberg, M.J.E.; Snary, D.; Freemont, P.S.
Deposited on : 1997-12-23
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	215	
2	H	218	
3	P	13	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3564 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SM3 ANTIBODY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	210	Total	C	N	O	S	0	0	0
			1581	991	266	318	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	108	SER	GLN	conflict	UNP P01723
L	109	GLU	PRO	conflict	UNP P01723

- Molecule 2 is a protein called SM3 ANTIBODY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	211	Total	C	N	O	S	0	0	0
			1580	1006	266	300	8			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	3	GLN	LYS	conflict	UNP P01801
H	5	GLN	GLU	conflict	UNP P01801
H	20	LEU	VAL	conflict	UNP P01801
H	28	THR	ALA	conflict	UNP P01801
H	31	ASN	TYR	conflict	UNP P01801
H	43	LYS	ARG	conflict	UNP P01801
H	48	VAL	ILE	conflict	UNP P01801
H	52A	LEU	PHE	conflict	UNP P01801
H	56	ALA	-	insertion	UNP P01801
H	94	GLY	ARG	conflict	UNP P01801
H	95	VAL	GLU	conflict	UNP P01801
H	?	-	ILE	deletion	UNP P01801
H	?	-	TYR	deletion	UNP P01801
H	?	-	TYR	deletion	UNP P01801

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	97	GLN	PRO	conflict	UNP P01801
H	108	THR	LEU	conflict	UNP P01801
H	113	SER	ALA	conflict	UNP P01801
H	120	THR	SER	conflict	UNP P01801
H	129	ASN	ALA	conflict	UNP P01801
H	131	ALA	GLN	conflict	UNP P01801
H	132	SER	THR	conflict	UNP P01801
H	133	GLN	ASN	conflict	UNP P01801
H	160	ALA	SER	conflict	UNP P01801
H	208	ALA	LYS	conflict	UNP P01801

- Molecule 3 is a protein called PEPTIDE EPITOPE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	P	9	Total	C	N	O	0	0	0
			62	38	12	12			

- Molecule 4 is CADMIUM ION (CCD ID: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	4	Total	Cd	0	0
			4	4		
4	H	4	Total	Cd	0	0
			4	4		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	2	Total	Cl	0	0
			2	2		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	156	Total	O	0	0
			156	156		
6	H	162	Total	O	0	0
			162	162		
6	P	13	Total	O	0	0
			13	13		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: SM3 ANTIBODY



• Molecule 2: SM3 ANTIBODY



• Molecule 3: PEPTIDE EPITOPE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.20Å 83.73Å 67.24Å 90.00° 93.30° 90.00°	Depositor
Resolution (Å)	8.00 – 1.95	Depositor
% Data completeness (in resolution range)	85.6 (8.00-1.95)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.213 , 0.283	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3564	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	1.19	2/1618 (0.1%)	1.03	0/2212
2	H	1.20	5/1622 (0.3%)	1.10	3/2219 (0.1%)
3	P	1.21	0/64	1.18	0/89
All	All	1.20	7/3304 (0.2%)	1.07	3/4520 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1
2	H	0	1
All	All	0	2

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	163	GLN	CA-C	-5.48	1.46	1.52
1	L	74	THR	CA-C	-5.40	1.46	1.52
2	H	72	ASP	N-CA	-5.30	1.40	1.46
2	H	74	SER	CA-CB	5.22	1.62	1.53
2	H	74	SER	N-CA	5.20	1.52	1.46

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	181	VAL	CB-CA-C	-7.72	98.33	110.69
2	H	147	PRO	CA-N-CD	-5.70	103.52	111.50
2	H	149	PRO	CA-N-CD	-5.22	104.20	111.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	79	TYR	Sidechain
1	L	86	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1581	0	1524	56	0
2	H	1580	0	1529	33	0
3	P	62	0	56	2	0
4	H	4	0	0	0	0
4	L	4	0	0	0	0
5	H	2	0	0	0	0
6	H	162	0	0	1	0
6	L	156	0	0	7	0
6	P	13	0	0	0	0
All	All	3564	0	3109	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 88 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:3:ILE:HD13	1:L:27:THR:HG23	1.56	0.86
2:H:20:LEU:HG	2:H:82:MET:HE2	1.69	0.74
1:L:159:MET:H	1:L:159:MET:HE2	1.50	0.74
1:L:3:ILE:HD12	1:L:26:SER:OG	1.87	0.74
2:H:63:VAL:HG13	2:H:67:PHE:HB2	1.71	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	208/215 (97%)	197 (95%)	10 (5%)	1 (0%)	24	16
2	H	207/218 (95%)	200 (97%)	5 (2%)	2 (1%)	12	5
3	P	7/13 (54%)	6 (86%)	1 (14%)	0	100	100
All	All	422/446 (95%)	403 (96%)	16 (4%)	3 (1%)	18	10

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	41	ASP
2	H	172	SER
2	H	95	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	176/183 (96%)	169 (96%)	7 (4%)	28	17
2	H	172/185 (93%)	157 (91%)	15 (9%)	9	2
3	P	6/10 (60%)	6 (100%)	0	100	100
All	All	354/378 (94%)	332 (94%)	22 (6%)	16	6

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	52(C)	SER
2	H	93	THR
2	H	74	SER
2	H	140	CYS
1	L	208	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	128	ASN
1	L	194	GLN
1	L	200	HIS
2	H	81	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.