



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:36 PM UTC

PDB ID : 3SMI / pdb_00003smi
Title : Human poly(ADP-ribose) polymerase 14 (Parp14/Artd8) - catalytic domain in complex with a quinazoline inhibitor
Authors : Karlberg, T.; Moche, M.; Arrowsmith, C.H.; Berglund, H.; Bountra, C.; Edwards, A.M.; Ekblad, T.; Graslund, S.; Kouznetsova, E.; Nordlund, P.; Nyman, T.; Thorsell, A.G.; Tresaugues, L.; Weigelt, J.; Schuler, H.; Structural Genomics Consortium (SGC)
Deposited on : 2011-06-28
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	193	13% 78% 12% 8%
1	B	193	4% 79% 9% 10%

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2940 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Poly [ADP-ribose] polymerase 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	177	Total	C	N	O	S	0	0	0
			1428	906	248	270	4			
1	B	174	Total	C	N	O	S	0	0	0
			1411	892	246	269	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1528	SER	-	expression tag	UNP Q460N5
A	1529	MET	-	expression tag	UNP Q460N5
B	1528	SER	-	expression tag	UNP Q460N5
B	1529	MET	-	expression tag	UNP Q460N5

- Molecule 2 is 2-[[[(3-amino-1H-1,2,4-triazol-5-yl)sulfanyl]methyl]-8-methylquinazolin-4(3H)-one (CCD ID: QDR) (formula: C₁₂H₁₂N₆OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			20	12	6	1	1		
2	B	1	Total	C	N	O	S	0	0
			20	12	6	1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	25	Total	O	0	0
			25	25		
3	B	36	Total	O	0	0
			36	36		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly [ADP-ribose] polymerase 14



- Molecule 1: Poly [ADP-ribose] polymerase 14



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	40.56Å 66.42Å 145.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.67 – 2.40 33.67 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.67-2.40) 99.8 (33.67-2.40)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.23 (at 2.39Å)	Xtriage
Refinement program	BUSTER 2.9.2	Depositor
R, R_{free}	0.209 , 0.252 0.232 , 0.268	Depositor DCC
R_{free} test set	1601 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.991	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2940	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QDR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.10	2/1468 (0.1%)	1.27	9/1995 (0.5%)
1	B	1.39	9/1450 (0.6%)	1.20	4/1972 (0.2%)
All	All	1.25	11/2918 (0.4%)	1.23	13/3967 (0.3%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1568	GLU	C-O	-6.25	1.17	1.23
1	A	1704	ALA	CA-CB	-6.20	1.43	1.53
1	B	1572	ASN	C-O	-5.91	1.19	1.24
1	B	1551	VAL	C-O	-5.72	1.17	1.24
1	B	1567	ILE	C-O	-5.62	1.18	1.24
1	B	1666	ILE	C-O	-5.47	1.18	1.24
1	B	1539	VAL	C-O	-5.41	1.18	1.24
1	B	1570	ILE	C-O	-5.35	1.18	1.24
1	B	1595	ASN	C-O	-5.29	1.17	1.24
1	B	1596	GLU	C-O	-5.26	1.17	1.23
1	A	1570	ILE	CA-CB	-5.05	1.48	1.54

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1656	HIS	N-CA-C	7.02	119.85	108.41
1	A	1671	ASN	CA-C-N	6.86	129.78	120.38
1	A	1671	ASN	C-N-CA	6.86	129.78	120.38
1	B	1630	LYS	N-CA-C	6.71	119.17	111.11
1	A	1675	ILE	N-CA-C	-6.45	106.51	112.96
1	B	1701	LEU	N-CA-C	6.32	118.37	108.96
1	A	1695	ASN	CA-CB-CG	5.83	118.44	112.60
1	A	1544	SER	N-CA-C	-5.77	106.09	113.01

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1700	SER	N-CA-C	-5.69	106.98	114.04
1	A	1616	PHE	CA-C-N	5.50	130.73	120.95
1	A	1616	PHE	C-N-CA	5.50	130.73	120.95
1	B	1639	ASN	CA-CB-CG	5.41	118.01	112.60
1	A	1650	ASP	CA-CB-CG	5.05	117.65	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1428	0	1332	14	0
1	B	1411	0	1316	4	0
2	A	20	0	12	0	0
2	B	20	0	12	0	0
3	A	25	0	0	0	0
3	B	36	0	0	0	0
All	All	2940	0	2672	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1668:THR:O	1:B:1669:HIS:HB2	1.79	0.83
1:A:1534:GLN:CD	1:A:1614:ASN:OD1	2.43	0.62
1:A:1542:LEU:O	1:A:1545:ASP:HB2	2.00	0.62
1:A:1543:PRO:C	1:A:1545:ASP:H	2.09	0.58
1:A:1609:PRO:O	1:A:1613:ARG:HG2	2.07	0.55
1:A:1570:ILE:HD12	1:A:1612:ASN:OD1	2.12	0.50
1:A:1542:LEU:O	1:A:1545:ASP:CB	2.60	0.50
1:A:1572:ASN:OD1	1:A:1572:ASN:C	2.56	0.48
1:A:1543:PRO:C	1:A:1545:ASP:N	2.67	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1647:SER:O	1:A:1655:LYS:HG2	2.14	0.47
1:A:1707:ASP:O	1:A:1708:TYR:HB2	2.15	0.47
1:B:1547:GLU:OE2	1:B:1569:ARG:NH1	2.47	0.46
1:A:1534:GLN:OE1	1:A:1614:ASN:OD1	2.34	0.45
1:A:1536:PHE:HE2	1:A:1613:ARG:HE	1.63	0.45
1:A:1548:TYR:CE1	1:A:1567:ILE:HD12	2.53	0.44
1:B:1659:TYR:HB3	1:B:1714:TYR:HB2	2.01	0.42
1:B:1669:HIS:HB2	1:B:1692:VAL:HB	2.02	0.42
1:A:1534:GLN:NE2	1:A:1614:ASN:OD1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/193 (89%)	164 (96%)	7 (4%)	0	100	100
1	B	168/193 (87%)	166 (99%)	2 (1%)	0	100	100
All	All	339/386 (88%)	330 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/170 (90%)	150 (98%)	3 (2%)	48	70
1	B	153/170 (90%)	148 (97%)	5 (3%)	33	55
All	All	306/340 (90%)	298 (97%)	8 (3%)	40	63

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1542	LEU
1	A	1570	ILE
1	A	1688	LEU
1	B	1547	GLU
1	B	1666	ILE
1	B	1680	LYS
1	B	1692	VAL
1	B	1700	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1590	ASN
1	A	1652	ASN
1	B	1637	ASN
1	B	1681	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	QDR	B	2	-	20,22,22	3.36	9 (45%)	26,31,31	2.68	10 (38%)
2	QDR	A	1	-	20,22,22	3.09	8 (40%)	26,31,31	2.36	8 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QDR	B	2	-	-	1/3/5/5	0/3/3/3
2	QDR	A	1	-	-	2/3/5/5	0/3/3/3

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	QDR	CAO-NAI	-7.57	1.29	1.37
2	A	1	QDR	CAS-CAR	-7.36	1.34	1.47
2	B	2	QDR	NAH-NAK	-7.18	1.21	1.36
2	A	1	QDR	NAH-NAK	-6.75	1.22	1.36
2	B	2	QDR	CAQ-SAM	-5.58	1.68	1.75
2	B	2	QDR	CAS-CAR	-4.82	1.39	1.47
2	A	1	QDR	CAO-NAI	-4.65	1.32	1.37
2	A	1	QDR	CAR-NAL	-4.56	1.30	1.37
2	B	2	QDR	CAQ-NAH	-3.54	1.31	1.33
2	B	2	QDR	CAT-NAJ	-3.47	1.31	1.38
2	A	1	QDR	CAT-NAJ	-3.39	1.32	1.38
2	A	1	QDR	CAS-CAT	-3.35	1.36	1.41
2	B	2	QDR	CAR-NAL	-3.00	1.33	1.37
2	A	1	QDR	CAF-CAS	-2.91	1.35	1.39
2	B	2	QDR	CAS-CAT	-2.78	1.37	1.41
2	B	2	QDR	CAO-NAB	-2.26	1.29	1.34
2	A	1	QDR	CAQ-SAM	2.14	1.78	1.75

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	QDR	CAO-NAK-NAH	8.10	109.73	102.05
2	A	1	QDR	CAO-NAK-NAH	6.93	108.62	102.05
2	B	2	QDR	CAR-NAL-CAP	-6.51	120.14	123.90
2	A	1	QDR	CAR-NAL-CAP	-5.29	120.84	123.90
2	B	2	QDR	CAS-CAR-NAL	3.79	118.94	114.93
2	A	1	QDR	NAB-CAO-NAI	3.40	126.13	121.29
2	A	1	QDR	CAS-CAR-NAL	3.38	118.50	114.93
2	B	2	QDR	NAB-CAO-NAI	3.36	126.07	121.29
2	B	2	QDR	CAS-CAT-NAJ	-3.23	119.14	122.33
2	B	2	QDR	CAG-SAM-CAQ	2.96	107.52	100.98
2	A	1	QDR	NAK-CAO-NAI	-2.88	109.72	115.61
2	B	2	QDR	NAK-CAO-NAI	-2.47	110.55	115.61
2	B	2	QDR	CAQ-NAH-NAK	-2.45	107.16	116.48
2	A	1	QDR	CAS-CAT-NAJ	-2.27	120.09	122.33
2	B	2	QDR	NAL-CAP-NAJ	-2.17	120.32	123.08
2	B	2	QDR	CAP-CAG-SAM	-2.13	105.47	111.09
2	A	1	QDR	CAP-CAG-SAM	2.09	116.60	111.09
2	A	1	QDR	CAS-CAT-CAN	-2.07	118.82	120.56

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1	QDR	NAH-CAQ-SAM-CAG
2	A	1	QDR	NAI-CAQ-SAM-CAG
2	B	2	QDR	NAH-CAQ-SAM-CAG

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand QDR B 2



Ligand QDR A 1



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	177/193 (91%)	0.76	26 (14%) 6 4	26, 42, 80, 106	0
1	B	174/193 (90%)	0.30	8 (4%) 37 33	21, 34, 59, 82	0
All	All	351/386 (90%)	0.53	34 (9%) 13 10	21, 37, 71, 106	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1681	ASN	5.8
1	B	1669	HIS	4.9
1	A	1687	ASP	4.5
1	A	1621	ALA	4.3
1	A	1675	ILE	3.9
1	A	1627	ALA	3.8
1	B	1676	VAL	3.7
1	A	1698	HIS	3.3
1	A	1585	THR	3.1
1	A	1669	HIS	3.1
1	A	1588	ALA	3.0
1	A	1680	LYS	3.0
1	B	1533	GLN	2.8
1	A	1676	VAL	2.7
1	A	1672	HIS	2.7
1	A	1628	TYR	2.6
1	A	1620	TYR	2.5
1	B	1697	HIS	2.5
1	A	1619	SER	2.5
1	A	1534	GLN	2.5
1	A	1678	PRO	2.4
1	B	1720	LYS	2.4
1	A	1673	SER	2.3
1	B	1619	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1628	TYR	2.3
1	A	1674	LEU	2.3
1	B	1680	LYS	2.3
1	A	1581	ALA	2.2
1	A	1706	TYR	2.2
1	A	1589	LYS	2.2
1	A	1677	PRO	2.2
1	A	1582	LYS	2.1
1	A	1583	LYS	2.1
1	A	1671	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	QDR	B	2	20/20	0.90	0.15	20,43,160,194	0
2	QDR	A	1	20/20	0.94	0.12	16,30,162,230	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around QDR B 2:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around QDR A 1:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.