



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:51 PM UTC

PDB ID : 1SQA / pdb_00001sqa
Title : Substituted 2-Naphthamidine Inhibitors of Urokinase
Authors : Wendt, M.D.; Geyer, A.; McClellan, W.J.; Rockway, T.W.; Weitzberg, M.;
Zhao, X.; Stewart, K.; Nienaber, V.; Klinghofer, V.; Giranda, V.L.
Deposited on : 2004-03-18
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	245	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2041 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urokinase-type plasminogen activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	1934	1219	338	361	16	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	156	GLN	ASN	conflict	UNP P00749

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is 6-[(Z)-AMINO(IMINO)METHYL]-N-[4-(AMINOMETHYL)PHENYL]-4-(PYRIMIDIN-2-YLAMINO)-2-NAPHTHAMIDE (CCD ID: UI1) (formula: C₂₃H₂₁N₇O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			31	23	7	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	61	Total O 61 61	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Urokinase-type plasminogen activator



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.16Å 53.00Å 82.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.00 10.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	67.4 (10.00-2.00) 91.7 (10.00-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.92Å)	Xtriage
Refinement program	X-PLOR 98.0	Depositor
R, R_{free}	0.210 , 0.266 0.234 , 0.233	Depositor DCC
R_{free} test set	1702 reflections (10.03%)	wwPDB-VP
Wilson B-factor (Å ²)	11.6	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.50 , 61.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.069 for k,h,-l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	2041	wwPDB-VP
Average B, all atoms (Å ²)	7.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, UI1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.85	49/1983 (2.5%)	1.90	57/2685 (2.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (49) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	228	ILE	CA-CB	10.15	1.67	1.54
1	A	50	ILE	CA-CB	9.36	1.67	1.54
1	A	51	SER	CA-CB	8.76	1.68	1.53
1	A	245	THR	N-CA	7.84	1.55	1.46
1	A	145	THR	CA-C	7.82	1.62	1.52
1	A	230	SER	CA-C	-7.53	1.46	1.52
1	A	215	LEU	CA-CB	7.51	1.63	1.53
1	A	162	TYR	CA-C	7.45	1.62	1.53
1	A	154	LYS	N-CA	6.82	1.54	1.46
1	A	146	SER	CA-CB	-6.42	1.44	1.53
1	A	53	THR	CA-C	6.38	1.61	1.52
1	A	172	VAL	CA-CB	6.32	1.65	1.54
1	A	202	TRP	CA-CB	6.28	1.63	1.53
1	A	110	ILE	N-CA	-6.20	1.39	1.46
1	A	175	ILE	CB-CG1	6.17	1.65	1.53
1	A	99	TYR	CA-CB	6.09	1.63	1.53
1	A	198	ALA	N-CA	6.08	1.53	1.46
1	A	169	MET	CA-CB	5.90	1.66	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	99	TYR	N-CA	5.86	1.53	1.46
1	A	96	HIS	N-CA	5.86	1.53	1.46
1	A	109	ASP	CA-CB	5.85	1.61	1.52
1	A	251	LEU	CA-CB	5.84	1.61	1.53
1	A	179	GLU	CA-CB	5.83	1.62	1.53
1	A	190	VAL	CA-CB	5.83	1.61	1.54
1	A	74	SER	N-CA	5.81	1.53	1.45
1	A	175	ILE	N-CA	-5.80	1.39	1.46
1	A	94	ILE	CA-CB	5.79	1.61	1.53
1	A	41	MET	CB-CG	-5.73	1.35	1.52
1	A	67	TYR	C-O	5.64	1.30	1.23
1	A	196	CYS	CA-CB	-5.63	1.44	1.53
1	A	183	PRO	C-O	-5.63	1.17	1.24
1	A	173	LYS	N-CA	5.61	1.52	1.46
1	A	69	VAL	CA-CB	5.58	1.61	1.54
1	A	32	VAL	N-CA	5.56	1.53	1.46
1	A	54	HIS	CA-CB	5.53	1.63	1.53
1	A	50	ILE	C-O	5.51	1.29	1.24
1	A	113	LEU	CA-CB	5.50	1.62	1.53
1	A	189	GLU	N-CA	5.41	1.53	1.46
1	A	129	ILE	N-CA	5.39	1.54	1.46
1	A	74	SER	CA-CB	5.36	1.61	1.54
1	A	62	LYS	CA-CB	-5.27	1.44	1.53
1	A	160	TYR	CA-C	5.20	1.59	1.52
1	A	106	HIS	N-CA	5.19	1.52	1.45
1	A	96	HIS	CA-C	5.17	1.59	1.52
1	A	57	ILE	N-CA	5.14	1.52	1.46
1	A	247	VAL	CA-C	5.08	1.59	1.52
1	A	56	PHE	CA-CB	5.07	1.61	1.54
1	A	18	ILE	CA-CB	5.07	1.60	1.54
1	A	103	THR	CA-C	5.03	1.59	1.52

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	ILE	O-C-N	-11.65	110.68	123.26
1	A	169	MET	CA-CB-CG	10.47	135.05	114.10
1	A	65	ASP	CA-CB-CG	9.36	121.96	112.60
1	A	41	MET	CA-CB-CG	-9.20	95.70	114.10
1	A	13	PRO	N-CA-C	8.78	124.83	114.03
1	A	214	PRO	N-CA-C	8.62	124.72	111.19
1	A	238	LYS	CA-C-N	8.59	134.71	122.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	238	LYS	C-N-CA	8.59	134.71	122.08
1	A	20	ARG	NE-CZ-NH2	8.58	126.93	119.20
1	A	11	ASN	CA-CB-CG	-8.30	104.30	112.60
1	A	53	THR	O-C-N	-8.29	112.75	122.20
1	A	20	ARG	NE-CZ-NH1	-8.17	113.33	121.50
1	A	95	LEU	O-C-N	-8.14	113.94	123.06
1	A	74	SER	O-C-N	-8.10	112.68	122.48
1	A	241	PRO	N-CA-C	7.67	123.22	111.34
1	A	92	ASN	OD1-CG-ND2	-7.57	115.03	122.60
1	A	67	TYR	O-C-N	7.57	131.94	123.16
1	A	199	ASP	CA-CB-CG	7.17	119.77	112.60
1	A	220	GLN	CA-CB-CG	-6.94	100.22	114.10
1	A	181	GLN	OE1-CD-NE2	-6.94	115.66	122.60
1	A	73	ARG	CA-C-N	6.94	133.48	122.11
1	A	73	ARG	C-N-CA	6.94	133.48	122.11
1	A	69	VAL	CG1-CB-CG2	-6.87	95.69	110.80
1	A	130	GLN	OE1-CD-NE2	-6.63	115.97	122.60
1	A	68	ILE	O-C-N	-6.58	116.07	123.18
1	A	201	GLN	O-C-N	-6.50	114.54	122.34
1	A	106	HIS	O-C-N	-6.48	114.76	123.19
1	A	161	LEU	CD1-CG-CD2	-6.41	96.70	110.80
1	A	238	LYS	O-C-N	-6.39	115.39	123.06
1	A	89	GLU	O-C-N	-6.26	114.54	122.43
1	A	39	LEU	CA-C-N	6.17	134.78	121.64
1	A	39	LEU	C-N-CA	6.17	134.78	121.64
1	A	48	TRP	O-C-N	-6.11	116.16	123.31
1	A	210	ASP	CA-CB-CG	5.88	118.48	112.60
1	A	60	PRO	O-C-N	-5.69	114.96	122.64
1	A	201	GLN	OE1-CD-NE2	-5.69	116.91	122.60
1	A	21	ARG	NE-CZ-NH2	-5.64	114.12	119.20
1	A	127	ARG	CA-CB-CG	-5.47	103.15	114.10
1	A	255	ARG	NE-CZ-NH2	-5.46	114.29	119.20
1	A	197	ALA	O-C-N	-5.33	116.89	123.33
1	A	181	GLN	O-C-N	-5.30	114.55	122.39
1	A	226	THR	O-C-N	-5.29	115.36	122.23
1	A	109	ASP	CA-CB-CG	5.27	117.87	112.60
1	A	83	GLN	O-C-N	5.26	129.47	122.95
1	A	216	VAL	CG1-CB-CG2	-5.23	99.29	110.80
1	A	250	PHE	O-C-N	-5.19	115.97	122.35
1	A	83	GLN	OE1-CD-NE2	-5.18	117.42	122.60
1	A	148	GLU	CA-C-N	5.17	130.92	122.69
1	A	148	GLU	C-N-CA	5.17	130.92	122.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	TYR	O-C-N	-5.14	116.02	122.24
1	A	139	ASN	CA-CB-CG	5.11	117.71	112.60
1	A	160	TYR	CA-C-N	5.10	130.82	122.81
1	A	160	TYR	C-N-CA	5.10	130.82	122.81
1	A	7	THR	O-C-N	-5.10	116.86	122.72
1	A	206	SER	O-C-N	-5.09	117.05	122.85
1	A	128	THR	O-C-N	-5.06	116.12	122.24
1	A	145	THR	OG1-CB-CG2	-5.06	99.19	109.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	153	GLY	Mainchain
1	A	250	PHE	Mainchain
1	A	255	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1934	0	1880	29	0
2	A	15	0	0	0	0
3	A	31	0	20	0	0
4	A	61	0	0	3	0
All	All	2041	0	1900	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:HIS:HD2	1:A:98:ASP:H	1.26	0.82
1:A:76:ARG:HG2	1:A:76:ARG:HH11	1.57	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:MET:HG2	1:A:251:LEU:HD12	1.80	0.64
1:A:255:ARG:HG2	1:A:255:ARG:HH11	1.61	0.64
1:A:76:ARG:HG2	1:A:76:ARG:NH1	2.18	0.59
1:A:41:MET:HE2	1:A:225:LEU:HD22	1.84	0.58
1:A:62:LYS:HD2	1:A:90:VAL:O	2.04	0.58
1:A:195:LEU:HB3	4:A:1029:HOH:O	2.05	0.55
1:A:253:TRP:O	1:A:257:HIS:HD2	1.92	0.53
1:A:195:LEU:HG	1:A:246:ARG:HB2	1.90	0.52
1:A:71:LEU:HD12	1:A:88:PHE:CE1	2.49	0.48
1:A:39:LEU:HG	1:A:131:THR:HG22	1.96	0.48
1:A:41:MET:HE3	1:A:134:LEU:HD21	1.96	0.47
1:A:34:CYS:CB	1:A:55:CYS:SG	3.01	0.47
1:A:137:MET:HE1	1:A:252:PRO:HG3	1.97	0.47
1:A:105:ALA:HA	1:A:231:TRP:CE2	2.51	0.46
1:A:99:TYR:HA	1:A:107:HIS:O	2.16	0.45
1:A:96:HIS:CD2	1:A:98:ASP:H	2.18	0.45
1:A:41:MET:HE2	1:A:225:LEU:CD2	2.47	0.44
1:A:177:HIS:HD2	4:A:1010:HOH:O	2.01	0.43
1:A:112:LEU:C	1:A:113:LEU:HD12	2.44	0.42
1:A:76:ARG:HH11	1:A:165:GLN:HE21	1.68	0.42
1:A:135:PRO:O	1:A:251:LEU:HD11	2.20	0.41
1:A:178:ARG:HD3	1:A:178:ARG:HA	1.87	0.41
1:A:225:LEU:HD21	1:A:228:ILE:HD11	2.02	0.41
1:A:121:ARG:HH11	1:A:121:ARG:HD2	1.77	0.41
1:A:76:ARG:HH11	1:A:165:GLN:NE2	2.19	0.40
1:A:207:CYS:HA	1:A:235:CYS:SG	2.61	0.40
1:A:22:HIS:HD2	4:A:1043:HOH:O	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/245 (99%)	234 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/214 (100%)	200 (94%)	13 (6%)	17	14

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	ILE
1	A	21	ARG
1	A	53	THR
1	A	54	HIS
1	A	60	PRO
1	A	61	LYS
1	A	76	ARG
1	A	83	GLN
1	A	137	MET
1	A	150	THR
1	A	169	MET
1	A	255	ARG
1	A	256	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	HIS
1	A	92	ASN
1	A	96	HIS
1	A	107	HIS
1	A	130	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	156	GLN
1	A	165	GLN
1	A	177	HIS
1	A	181	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	302	-	4,4,4	0.82	0	6,6,6	1.94	2 (33%)
2	SO4	A	301	-	4,4,4	1.06	0	6,6,6	1.00	0
3	UI1	A	1001	-	34,34,34	1.45	7 (20%)	44,47,47	2.20	9 (20%)
2	SO4	A	303	-	4,4,4	0.56	0	6,6,6	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UI1	A	1001	-	-	8/18/18/18	0/4/4/4

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	UI1	C26-C10	3.43	1.43	1.37
3	A	1001	UI1	C31-C30	2.96	1.43	1.38
3	A	1001	UI1	C14-N13	2.80	1.42	1.36
3	A	1001	UI1	C24-C23	2.52	1.42	1.36
3	A	1001	UI1	C5-C6	2.46	1.43	1.39
3	A	1001	UI1	C21-C22	2.27	1.41	1.37
3	A	1001	UI1	C12-N13	2.16	1.44	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	UI1	C18-N19-C14	6.52	120.87	115.42
3	A	1001	UI1	C16-N15-C14	6.44	120.80	115.42
3	A	1001	UI1	C21-C20-C12	-6.09	119.25	123.67
3	A	1001	UI1	N19-C14-N15	-5.52	120.83	126.43
2	A	302	SO4	O4-S-O2	3.19	126.26	109.56
2	A	302	SO4	O4-S-O3	-2.97	92.19	108.54
3	A	1001	UI1	C17-C16-N15	-2.36	119.69	123.42
3	A	1001	UI1	C17-C18-N19	-2.30	119.78	123.42
3	A	1001	UI1	C11-C12-N13	-2.06	118.89	123.20
3	A	1001	UI1	C5-C6-N7	-2.04	113.57	120.41
3	A	1001	UI1	C6-N7-C8	-2.03	121.25	126.61

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1001	UI1	C5-C6-N7-C8
3	A	1001	UI1	C30-C6-N7-C8
3	A	1001	UI1	C20-C12-N13-C14
3	A	1001	UI1	C21-C22-C27-N29
3	A	1001	UI1	C23-C22-C27-N29
3	A	1001	UI1	C11-C12-N13-C14
3	A	1001	UI1	C21-C22-C27-N28
3	A	1001	UI1	C23-C22-C27-N28

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/245 (100%)	0.02	3 (1%) 76 76	2, 4, 18, 33	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	41	MET	2.9
1	A	23	ARG	2.7
1	A	178	ARG	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	302	5/5	0.74	0.20	21,21,26,29	0
3	UI1	A	1001	31/31	0.78	0.12	2,9,15,19	0
2	SO4	A	303	5/5	0.95	0.09	35,36,37,38	0
2	SO4	A	301	5/5	0.98	0.04	8,11,13,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.