



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:25 PM UTC

PDB ID : 3SQ2 / pdb_00003sq2
Title : RB69 DNA Polymerase Ternary Complex with dTTP Opposite 2AP (AT rich sequence)
Authors : Xia, S.; Konigsberg, W.H.; Wang, J.
Deposited on : 2011-07-04
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	902	3% 92% 7%
2	T	16	81% 12% 6%
3	P	13	8% 69% 31%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	902	7375	4736	1228	1378	33	0	2	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	ALA	ASP	engineered mutation	UNP Q38087
A	327	ALA	ASP	engineered mutation	UNP Q38087


- Molecule 2 is a DNA chain called 5'-D(P*CP*(2PR)P*TP*AP*AP*TP*TP*AP*AP*TP*T P*AP*AP*TP*TP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	T	16	328	159	57	96	16	0	0	0

- Molecule 3 is a DNA chain called 5'-D(*AP*AP*TP*TP*AP*AP*TP*TP*AP*AP*TP*TP *(2DA))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	P	13	263	130	47	74	12	0	0	0

- Molecule 4 is THYMIDINE-5'-TRIPHOSPHATE (CCD ID: TTP) (formula: C₁₀H₁₇N₂O₁₄P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			29	10	2	14	3		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	5	Total Ca 5 5	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	311	Total O 313 313	0	2
6	T	26	Total O 26 26	0	0
6	P	11	Total O 11 11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase



- Molecule 2: 5'-D(P*CP*(2PR)P*TP*AP*AP*TP*TP*AP*AP*TP*TP*AP*AP*TP*TP*G)-3'



- Molecule 3: 5'-D(*AP*AP*TP*TP*AP*AP*TP*TP*AP*AP*TP*TP*(2DA))-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.95Å 119.88Å 130.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.26 – 2.10 88.26 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.7 (88.26-2.10) 98.0 (88.26-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.202 , 0.244 0.206 , 0.249	Depositor DCC
R_{free} test set	3447 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtriage
Anisotropy	0.523	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8350	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2PR, TTP, CA, 2DA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/7562	0.75	0/10219
2	T	0.29	0/342	0.95	0/523
3	P	0.31	0/272	0.98	0/418
All	All	0.47	0/8176	0.77	0/11160

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7375	0	7271	38	0
2	T	328	0	184	3	0
3	P	263	0	151	2	0
4	A	29	0	13	0	0
5	A	5	0	0	0	0
6	A	313	0	0	5	0
6	P	11	0	0	0	0
6	T	26	0	0	1	0
All	All	8350	0	7619	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:408:MET:HE1	1:A:655:ALA:HB2	1.68	0.76
1:A:508:LEU:HD12	1:A:508:LEU:H	1.54	0.72
1:A:206:GLN:HE22	1:A:241:ARG:HE	1.39	0.69
1:A:739:LYS:O	1:A:739:LYS:HE3	1.95	0.66
1:A:449:ARG:NH1	1:A:452:ASP:OD1	2.32	0.62
2:T:17:DT:H6	6:T:2:HOH:O	1.83	0.60
1:A:131:HIS:HD2	1:A:156:TYR:OH	1.85	0.60
1:A:206:GLN:NE2	1:A:241:ARG:HE	2.00	0.59
1:A:705:LYS:HD3	2:T:8:DT:H5''	1.86	0.56
1:A:779:ILE:HD11	1:A:866:MET:HE1	1.88	0.56
1:A:191:PHE:HD2	1:A:196:GLU:HG3	1.71	0.56
3:P:105:DT:H2''	3:P:106:DT:OP2	2.09	0.53
1:A:514:LEU:HD13	1:A:533:LEU:HD11	1.91	0.53
1:A:35:PRO:HG3	1:A:65:MET:HG2	1.92	0.51
1:A:126:PRO:HB3	1:A:224:PRO:HB2	1.91	0.51
1:A:41:CYS:HB2	1:A:42:PRO:HD2	1.92	0.50
1:A:286:PRO:HB3	1:A:782:VAL:HG21	1.93	0.50
1:A:112[B]:ASN:ND2	6:A:939:HOH:O	2.44	0.50
1:A:873:GLU:HG2	6:A:963:HOH:O	2.14	0.48
1:A:253:ILE:HD13	1:A:254:GLU:H	1.79	0.48
1:A:203:ASN:ND2	1:A:241:ARG:HH22	2.12	0.47
1:A:441:ASP:HB3	1:A:447:ALA:HB2	1.96	0.46
1:A:668:ARG:HD3	6:A:964:HOH:O	2.14	0.46
1:A:191:PHE:CD2	1:A:196:GLU:HG3	2.50	0.46
1:A:878:LYS:HB3	1:A:879:PRO:HD3	1.98	0.46
3:P:104:DA:H2''	3:P:105:DT:OP2	2.15	0.46
1:A:121:ASP:HA	1:A:819:ILE:HD13	1.98	0.45
1:A:405:LYS:O	1:A:690:GLY:HA2	2.17	0.44
1:A:508:LEU:HD12	1:A:508:LEU:N	2.28	0.43
1:A:819:ILE:H	1:A:819:ILE:HG13	1.70	0.43
1:A:818:ASN:HB3	6:A:968:HOH:O	2.18	0.43
1:A:112[A]:ASN:HD22	1:A:332:LEU:HG	1.84	0.43
1:A:749:ILE:O	1:A:753:LEU:HG	2.19	0.42
1:A:870:VAL:HG13	1:A:874:LYS:HD3	2.01	0.42
1:A:482:ARG:HD3	1:A:482:ARG:C	2.45	0.42
1:A:153:ASN:HB2	1:A:192:ASP:O	2.20	0.42
1:A:415:LEU:HD22	1:A:623:ASP:HB3	2.02	0.42
1:A:347:MET:HE1	6:A:1025:HOH:O	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:813:ARG:HA	1:A:816:LYS:HE3	2.00	0.42
1:A:360:SER:OG	2:T:4:2PR:OP1	2.24	0.41
1:A:412:LEU:HD13	1:A:415:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	902/902 (100%)	874 (97%)	27 (3%)	1 (0%)	48 50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	801/799 (100%)	779 (97%)	22 (3%)	39 45

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	LEU
1	A	206	GLN
1	A	217	ASN
1	A	253	ILE
1	A	254	GLU
1	A	295	GLU
1	A	303	LEU
1	A	309	ILE
1	A	482	ARG
1	A	508	LEU
1	A	580	LEU
1	A	612	GLU
1	A	636	VAL
1	A	686	GLU
1	A	693	LEU
1	A	739	LYS
1	A	760	LEU
1	A	765	LYS
1	A	769	LYS
1	A	819	ILE
1	A	835	LEU
1	A	897	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	131	HIS
1	A	203	ASN
1	A	206	GLN
1	A	317	HIS
1	A	333	GLN
1	A	539	ASN
1	A	546	GLN
1	A	733	GLN
1	A	761	GLN
1	A	773	GLN
1	A	787	ASN
1	A	823	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	2DA	P	115	2,3	19,22,23	1.06	1 (5%)	25,31,34	2.62	11 (44%)
2	2PR	T	4	2	20,23,24	1.11	1 (5%)	27,33,36	2.21	13 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2DA	P	115	2,3	-	4/7/18/19	0/3/3/3
2	2PR	T	4	2	-	1/7/21/22	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	4	2PR	C8-N7	2.55	1.36	1.31
3	P	115	2DA	C8-N7	2.40	1.36	1.31

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	115	2DA	C4'-O4'-C1'	-5.64	104.49	109.81
3	P	115	2DA	N3-C2-N1	-5.40	120.41	128.58
2	T	4	2PR	C5-C4-N3	-4.57	120.38	126.55
2	T	4	2PR	N1-C2-N3	-4.52	120.67	125.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	115	2DA	C5-C4-N3	-4.45	120.58	126.72
3	P	115	2DA	N9-C8-N7	-3.91	108.39	113.94
2	T	4	2PR	N9-C8-N7	-3.75	108.61	113.94
2	T	4	2PR	N3-C4-N9	3.52	131.46	126.99
3	P	115	2DA	C2-N3-C4	3.45	120.25	111.83
2	T	4	2PR	N2-C2-N1	3.38	120.89	117.40
3	P	115	2DA	C4-N9-C8	3.19	109.09	105.74
3	P	115	2DA	O4'-C4'-C5'	3.05	114.84	109.34
3	P	115	2DA	O4'-C1'-N9	2.89	113.15	107.96
3	P	115	2DA	N3-C4-N9	2.88	132.06	127.17
2	T	4	2PR	C5-N7-C8	2.82	107.86	104.40
2	T	4	2PR	C2'-C1'-N9	-2.74	108.19	114.63
2	T	4	2PR	C4-N9-C8	2.66	108.53	105.74
2	T	4	2PR	C6-N1-C2	2.57	120.81	116.29
3	P	115	2DA	C5-N7-C8	2.52	107.42	103.45
3	P	115	2DA	C4-C5-N7	-2.48	107.75	110.58
2	T	4	2PR	C2-N3-C4	2.40	119.93	113.51
2	T	4	2PR	O4'-C1'-N9	2.26	112.02	107.96
2	T	4	2PR	C6-C5-C4	2.24	118.69	116.24
2	T	4	2PR	C4-C5-N7	-2.05	106.83	109.94

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	P	115	2DA	O4'-C4'-C5'-O5'
3	P	115	2DA	C2'-C1'-N9-C4
3	P	115	2DA	O4'-C1'-N9-C4
3	P	115	2DA	C3'-C4'-C5'-O5'
2	T	4	2PR	O4'-C1'-N9-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	T	4	2PR	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	TTP	A	903	5	29,30,30	1.28	4 (13%)	43,47,47	1.83	10 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TTP	A	903	5	-	3/22/34/34	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	903	TTP	C6-C5	3.06	1.39	1.34
4	A	903	TTP	C4-C5	2.57	1.49	1.44
4	A	903	TTP	C2-N1	2.47	1.42	1.38
4	A	903	TTP	C4-N3	-2.17	1.34	1.38

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	903	TTP	N3-C2-N1	4.92	121.30	114.89
4	A	903	TTP	C4-N3-C2	-4.87	120.96	127.34
4	A	903	TTP	C5-C4-N3	4.28	119.04	115.32
4	A	903	TTP	O4-C4-C5	-3.66	120.73	124.92
4	A	903	TTP	O4'-C1'-N1	3.37	113.85	107.86
4	A	903	TTP	C2'-C1'-N1	-3.08	106.12	113.81
4	A	903	TTP	C5-C6-N1	-3.02	120.04	123.31
4	A	903	TTP	O2-C2-N1	-2.65	119.34	122.80
4	A	903	TTP	O2B-PB-O3A	2.62	114.36	107.27
4	A	903	TTP	C5M-C5-C4	2.16	121.08	118.78

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	903	TTP	PB-O3B-PG-O2G
4	A	903	TTP	PB-O3B-PG-O3G
4	A	903	TTP	PB-O3B-PG-O1G

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	902/902 (100%)	0.07	27 (2%) 52 55	16, 35, 64, 121	2 (0%)
2	T	15/16 (93%)	-0.43	0 100 100	26, 36, 60, 62	0
3	P	12/13 (92%)	0.26	1 (8%) 17 18	27, 50, 91, 95	0
All	All	929/931 (99%)	0.07	28 (3%) 52 55	16, 35, 66, 121	2 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	253	ILE	9.0
1	A	257	TYR	5.6
1	A	252	VAL	5.3
1	A	255	ASN	4.8
1	A	259	SER	4.8
1	A	508	LEU	4.6
1	A	256	MET	4.3
1	A	857	LEU	4.3
1	A	771	PHE	3.8
1	A	901	PHE	3.7
1	A	258	GLY	3.5
1	A	819	ILE	3.4
1	A	852	THR	3.3
1	A	254	GLU	3.1
1	A	260	ARG	2.9
1	A	251	LYS	2.8
3	P	103	DA	2.8
1	A	510	VAL	2.7
1	A	851	GLY	2.7
1	A	895	ALA	2.6
1	A	821	ALA	2.5
1	A	42	PRO	2.3
1	A	739	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	789	ALA	2.2
1	A	43	GLU	2.1
1	A	898	PHE	2.1
1	A	303	LEU	2.0
1	A	525	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	2PR	T	4	21/22	0.95	0.07	22,26,31,37	0
3	2DA	P	115	20/21	0.98	0.05	23,25,32,32	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CA	A	907	1/1	0.90	0.18	80,80,80,80	0
4	TTP	A	903	29/29	0.94	0.12	33,42,47,49	0
5	CA	A	908	1/1	0.95	0.07	55,55,55,55	0
5	CA	A	905	1/1	0.97	0.04	51,51,51,51	0
5	CA	A	906	1/1	0.97	0.08	54,54,54,54	0
5	CA	A	904	1/1	0.99	0.02	31,31,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.