



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 01:15 PM UTC

PDB ID : 3SQN / pdb_00003sqn
Title : Putative Mga family transcriptional regulator from *Enterococcus faecalis*
Authors : Osipiuk, J.; Wu, R.; Jedrzejczak, R.; Moy, S.; Joachimiak, A.; Midwest Center
for Structural Genomics (MCSG)
Deposited on : 2011-07-05
Resolution : 2.31 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	7% 70% 24% • •
1	B	485	13% 51% 22% 6% 21%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7103 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Conserved domain protein.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	468	Total	C	N	O	S	Se	0	1	0
			3883	2519	628	728	3	5			
1	B	381	Total	C	N	O	S	Se	0	3	0
			3181	2076	498	600	3	4			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q82ZN7
A	-1	ASN	-	expression tag	UNP Q82ZN7
A	0	ALA	-	expression tag	UNP Q82ZN7
B	-2	SER	-	expression tag	UNP Q82ZN7
B	-1	ASN	-	expression tag	UNP Q82ZN7
B	0	ALA	-	expression tag	UNP Q82ZN7

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	26	Total	O	0	0
			26	26		
2	B	13	Total	O	0	0
			13	13		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.80Å 101.76Å 89.90Å 90.00° 96.75° 90.00°	Depositor
Resolution (Å)	36.17 – 2.31 36.17 – 2.31	Depositor EDS
% Data completeness (in resolution range)	97.2 (36.17-2.31) 94.8 (36.17-2.31)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.224 , 0.282 (Not available) , 0.286	Depositor DCC
R_{free} test set	2439 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7103	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	5/3965 (0.1%)	1.05	5/5365 (0.1%)
1	B	1.10	2/3260 (0.1%)	1.09	5/4408 (0.1%)
All	All	1.08	7/7225 (0.1%)	1.07	10/9773 (0.1%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	111	PHE	CA-C	7.28	1.62	1.52
1	A	213	THR	N-CA	-5.43	1.39	1.46
1	A	418	ALA	C-O	-5.41	1.17	1.24
1	A	112	LEU	N-CA	5.13	1.52	1.46
1	B	156	ILE	CA-CB	5.11	1.60	1.54

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	447	SER	N-CA-C	6.75	118.45	108.60
1	B	312	THR	N-CA-C	-5.70	101.56	110.17
1	B	164	LEU	N-CA-C	5.54	121.81	114.12
1	B	338	GLU	N-CA-C	5.52	117.30	111.28
1	A	113	SER	N-CA-C	5.46	117.67	111.11

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3883	0	3876	87	0
1	B	3181	0	3130	126	0
2	A	26	0	0	0	0
2	B	13	0	0	1	0
All	All	7103	0	7006	204	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 204 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:114:THR:HG22	1:B:114:THR:O	1.68	0.92
1:B:112:LEU:HD11	1:B:119:TYR:HA	1.53	0.90
1:B:340:GLN:O	1:B:344:THR:HG23	1.72	0.90
1:A:451:LEU:HD23	1:A:451:LEU:O	1.73	0.88
1:B:133:LEU:HD11	1:B:140:ILE:CD1	2.05	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	461/485 (95%)	434 (94%)	24 (5%)	3 (1%)	18	22
1	B	374/485 (77%)	348 (93%)	22 (6%)	4 (1%)	11	12
All	All	835/970 (86%)	782 (94%)	46 (6%)	7 (1%)	16	19

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	36	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	94	ILE
1	A	35	LEU
1	B	432	VAL
1	B	93	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	434/439 (99%)	407 (94%)	27 (6%)	16	23
1	B	356/439 (81%)	315 (88%)	41 (12%)	5	6
All	All	790/878 (90%)	722 (91%)	68 (9%)	9	12

5 of 68 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	409	LYS
1	B	433	GLU
1	B	471	ARG
1	A	451	LEU
1	A	440	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	95	GLN
1	B	131	GLN
1	B	386	GLN
1	B	299	ASN
1	A	336	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	463/485 (95%)	0.59	35 (7%) 20 22	26, 53, 90, 105	1 (0%)
1	B	377/485 (77%)	1.03	63 (16%) 4 5	29, 59, 91, 110	3 (0%)
All	All	840/970 (86%)	0.79	98 (11%) 9 10	26, 55, 90, 110	4 (0%)

The worst 5 of 98 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	149	LEU	5.4
1	B	393	ALA	5.3
1	B	147	ILE	5.2
1	B	92	ILE	5.2
1	A	351	LEU	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.