



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 06:58 PM UTC

PDB ID : 6SSI / pdb_00006ssi
Title : Structure of the pentameric ligand-gated ion channel ELIC in complex with a PAM nanobody
Authors : Ulens, C.; Brams, M.; Evans, G.L.; Spurny, R.; Govaerts, C.; Pardon, E.; Steyaert, J.
Deposited on : 2019-09-07
Resolution : 2.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	12% 89% 6% 5%
1	B	318	13% 92% 5% .
1	C	318	14% 93% . .
1	D	318	15% 90% 6% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	318	14% 90% 6%
2	F	121	33% 91% 9%
2	G	121	11% 90% 8%
2	H	121	7% 88% 7%
2	I	121	20% 88% 8%
2	J	121	37% 89% 9%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 16941 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cys-loop ligand-gated ion channel.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	303	Total	C	N	O	S	0	0	0
			2453	1600	410	437	6			
1	B	308	Total	C	N	O	S	0	1	0
			2477	1612	412	447	6			
1	C	307	Total	C	N	O	S	0	0	0
			2461	1608	406	441	6			
1	D	305	Total	C	N	O	S	0	1	0
			2424	1583	403	432	6			
1	E	304	Total	C	N	O	S	0	0	0
			2457	1606	406	439	6			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	GLY	-	expression tag	UNP P0C7B7
A	6	PRO	-	expression tag	UNP P0C7B7
A	7	VAL	-	expression tag	UNP P0C7B7
A	164	GLY	-	insertion	UNP P0C7B7
A	289	ASN	MET	conflict	UNP P0C7B7
A	322	GLY	-	expression tag	UNP P0C7B7
B	5	GLY	-	expression tag	UNP P0C7B7
B	6	PRO	-	expression tag	UNP P0C7B7
B	7	VAL	-	expression tag	UNP P0C7B7
B	164	GLY	-	insertion	UNP P0C7B7
B	289	ASN	MET	conflict	UNP P0C7B7
B	322	GLY	-	expression tag	UNP P0C7B7
C	5	GLY	-	expression tag	UNP P0C7B7
C	6	PRO	-	expression tag	UNP P0C7B7
C	7	VAL	-	expression tag	UNP P0C7B7
C	164	GLY	-	insertion	UNP P0C7B7
C	289	ASN	MET	conflict	UNP P0C7B7
C	322	GLY	-	expression tag	UNP P0C7B7
D	5	GLY	-	expression tag	UNP P0C7B7

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	6	PRO	-	expression tag	UNP P0C7B7
D	7	VAL	-	expression tag	UNP P0C7B7
D	164	GLY	-	insertion	UNP P0C7B7
D	289	ASN	MET	conflict	UNP P0C7B7
D	322	GLY	-	expression tag	UNP P0C7B7
E	5	GLY	-	expression tag	UNP P0C7B7
E	6	PRO	-	expression tag	UNP P0C7B7
E	7	VAL	-	expression tag	UNP P0C7B7
E	164	GLY	-	insertion	UNP P0C7B7
E	289	ASN	MET	conflict	UNP P0C7B7
E	322	GLY	-	expression tag	UNP P0C7B7

- Molecule 2 is a protein called NANOBODY 22.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	110	Total	C	N	O	S	0	0	0
			815	502	146	162	5			
2	G	111	Total	C	N	O	S	0	0	0
			838	516	151	166	5			
2	H	112	Total	C	N	O	S	0	0	0
			843	519	152	167	5			
2	I	111	Total	C	N	O	S	0	0	0
			841	516	154	166	5			
2	J	110	Total	C	N	O	S	0	0	0
			831	510	152	164	5			

- Molecule 3 is GAMMA-AMINO-BUTANOIC ACID (CCD ID: ABU) (formula: C₄H₉NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			7	4	1	2		
3	B	1	Total	C	N	O	0	0
			7	4	1	2		
3	C	1	Total	C	N	O	0	0
			7	4	1	2		
3	D	1	Total	C	N	O	0	0
			7	4	1	2		
3	E	1	Total	C	N	O	0	0
			7	4	1	2		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		
5	E	1	Total	C	O	0	0
			6	3	3		
5	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		
6	B	1	Total	Ca	0	0
			1	1		
6	C	1	Total	Ca	0	0
			1	1		
6	D	1	Total	Ca	0	0
			1	1		
6	E	1	Total	Ca	0	0
			1	1		

- Molecule 7 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			13	8	5		
7	C	1	Total	C	O	0	0
			13	8	5		
7	D	1	Total	C	O	0	0
			13	8	5		
7	E	1	Total	C	O	0	0
			13	8	5		

- Molecule 8 is UNDECYL-MALTOSIDE (CCD ID: UMQ) (formula: C₂₃H₄₄O₁₁).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			32	21	11		
8	B	1	Total	C	O	0	0
			32	21	11		
8	C	1	Total	C	O	0	0
			32	21	11		
8	E	1	Total	C	O	0	0
			32	21	11		
8	E	1	Total	C	O	0	0
			32	21	11		

- Molecule 9 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	D	1	Total	C	O	0	0
			4	2	2		
9	E	1	Total	C	O	0	0
			4	2	2		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	18	Total	O	0	0
			18	18		
10	B	22	Total	O	0	0
			22	22		
10	C	31	Total	O	0	0
			31	31		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	D	18	Total 18	O 18	0	0
10	E	16	Total 16	O 16	0	0
10	G	14	Total 14	O 14	0	0
10	H	18	Total 18	O 18	0	0
10	I	6	Total 6	O 6	0	0
10	J	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cys-loop ligand-gated ion channel



• Molecule 1: Cys-loop ligand-gated ion channel




• Molecule 1: Cys-loop ligand-gated ion channel





• Molecule 1: Cys-loop ligand-gated ion channel








- Molecule 1: Cys-loop ligand-gated ion channel




- Molecule 2: NANOBODY 22





- Molecule 2: NANOBODY 22




- Molecule 2: NANOBODY 22



- Molecule 2: NANOBODY 22



- Molecule 2: NANOBODY 22



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	105.60Å 146.56Å 140.24Å 90.00° 111.39° 90.00°	Depositor
Resolution (Å)	47.06 – 2.59 47.06 – 2.59	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.06-2.59) 99.9 (47.06-2.59)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.58Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.230 , 0.246 0.231 , 0.246	Depositor DCC
R_{free} test set	6188 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	68.3	Xtriage
Anisotropy	0.211	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.013 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	16941	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, ABU, PG4, GOL, UMQ, CA, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.66	0/2519	1.07	2/3432 (0.1%)
1	B	0.66	0/2543	1.06	3/3467 (0.1%)
1	C	0.66	0/2527	1.06	1/3448 (0.0%)
1	D	0.67	0/2490	1.05	1/3401 (0.0%)
1	E	0.66	0/2523	1.06	6/3439 (0.2%)
2	F	0.60	0/827	0.93	0/1121
2	G	0.64	0/850	0.92	0/1149
2	H	0.71	1/855 (0.1%)	0.99	0/1156
2	I	0.63	1/853 (0.1%)	0.93	0/1152
2	J	0.64	0/843	0.96	0/1140
All	All	0.66	2/16830 (0.0%)	1.03	13/22905 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	82	MET	SD-CE	-6.24	1.64	1.79
2	I	82	MET	SD-CE	-5.19	1.66	1.79

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	136	ASN	CA-C-N	5.73	128.24	120.38
1	E	136	ASN	C-N-CA	5.73	128.24	120.38
1	B	138	GLN	N-CA-C	-5.32	106.84	113.38
1	D	286	ARG	N-CA-C	-5.26	106.71	113.23
1	A	306	LEU	CA-C-N	5.24	125.79	119.98

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2453	0	2404	13	0
1	B	2477	0	2399	9	0
1	C	2461	0	2396	7	0
1	D	2424	0	2330	11	0
1	E	2457	0	2416	8	0
2	F	815	0	773	0	0
2	G	838	0	813	1	0
2	H	843	0	815	3	0
2	I	841	0	815	2	0
2	J	831	0	799	2	0
3	A	7	0	0	0	0
3	B	7	0	0	0	0
3	C	7	0	0	0	0
3	D	7	0	0	0	0
3	E	7	0	0	1	0
4	A	12	0	13	0	0
4	B	12	0	13	0	0
4	C	12	0	13	0	0
4	D	12	0	13	0	0
4	E	12	0	13	0	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
5	C	6	0	8	0	0
5	D	6	0	8	0	0
5	E	12	0	16	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	E	1	0	0	0	0
7	B	13	0	18	0	0
7	C	13	0	18	0	0
7	D	13	0	18	2	0
7	E	13	0	18	0	0
8	B	64	0	74	0	0
8	C	32	0	37	0	0
8	E	64	0	74	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	D	4	0	3	0	0
9	E	4	0	3	0	0
10	A	18	0	0	0	0
10	B	22	0	0	1	0
10	C	31	0	0	0	0
10	D	18	0	0	0	0
10	E	16	0	0	0	0
10	G	14	0	0	0	0
10	H	18	0	0	0	0
10	I	6	0	0	0	0
10	J	2	0	0	0	0
All	All	16941	0	16336	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:ALA:HB2	1:A:297:ILE:HG21	1.85	0.59
2:I:40:ALA:HB3	2:I:43:LYS:HB2	1.85	0.58
1:E:152:ILE:HG21	1:E:162:ILE:HG13	1.85	0.57
1:D:123:ARG:HG2	1:D:198:VAL:HG22	1.86	0.57
1:D:76:LEU:H	7:D:404:PG4:H52	1.69	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	297/318 (93%)	289 (97%)	8 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	303/318 (95%)	292 (96%)	10 (3%)	1 (0%)	36	58
1	C	301/318 (95%)	292 (97%)	9 (3%)	0	100	100
1	D	300/318 (94%)	293 (98%)	7 (2%)	0	100	100
1	E	298/318 (94%)	293 (98%)	5 (2%)	0	100	100
2	F	108/121 (89%)	106 (98%)	2 (2%)	0	100	100
2	G	109/121 (90%)	109 (100%)	0	0	100	100
2	H	110/121 (91%)	107 (97%)	3 (3%)	0	100	100
2	I	109/121 (90%)	104 (95%)	5 (5%)	0	100	100
2	J	108/121 (89%)	100 (93%)	8 (7%)	0	100	100
All	All	2043/2195 (93%)	1985 (97%)	57 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	295	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/282 (94%)	264 (100%)	0	100	100
1	B	264/282 (94%)	262 (99%)	2 (1%)	73	88
1	C	263/282 (93%)	262 (100%)	1 (0%)	84	93
1	D	254/282 (90%)	251 (99%)	3 (1%)	63	83
1	E	266/282 (94%)	261 (98%)	5 (2%)	50	75
2	F	83/98 (85%)	83 (100%)	0	100	100
2	G	88/98 (90%)	88 (100%)	0	100	100
2	H	88/98 (90%)	88 (100%)	0	100	100
2	I	88/98 (90%)	88 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	J	86/98 (88%)	86 (100%)	0	100	100
All	All	1744/1900 (92%)	1733 (99%)	11 (1%)	78	91

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	150	GLU
1	E	247	PHE
1	E	302	LEU
1	E	298	GLN
1	D	157	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	264	GLN
2	J	13	GLN
1	E	138	GLN
2	H	106	GLN
1	E	137	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 32 ligands modelled in this entry, 5 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	UMQ	B	407	-	33,33,35	0.22	0	44,44,46	0.65	1 (2%)
5	GOL	C	403	-	5,5,5	0.12	0	5,5,5	0.45	0
4	MES	E	402	-	12,12,12	1.19	1 (8%)	15,16,16	0.48	0
8	UMQ	E	409	-	33,33,35	0.21	0	44,44,46	0.54	0
3	ABU	C	401	-	6,6,6	1.63	2 (33%)	6,6,6	1.59	2 (33%)
3	ABU	B	401	-	6,6,6	0.11	0	6,6,6	0.21	0
7	PG4	E	405	-	12,12,12	0.19	0	11,11,11	0.20	0
4	MES	B	402	-	12,12,12	1.24	2 (16%)	15,16,16	0.47	0
5	GOL	E	404	-	5,5,5	0.11	0	5,5,5	0.40	0
8	UMQ	B	406	-	33,33,35	0.21	0	44,44,46	0.53	0
7	PG4	C	404	-	12,12,12	0.22	0	11,11,11	0.17	0
3	ABU	D	401	-	6,6,6	1.59	2 (33%)	6,6,6	1.80	2 (33%)
9	ACT	D	406	-	3,3,3	1.20	0	3,3,3	1.01	0
4	MES	A	402	-	12,12,12	0.83	0	15,16,16	0.41	0
4	MES	C	402	-	12,12,12	1.08	1 (8%)	15,16,16	0.43	0
5	GOL	E	403	-	5,5,5	0.16	0	5,5,5	0.26	0
9	ACT	E	407	-	3,3,3	1.27	0	3,3,3	0.92	0
5	GOL	B	403	-	5,5,5	0.05	0	5,5,5	0.46	0
5	GOL	A	403	-	5,5,5	0.10	0	5,5,5	0.26	0
4	MES	D	402	-	12,12,12	0.83	0	15,16,16	0.26	0
5	GOL	D	403	-	5,5,5	0.20	0	5,5,5	0.14	0
8	UMQ	E	408	-	33,33,35	0.24	0	44,44,46	0.81	2 (4%)
7	PG4	D	404	-	12,12,12	0.29	0	11,11,11	0.33	0
3	ABU	A	401	-	6,6,6	1.53	2 (33%)	6,6,6	2.03	2 (33%)
3	ABU	E	401	-	6,6,6	1.67	2 (33%)	6,6,6	1.65	2 (33%)
7	PG4	B	404	-	12,12,12	0.23	0	11,11,11	0.11	0
8	UMQ	C	406	-	33,33,35	0.23	0	44,44,46	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	UMQ	B	407	-	-	6/18/58/60	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	C	403	-	-	0/4/4/4	-
4	MES	E	402	-	-	0/6/14/14	0/1/1/1
8	UMQ	E	409	-	-	6/18/58/60	0/2/2/2
3	ABU	C	401	-	-	0/4/4/4	-
3	ABU	B	401	-	-	2/4/4/4	-
7	PG4	E	405	-	-	6/10/10/10	-
4	MES	B	402	-	-	0/6/14/14	0/1/1/1
5	GOL	E	404	-	-	3/4/4/4	-
8	UMQ	B	406	-	-	6/18/58/60	0/2/2/2
7	PG4	C	404	-	-	6/10/10/10	-
3	ABU	D	401	-	-	2/4/4/4	-
4	MES	A	402	-	-	3/6/14/14	0/1/1/1
4	MES	C	402	-	-	1/6/14/14	0/1/1/1
5	GOL	E	403	-	-	0/4/4/4	-
5	GOL	B	403	-	-	2/4/4/4	-
5	GOL	A	403	-	-	0/4/4/4	-
4	MES	D	402	-	-	2/6/14/14	0/1/1/1
5	GOL	D	403	-	-	0/4/4/4	-
8	UMQ	E	408	-	-	9/18/58/60	0/2/2/2
7	PG4	D	404	-	-	5/10/10/10	-
3	ABU	A	401	-	-	0/4/4/4	-
3	ABU	E	401	-	-	2/4/4/4	-
7	PG4	B	404	-	-	4/10/10/10	-
8	UMQ	C	406	-	-	8/18/58/60	0/2/2/2

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	401	ABU	O-C	2.96	1.31	1.22
3	E	401	ABU	O-C	2.95	1.31	1.22
3	D	401	ABU	O-C	2.88	1.31	1.22
3	E	401	ABU	OXT-C	-2.84	1.21	1.30
3	A	401	ABU	O-C	2.70	1.30	1.22

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	ABU	OXT-C-CG	3.48	125.01	114.00
3	A	401	ABU	O-C-CG	-3.45	112.14	123.09

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	401	ABU	O-C-CG	-3.12	113.20	123.09
3	D	401	ABU	OXT-C-CG	3.09	123.78	114.00
3	E	401	ABU	OXT-C-CG	2.83	122.95	114.00

There are no chirality outliers.

5 of 73 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
4	A	402	MES	C7-C8-S-O1S
4	A	402	MES	C7-C8-S-O3S
5	B	403	GOL	C1-C2-C3-O3
8	B	406	UMQ	C2'-C1'-O1'-CA
8	B	407	UMQ	O5-C1-O1-C4'


There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	404	GOL	1	0
7	D	404	PG4	2	0
3	E	401	ABU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/318 (95%)	0.89	37 (12%) 8 6	48, 74, 113, 155	0
1	B	308/318 (96%)	0.95	42 (13%) 7 5	38, 71, 127, 158	1 (0%)
1	C	307/318 (96%)	0.88	43 (14%) 6 5	46, 71, 119, 143	0
1	D	305/318 (95%)	1.00	49 (16%) 4 3	27, 74, 124, 149	1 (0%)
1	E	304/318 (95%)	0.96	45 (14%) 5 4	48, 74, 112, 155	0
2	F	110/121 (90%)	1.63	40 (36%) 1 0	64, 97, 130, 138	0
2	G	111/121 (91%)	0.89	13 (11%) 9 7	50, 64, 87, 117	0
2	H	112/121 (92%)	0.55	9 (8%) 18 14	47, 62, 91, 105	0
2	I	111/121 (91%)	1.20	24 (21%) 2 2	54, 75, 116, 134	0
2	J	110/121 (90%)	1.86	45 (40%) 0 0	61, 99, 129, 138	0
All	All	2081/2195 (94%)	1.01	347 (16%) 4 3	27, 74, 123, 158	2 (0%)

The worst 5 of 347 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	181	VAL	8.8
2	H	112	HIS	6.2
1	B	319	GLY	6.1
1	D	7	VAL	5.6
1	C	7	VAL	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	ACT	E	407	4/4	0.50	0.24	81,81,81,81	0
9	ACT	D	406	4/4	0.57	0.24	75,76,76,77	0
8	UMQ	E	409	32/34	0.75	0.21	99,131,137,138	0
4	MES	C	402	12/12	0.76	0.26	88,92,111,112	0
8	UMQ	B	406	32/34	0.77	0.19	98,121,137,140	0
5	GOL	C	403	6/6	0.77	0.27	68,69,71,71	0
8	UMQ	E	408	32/34	0.78	0.18	95,120,134,135	0
8	UMQ	C	406	32/34	0.79	0.19	106,122,131,134	0
6	CA	C	405	1/1	0.80	0.13	110,110,110,110	0
7	PG4	E	405	13/13	0.80	0.27	82,83,86,86	0
4	MES	A	402	12/12	0.80	0.23	87,92,116,116	0
6	CA	A	404	1/1	0.80	0.10	107,107,107,107	0
5	GOL	E	404	6/6	0.81	0.24	89,91,91,92	0
6	CA	E	406	1/1	0.81	0.13	126,126,126,126	0
8	UMQ	B	407	32/34	0.81	0.19	100,114,130,135	0
7	PG4	D	404	13/13	0.81	0.27	81,83,86,86	0
5	GOL	B	403	6/6	0.83	0.23	73,77,78,78	0
6	CA	B	405	1/1	0.83	0.12	118,118,118,118	0
7	PG4	C	404	13/13	0.83	0.25	97,101,103,103	0
4	MES	B	402	12/12	0.84	0.20	89,92,105,106	0
7	PG4	B	404	13/13	0.84	0.23	82,84,86,87	0
3	ABU	A	401	7/7	0.86	0.17	68,74,83,84	0
4	MES	D	402	12/12	0.86	0.19	82,86,106,107	0
5	GOL	A	403	6/6	0.86	0.17	72,73,75,75	0
4	MES	E	402	12/12	0.87	0.18	79,83,104,104	0
5	GOL	D	403	6/6	0.89	0.18	74,76,79,81	0
5	GOL	E	403	6/6	0.89	0.18	73,74,74,75	0
3	ABU	B	401	7/7	0.89	0.15	64,69,77,77	0
3	ABU	E	401	7/7	0.90	0.17	63,68,76,78	0
3	ABU	C	401	7/7	0.91	0.12	58,62,71,72	0
3	ABU	D	401	7/7	0.93	0.14	68,70,77,77	0
6	CA	D	405	1/1	0.95	0.08	117,117,117,117	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around UMQ B 406:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UMQ E 408:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UMQ C 406:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UMQ B 407:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.