



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 08:26 AM UTC

PDB ID : 5STW / pdb_00005stw
Title : PanDDA analysis group deposition – Aar2/RNaseH in complex with fragment P03C12 from the F2X-Universal Library
Authors : Barthel, T.; Wollenhaupt, J.; Lima, G.M.A.; Wahl, M.C.; Weiss, M.S.
Deposited on : 2022-08-26
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	50% 75% 16% 8%
2	B	308	52% 77% 19% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	V8L	B	401	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9173 atoms, of which 4524 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	237	4068	1287	2060	336	373	12	18	21	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	300	5044	1654	2464	421	485	20	0	17	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	?	-	LEU	deletion	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	170	SER	ASP	conflict	UNP P32357

- Molecule 3 is N¹-phenylacetohydrazide (CCD ID: V8L) (formula: C₈H₁₀N₂O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			11	8	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	21	Total	O	0	0
			21	21		
4	B	29	Total	O	0	0
			29	29		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pre-mRNA-splicing factor 8



• Molecule 2: A1 cistron-splicing factor AAR2



Y288	W289	W290	I291	C292	L293	Y294	S295	S296	S301	L302	H303	H304	T305	E306	K307	I308	M309	E310	M311	K312	Y313	L316	L317
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.10Å 82.20Å 93.53Å 90.00° 108.52° 90.00°	Depositor
Resolution (Å)	44.34 – 1.89 44.34 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.2 (44.34-1.89) 99.3 (44.34-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.89Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.283 , 0.341 0.297 , 0.346	Depositor DCC
R_{free} test set	2100 reflections (4.15%)	wwPDB-VP
Wilson B-factor (Å ²)	51.6	Xtriage
Anisotropy	0.385	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 75.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9173	wwPDB-VP
Average B, all atoms (Å ²)	107.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: V8L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/2149	0.99	2/2911 (0.1%)
2	B	0.95	6/2739 (0.2%)	1.18	16/3699 (0.4%)
All	All	0.88	6/4888 (0.1%)	1.10	18/6610 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	123[A]	MET	N-CA	8.49	1.56	1.46
2	B	123[B]	MET	N-CA	8.49	1.56	1.46
2	B	122[A]	GLN	C-N	6.68	1.43	1.33
2	B	122[B]	GLN	C-N	6.68	1.43	1.33
2	B	42[A]	HIS	CB-CG	5.39	1.57	1.50
2	B	42[B]	HIS	CB-CG	5.39	1.57	1.50

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	122[A]	GLN	CA-C-N	-7.57	109.53	120.29
2	B	122[A]	GLN	C-N-CA	-7.57	109.53	120.29
2	B	122[B]	GLN	CA-C-N	-7.57	109.53	120.29
2	B	122[B]	GLN	C-N-CA	-7.57	109.53	120.29
2	B	214	PHE	O-C-N	6.76	129.28	122.12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	35	ASP	CA-C-N	-6.46	117.94	122.59
2	B	35	ASP	C-N-CA	-6.46	117.94	122.59
2	B	214	PHE	CA-C-N	6.33	133.63	121.54
2	B	214	PHE	C-N-CA	6.33	133.63	121.54
2	B	223	GLU	CA-C-N	6.15	128.52	120.28
2	B	223	GLU	C-N-CA	6.15	128.52	120.28
2	B	227	ALA	CA-C-O	-5.40	115.12	121.07
1	A	1910	LYS	CA-C-N	5.39	127.50	120.28
1	A	1910	LYS	C-N-CA	5.39	127.50	120.28
2	B	176	LEU	CA-C-O	-5.28	115.04	121.58
2	B	223	GLU	O-C-N	5.27	127.71	122.12
2	B	180	VAL	CA-C-N	5.14	129.80	123.12
2	B	180	VAL	C-N-CA	5.14	129.80	123.12

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	122[A]	GLN	Mainchain
2	B	122[B]	GLN	Mainchain
2	B	176	LEU	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2008	2060	1974	26	0
2	B	2580	2464	2396	46	0
3	B	11	0	0	0	0
4	A	21	0	0	1	0
4	B	29	0	0	3	0
All	All	4649	4524	4370	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1962:ARG:O	1:A:2013:ARG:NH1	1.97	0.97
2:B:129:ILE:HG23	2:B:177[A]:ASN:C	2.01	0.86
2:B:129:ILE:HG23	2:B:177[B]:ASN:C	2.02	0.85
2:B:287:ARG:O	2:B:291:ILE:HD13	1.95	0.67
2:B:111:ASP:OD2	4:B:501:HOH:O	2.13	0.66
2:B:226:PHE:CE1	2:B:230[A]:ASN:ND2	2.64	0.66
1:A:2049:ILE:O	1:A:2053[A]:SER:OG	2.13	0.66
1:A:2064:GLY:O	1:A:2068:ASN:N	2.31	0.64
2:B:70:GLN:HB3	2:B:81:MET:CE	2.29	0.62
2:B:70:GLN:HB3	2:B:81:MET:HE2	1.81	0.62
1:A:1911:TRP:CD1	2:B:195:MET:HE3	2.34	0.62
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	1.83	0.60
1:A:2061:THR:O	1:A:2064:GLY:N	2.36	0.57
1:A:1859:ARG:HH12	1:A:1979[A]:MET:CE	2.16	0.57
2:B:1:MET:N	4:B:502:HOH:O	2.37	0.57
2:B:130:VAL:HG12	2:B:130:VAL:O	2.03	0.56
2:B:1:MET:N	2:B:38:ILE:HD11	2.21	0.55
2:B:54[A]:MET:HE1	4:B:528:HOH:O	2.06	0.54
1:A:1993:ASP:CG	1:A:2038:HIS:HB3	2.33	0.54
2:B:17:ILE:HD13	2:B:44[B]:ILE:HG13	1.90	0.54
2:B:37:PRO:HD3	2:B:105:TYR:CD1	2.43	0.54
2:B:277:GLU:CD	2:B:277:GLU:H	2.16	0.52
2:B:1:MET:H2	2:B:38:ILE:HD11	1.76	0.51
2:B:242:GLN:O	2:B:246:MET:HG3	2.12	0.49
2:B:44[A]:ILE:O	2:B:44[A]:ILE:HG23	2.12	0.49
2:B:147:VAL:HG23	2:B:177[A]:ASN:HA	1.94	0.49
2:B:223:GLU:HG2	2:B:246:MET:SD	2.52	0.49
2:B:258:LYS:HD2	2:B:258:LYS:H	1.79	0.48
2:B:53:SER:O	2:B:54[A]:MET:HB3	2.12	0.48
2:B:190:ARG:HG3	2:B:203[B]:TYR:CZ	2.49	0.47
2:B:208:VAL:O	2:B:213:ILE:HG13	2.15	0.47
2:B:251:CYS:O	2:B:296:SER:HB2	2.14	0.47
2:B:129:ILE:HG23	2:B:177[B]:ASN:O	2.15	0.47
2:B:17:ILE:HD13	2:B:44[B]:ILE:CG1	2.45	0.47
1:A:1974:LEU:O	1:A:1977:VAL:HG12	2.15	0.47
2:B:130:VAL:HG22	2:B:176:LEU:HD23	1.95	0.46
1:A:1914:ALA:HB1	1:A:1944:LEU:HA	1.97	0.46
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ2	1.64	0.46
2:B:134:GLU:OE1	2:B:134:GLU:N	2.36	0.46
1:A:2060:LEU:O	1:A:2063:TYR:HB3	2.16	0.45
1:A:1853:ASP:OD2	1:A:1855[B]:THR:HG23	2.16	0.45
2:B:130:VAL:HG11	2:B:138:SER:HB3	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1878:CYS:SG	1:A:1969:MET:HE3	2.56	0.45
2:B:28:GLN:HG3	2:B:29:PRO:HD2	1.98	0.45
2:B:147:VAL:HG23	2:B:177[B]:ASN:HA	1.99	0.45
2:B:141:ASP:OD1	2:B:144:MET:HG3	2.17	0.44
2:B:114:TRP:CE2	2:B:118:THR:HG21	2.52	0.44
1:A:1948:MET:HE3	1:A:1948:MET:HB3	1.88	0.44
2:B:187:GLU:H	2:B:187:GLU:CD	2.24	0.43
2:B:277:GLU:CD	2:B:277:GLU:N	2.75	0.43
1:A:1940:MET:C	1:A:1943:PRO:HD2	2.43	0.43
1:A:1996:LEU:HB2	4:A:2108:HOH:O	2.18	0.43
1:A:1881:THR:O	1:A:1889:LEU:HD12	2.18	0.43
2:B:68:TYR:CE2	2:B:81:MET:HE3	2.54	0.43
2:B:129:ILE:HG23	2:B:177[A]:ASN:CA	2.48	0.43
1:A:1859:ARG:HB2	1:A:1875:ILE:HG13	2.01	0.42
1:A:2011:LEU:HD23	1:A:2055:MET:HE3	2.02	0.42
2:B:206:ASN:O	2:B:210:LEU:HB2	2.20	0.41
2:B:147:VAL:CG2	2:B:177[A]:ASN:HA	2.51	0.41
2:B:228:PHE:HB2	2:B:243:TRP:CD1	2.56	0.41
1:A:1859:ARG:NH1	1:A:1979[A]:MET:SD	2.90	0.41
2:B:31:HIS:HB3	2:B:96:PHE:CD2	2.56	0.41
2:B:306:GLU:O	2:B:310:GLU:HG3	2.20	0.41
1:A:1902:GLN:HB2	1:A:1905:LEU:CD2	2.51	0.41
2:B:105:TYR:HA	2:B:106:PRO:HD3	1.90	0.41
2:B:179:THR:HG21	2:B:214:PHE:CZ	2.56	0.41
1:A:1933:ILE:HB	1:A:1956:ILE:HD12	2.03	0.40
1:A:2061:THR:O	1:A:2063:TYR:N	2.54	0.40
1:A:1865:THR:HG23	1:A:1869:ASN:O	2.21	0.40
1:A:1835:MET:HE3	1:A:1835:MET:HB3	1.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/258 (100%)	247 (96%)	9 (4%)	2 (1%)	16	8
2	B	315/308 (102%)	290 (92%)	22 (7%)	3 (1%)	12	5
All	All	573/566 (101%)	537 (94%)	31 (5%)	5 (1%)	16	6

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1896	THR
2	B	177[A]	ASN
2	B	177[B]	ASN
1	A	2061	THR
2	B	215	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/233 (102%)	235 (99%)	2 (1%)	73	75
2	B	294/284 (104%)	290 (99%)	4 (1%)	59	59
All	All	531/517 (103%)	525 (99%)	6 (1%)	63	67

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1835	MET
1	A	1962	ARG
2	B	77	LEU
2	B	124	ASP
2	B	281	ASP
2	B	317	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	150	ASN
2	B	259	HIS
2	B	290	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	V8L	B	401	-	11,11,11	2.36	5 (45%)	13,13,13	4.23	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	V8L	B	401	-	-	4/5/5/5	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	V8L	C6-C7	-3.97	1.32	1.38
3	B	401	V8L	O-C1	-3.72	1.14	1.23
3	B	401	V8L	C3-C2	-2.96	1.34	1.39
3	B	401	V8L	C4-C3	-2.65	1.34	1.38
3	B	401	V8L	N1-N	-2.55	1.32	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	V8L	C-C1-N	10.66	124.52	114.88
3	B	401	V8L	O-C1-N	-7.93	113.95	122.04
3	B	401	V8L	C1-N-N1	5.70	131.01	119.72
3	B	401	V8L	C2-N1-N	-4.09	105.83	116.66

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	401	V8L	C7-C2-N1-N
3	B	401	V8L	C3-C2-N1-N
3	B	401	V8L	O-C1-N-N1
3	B	401	V8L	C-C1-N-N1

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/258 (91%)	2.35	128 (54%) 0 0	32, 101, 167, 222	12 (5%)
2	B	300/308 (97%)	2.30	160 (53%) 0 0	28, 98, 165, 287	9 (3%)
All	All	537/566 (94%)	2.32	288 (53%) 0 0	28, 100, 166, 287	21 (3%)

All (288) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	129	ILE	8.9
1	A	2039	LEU	7.7
1	A	2040	TRP	7.5
1	A	2069	VAL	7.0
1	A	1878	CYS	6.9
1	A	2017[A]	THR	6.6
2	B	77	LEU	6.3
2	B	226	PHE	6.0
2	B	288	VAL	5.8
1	A	2063	TYR	5.7
1	A	2048	TRP	5.7
2	B	213	ILE	5.4
2	B	111	ASP	5.4
2	B	30	PHE	5.3
1	A	2034	ILE	5.3
1	A	1958	PRO	4.9
2	B	126	ILE	4.8
1	A	1919[A]	ALA	4.8
1	A	1963	LEU	4.8
1	A	1896	THR	4.8
2	B	313	TYR	4.7
2	B	49	ALA	4.7
1	A	1862	VAL	4.6
2	B	293	LEU	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	218	SER	4.6
1	A	1979[A]	MET	4.5
1	A	2059	ILE	4.5
2	B	36	ILE	4.5
2	B	214	PHE	4.4
2	B	152	LEU	4.4
2	B	120	PHE	4.4
2	B	170	SER	4.4
2	B	282	ILE	4.3
2	B	256	VAL	4.3
1	A	2051	ILE	4.3
1	A	1851	PHE	4.2
2	B	1	MET	4.2
1	A	2038	HIS	4.2
1	A	2027	LEU	4.2
2	B	93	VAL	4.2
2	B	289	TRP	4.2
2	B	221	PHE	4.2
2	B	121	VAL	4.2
1	A	2037	TYR	4.1
2	B	220	TYR	4.1
1	A	2026	LEU	4.1
1	A	2060	LEU	4.1
2	B	172	PRO	4.1
2	B	124	ASP	4.1
1	A	1920	LEU	4.1
2	B	316	LEU	4.1
1	A	1995	TRP	4.0
2	B	56	TYR	4.0
2	B	181	ILE	4.0
1	A	1840	TYR	4.0
2	B	146	THR	3.9
1	A	1893	ILE	3.9
2	B	20	TYR	3.9
1	A	2049	ILE	3.9
2	B	183	PHE	3.8
1	A	1931[A]	LYS	3.8
2	B	176	LEU	3.8
2	B	96	PHE	3.8
1	A	2002[A]	TYR	3.8
1	A	1898[A]	VAL	3.8
2	B	130	VAL	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	275	LEU	3.7
2	B	15	ILE	3.7
1	A	2068	ASN	3.7
2	B	142	SER	3.6
2	B	54[A]	MET	3.6
2	B	272	ILE	3.6
2	B	246	MET	3.6
2	B	270	TYR	3.6
1	A	1857	VAL	3.6
1	A	2061	THR	3.6
2	B	11	ILE	3.6
2	B	137	PHE	3.6
2	B	147	VAL	3.5
1	A	2043	PHE	3.5
2	B	206	ASN	3.5
1	A	1933	ILE	3.5
1	A	1870	VAL	3.5
2	B	46	PHE	3.5
1	A	1951	PHE	3.5
1	A	1834	ALA	3.5
2	B	244	HIS	3.5
2	B	179	THR	3.4
2	B	308	ILE	3.4
1	A	1860	VAL	3.4
1	A	1874	ALA	3.4
2	B	217	SER	3.4
1	A	2025	ILE	3.4
1	A	1930	PRO	3.4
1	A	1989	PHE	3.3
1	A	1964	PRO	3.3
1	A	2036	SER	3.3
2	B	241	LEU	3.3
2	B	268	LEU	3.3
2	B	101	MET	3.3
2	B	108	ILE	3.3
2	B	128	LYS	3.3
1	A	1944	LEU	3.3
1	A	1996	LEU	3.3
2	B	22	PHE	3.3
2	B	203[A]	TYR	3.2
2	B	87	ALA	3.2
2	B	140	VAL	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	125	LYS	3.2
2	B	117	LEU	3.2
1	A	1833	GLY	3.2
2	B	294	TYR	3.2
1	A	1987	VAL	3.2
2	B	103	VAL	3.2
2	B	122[A]	GLN	3.2
2	B	295	SER	3.2
1	A	1955	ALA	3.1
2	B	123[A]	MET	3.1
2	B	13	VAL	3.1
2	B	229	LEU	3.1
2	B	134	GLU	3.1
1	A	1981	ALA	3.1
1	A	1943	PRO	3.1
2	B	5	PRO	3.1
1	A	1965	PHE	3.1
2	B	276	PRO	3.1
1	A	2058	LEU	3.1
2	B	178[A]	TYR	3.1
2	B	279	TYR	3.1
2	B	189	ILE	3.0
1	A	2008	LEU	3.0
2	B	8	SER	3.0
1	A	1936	THR	3.0
2	B	7	THR	3.0
1	A	1941	LEU	3.0
1	A	2010	LEU	3.0
1	A	2013	ARG	3.0
2	B	27	ASN	3.0
2	B	292	CYS	3.0
1	A	2066	LYS	3.0
1	A	2055	MET	3.0
1	A	2067	TYR	2.9
1	A	2065	ARG	2.9
1	A	1866	PHE	2.9
2	B	114	TRP	2.9
2	B	311	ASN	2.9
1	A	1967	ALA	2.9
1	A	1850	LEU	2.9
2	B	249	LEU	2.9
1	A	1998	ARG	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	4	VAL	2.9
2	B	24	VAL	2.9
1	A	1986	MET	2.9
2	B	317	LEU	2.9
1	A	1914	ALA	2.9
1	A	1849	LYS	2.9
1	A	1897	SER	2.9
1	A	1988	LEU	2.9
1	A	1966	SER	2.8
1	A	1970	SER	2.8
2	B	302	LEU	2.8
1	A	2064	GLY	2.8
1	A	1935	VAL	2.8
2	B	231	ALA	2.8
2	B	254	ALA	2.8
1	A	1863	HIS	2.8
1	A	1875	ILE	2.8
1	A	2032	ILE	2.8
2	B	224	LEU	2.8
1	A	2044	THR	2.8
2	B	75	ASP	2.8
2	B	235	GLY	2.8
1	A	1917	VAL	2.8
2	B	139	TYR	2.7
1	A	1961[A]	LEU	2.7
2	B	17	ILE	2.7
2	B	281	ASP	2.7
2	B	127	ARG	2.7
2	B	243	TRP	2.7
1	A	1901	GLY	2.7
2	B	33	ILE	2.7
1	A	1846	ASN	2.7
1	A	1873	LYS	2.7
1	A	1980[A]	LYS	2.7
1	A	2005	PHE	2.7
2	B	173	ALA	2.6
2	B	192	GLY	2.6
1	A	2030	PRO	2.6
2	B	92	ILE	2.6
2	B	62	CYS	2.6
1	A	1890	PHE	2.6
2	B	198	PHE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1949	LEU	2.6
1	A	1974	LEU	2.6
1	A	2012	LEU	2.6
2	B	210	LEU	2.6
2	B	180	VAL	2.6
1	A	1956	ILE	2.6
1	A	1841	ALA	2.6
1	A	1909	ALA	2.6
2	B	216	ASN	2.6
2	B	305	THR	2.6
1	A	2041	PRO	2.5
1	A	1929	GLN	2.5
2	B	67	PHE	2.5
2	B	233	PHE	2.5
1	A	1894	ILE	2.5
2	B	99	ARG	2.5
1	A	1985	GLN	2.5
2	B	69	ILE	2.5
1	A	1886	THR	2.5
2	B	6	PHE	2.5
2	B	283	LEU	2.5
1	A	1879	ILE	2.5
2	B	106	PRO	2.5
2	B	204[A]	TYR	2.4
2	B	230[A]	ASN	2.4
1	A	1844	PHE	2.4
1	A	2015	LEU	2.4
2	B	94	HIS	2.4
2	B	287	ARG	2.4
2	B	285	ASN	2.4
2	B	42[A]	HIS	2.4
2	B	21	SER	2.4
2	B	64[A]	MET	2.4
2	B	234	PHE	2.4
1	A	2056[A]	ARG	2.4
2	B	71	TYR	2.4
2	B	50	ASP	2.4
1	A	2062	GLU	2.4
1	A	1924	LEU	2.3
2	B	199	LEU	2.3
2	B	237	TYR	2.3
2	B	193	HIS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1910	LYS	2.3
2	B	136	GLN	2.3
2	B	284	LEU	2.3
2	B	89	PHE	2.3
2	B	53	SER	2.3
2	B	227	ALA	2.3
1	A	1937	ARG	2.3
1	A	1954	ILE	2.3
2	B	247	ILE	2.3
2	B	59	TRP	2.2
2	B	73	PRO	2.2
2	B	115	TYR	2.2
2	B	258	LYS	2.2
1	A	2009	THR	2.2
1	A	1939	ALA	2.2
2	B	232	MET	2.2
1	A	1977	VAL	2.2
1	A	2046	GLU	2.2
1	A	2052	GLU	2.2
1	A	1843	LEU	2.2
1	A	2024[A]	MET	2.2
1	A	1982	THR	2.2
2	B	301	SER	2.2
1	A	1984	PRO	2.2
2	B	10	PRO	2.2
2	B	74	LYS	2.2
2	B	219	ASN	2.1
2	B	38	ILE	2.1
2	B	267	ILE	2.1
1	A	1854	ASP	2.1
2	B	307	LYS	2.1
2	B	209	MET	2.1
1	A	1971	ILE	2.1
2	B	250	ILE	2.1
2	B	118	THR	2.1
2	B	175	SER	2.1
2	B	274	THR	2.1
1	A	1859	ARG	2.1
1	A	1962	ARG	2.1
2	B	143	SER	2.1
2	B	239	SER	2.1
2	B	107	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2004	ALA	2.1
2	B	109	ASP	2.1
1	A	1861	THR	2.0
2	B	174	HIS	2.0
1	A	1907	GLN	2.0
2	B	304	ASN	2.0
1	A	1946	VAL	2.0
2	B	171	ASP	2.0
1	A	1835	MET	2.0
1	A	1969	MET	2.0
2	B	222	GLY	2.0
2	B	257	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	V8L	B	401	11/11	0.73	0.15	20,20,20,20	11

6.5 Other polymers [i](#)

There are no such residues in this entry.