



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:26 AM UTC

PDB ID : 6STW / pdb_00006stw
Title : Adenovirus 15 Fiber Knob protein
Authors : Mundy, R.M.; Baker, A.T.; Rizkallah, P.J.; Parker, A.L.
Deposited on : 2019-09-11
Resolution : 1.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4403 (1.40-1.36)
Clashscore	190562	4528 (1.40-1.36)
Ramachandran outliers	187476	4459 (1.40-1.36)
Sidechain outliers	187428	4458 (1.40-1.36)
RSRZ outliers	180081	4399 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	9% 85% 12% .
1	B	190	6% 86% 12% ..
1	C	190	13% 81% 16% ...

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5429 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fiber protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	190	Total	C	N	O	S	0	9	0
			1575	1002	254	313	6			
1	B	190	Total	C	N	O	S	0	11	0
			1594	1011	255	322	6			
1	C	189	Total	C	N	O	S	0	9	0
			1565	995	252	312	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	179	LYS	MET	conflict	UNP P36847
A	180	LEU	ARG	conflict	UNP P36847
B	179	LYS	MET	conflict	UNP P36847
B	180	LEU	ARG	conflict	UNP P36847
C	179	LYS	MET	conflict	UNP P36847
C	180	LEU	ARG	conflict	UNP P36847


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	231	Total	O	0	0
			231	231		
2	B	278	Total	O	0	0
			278	278		
2	C	186	Total	O	0	0
			186	186		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fiber protein



- Molecule 1: Fiber protein



- Molecule 1: Fiber protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.58Å 89.18Å 106.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.64 – 1.37 39.64 – 1.37	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.64-1.37) 99.9 (39.64-1.37)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.43 (at 1.37Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.173 , 0.199 0.179 , 0.204	Depositor DCC
R_{free} test set	5905 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5429	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	0/1608	1.07	3/2176 (0.1%)
1	B	0.96	0/1627	1.10	1/2203 (0.0%)
1	C	0.94	0/1598	1.09	1/2165 (0.0%)
All	All	0.94	0/4833	1.09	5/6544 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	346	TYR	CB-CA-C	7.14	120.60	109.03
1	C	299	ASN	CA-CB-CG	6.44	119.04	112.60
1	A	344	LYS	CB-CG-CD	5.64	124.27	111.30
1	A	346	TYR	N-CA-C	-5.27	107.40	113.88
1	B	299	ASN	CA-CB-CG	5.25	117.86	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1575	0	1559	30	0
1	B	1594	0	1557	23	0
1	C	1565	0	1543	33	0
2	A	231	0	0	3	0
2	B	278	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	186	0	0	4	0
All	All	5429	0	4659	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:360[B]:ASN:ND2	2:C:401:HOH:O	1.80	1.15
1:A:204:LEU:HD13	1:A:217[A]:VAL:HG22	1.39	1.04
1:A:181:THR:HG23	1:A:205:ILE:HD11	1.38	1.03
1:C:204:LEU:HD13	1:C:217[A]:VAL:HG22	1.40	1.00
1:C:296:LYS:HB2	1:C:297:PRO:CD	2.02	0.89
1:C:296:LYS:HB2	1:C:297:PRO:HD3	1.57	0.86
1:A:204:LEU:CD1	1:A:217[A]:VAL:HG22	2.04	0.86
1:B:184[A]:THR:HG21	1:B:203:THR:HA	1.57	0.86
1:C:204:LEU:CD1	1:C:217[A]:VAL:HG22	2.08	0.84
1:B:231:ASN:HD21	1:B:345:TRP:HE1	1.26	0.84
1:A:181:THR:HG23	1:A:205:ILE:CD1	2.07	0.82
1:C:217[A]:VAL:HG23	1:C:361:PHE:HE1	1.45	0.81
1:A:299:ASN:HD22	1:A:301:GLU:H	1.24	0.81
1:C:231:ASN:HD21	1:C:345:TRP:HE1	1.26	0.81
1:A:232[B]:THR:HG22	2:A:452:HOH:O	1.81	0.81
1:A:231:ASN:HD21	1:A:345:TRP:HE1	1.26	0.81
1:A:217[A]:VAL:HG23	1:A:361:PHE:HE1	1.47	0.80
1:A:217[A]:VAL:HG21	1:A:341:PHE:CE1	2.19	0.78
1:B:196:GLU:HG2	2:B:405:HOH:O	1.84	0.77
1:C:217[A]:VAL:HG21	1:C:341:PHE:CE1	2.19	0.77
1:C:187:ASP:OD1	2:C:402:HOH:O	2.06	0.73
1:A:256:GLN:HE21	1:A:256:GLN:HA	1.53	0.73
1:B:348[A]:THR:HG22	2:B:455:HOH:O	1.90	0.71
1:C:295:THR:HG22	1:C:296:LYS:HG3	1.76	0.67
1:C:204:LEU:HD13	1:C:217[A]:VAL:CG2	2.24	0.65
1:A:232[B]:THR:HG21	1:A:350:GLU:OE2	1.97	0.65
1:A:181:THR:CG2	1:A:205:ILE:HD11	2.21	0.63
1:C:296:LYS:CB	1:C:297:PRO:CD	2.77	0.62
1:C:299:ASN:ND2	1:C:301:GLU:H	1.99	0.61
1:B:348[A]:THR:HG23	2:B:503:HOH:O	2.01	0.61
1:A:204:LEU:HD13	1:A:217[A]:VAL:CG2	2.23	0.60
1:A:217[A]:VAL:HG23	1:A:361:PHE:CE1	2.35	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:ASN:ND2	1:A:301:GLU:H	1.98	0.59
1:B:184[A]:THR:HG22	1:B:185:THR:O	2.03	0.59
1:B:299:ASN:ND2	1:B:301:GLU:H	1.99	0.59
1:B:348[A]:THR:HG21	2:B:620:HOH:O	2.04	0.58
1:A:183:TRP:HB3	1:A:205:ILE:HD13	1.86	0.57
1:C:299:ASN:HD21	1:C:301:GLU:HG2	1.69	0.57
1:C:217[A]:VAL:HG23	1:C:361:PHE:CE1	2.33	0.55
1:C:267:TYR:CE2	1:C:277:PRO:HB3	2.43	0.54
1:B:192[A]:CYS:SG	1:B:259:THR:HG22	2.49	0.53
1:A:293:LYS:NZ	2:A:401:HOH:O	2.41	0.53
1:B:290:ALA:HB2	1:C:186:PRO:HB3	1.90	0.53
1:A:344:LYS:HE3	1:A:346:TYR:CE2	2.45	0.52
1:A:181:THR:CG2	1:A:205:ILE:CD1	2.85	0.52
1:B:178:ASP:C	1:B:179:LYS:HD2	2.35	0.52
1:C:233[A]:THR:HG21	2:C:537:HOH:O	2.09	0.52
1:C:277:PRO:O	1:C:278:TYR:CD1	2.64	0.51
1:A:245[A]:LYS:HE3	2:A:410:HOH:O	2.10	0.51
1:C:307:ASN:HD21	1:C:308:LYS:NZ	2.09	0.50
1:B:307:ASN:HD21	1:B:308:LYS:NZ	2.10	0.50
1:C:289[A]:THR:HG21	2:C:508:HOH:O	2.13	0.49
1:B:307:ASN:HD22	1:B:307:ASN:H	1.60	0.49
1:A:307:ASN:HD21	1:A:308:LYS:NZ	2.11	0.48
1:B:184[A]:THR:CG2	1:B:185:THR:O	2.61	0.48
1:B:192[A]:CYS:HB3	1:B:200:SER:OG	2.13	0.48
1:C:307:ASN:HD22	1:C:307:ASN:H	1.59	0.48
1:B:205[A]:ILE:CG2	1:B:216[A]:SER:OG	2.61	0.48
1:A:205:ILE:HD12	1:A:206:LEU:N	2.29	0.48
1:B:205[A]:ILE:HG22	1:B:216[A]:SER:OG	2.13	0.48
1:A:205:ILE:CG2	1:A:216[A]:SER:OG	2.62	0.47
1:A:205:ILE:HG22	1:A:216[A]:SER:OG	2.15	0.47
1:A:186:PRO:HB3	1:C:290:ALA:HB2	1.97	0.46
1:B:299:ASN:HD22	1:B:299:ASN:C	2.25	0.45
1:A:294:GLN:HA	1:A:294:GLN:OE1	2.18	0.44
1:B:296:LYS:HB3	1:B:297:PRO:HD3	1.99	0.44
1:C:296:LYS:HB2	1:C:297:PRO:HD2	1.94	0.44
1:B:299:ASN:HD21	1:B:301:GLU:HB2	1.83	0.44
1:C:217[A]:VAL:HG21	1:C:341:PHE:CZ	2.53	0.43
1:C:230:ASN:CG	1:C:233[A]:THR:HG22	2.43	0.43
1:A:296:LYS:N	1:A:297:PRO:CD	2.82	0.42
1:A:217[A]:VAL:HG21	1:A:341:PHE:CZ	2.53	0.42
1:B:221:VAL:HG11	1:B:226:PHE:HB2	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:178:ASP:O	1:B:179:LYS:HD2	2.20	0.42
1:C:299:ASN:HD22	1:C:299:ASN:C	2.26	0.42
1:C:228:ASN:ND2	1:C:351:ASN:HA	2.34	0.42
1:C:180:LEU:HD21	1:C:280:LYS:HD2	2.02	0.42
1:B:299:ASN:HD22	1:B:300:LYS:N	2.17	0.41
1:C:299:ASN:HD22	1:C:300:LYS:N	2.19	0.41
1:C:230:ASN:HB3	1:C:233[A]:THR:CG2	2.50	0.41
1:C:190:PRO:HD3	1:C:201:LYS:HE3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/190 (104%)	182 (92%)	15 (8%)	0	100	100
1	B	199/190 (105%)	190 (96%)	9 (4%)	0	100	100
1	C	196/190 (103%)	185 (94%)	9 (5%)	2 (1%)	12	2
All	All	592/570 (104%)	557 (94%)	33 (6%)	2 (0%)	30	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	296	LYS
1	C	277	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/175 (105%)	176 (96%)	8 (4%)	26	3
1	B	186/175 (106%)	179 (96%)	7 (4%)	29	5
1	C	183/175 (105%)	172 (94%)	11 (6%)	17	1
All	All	553/525 (105%)	527 (95%)	26 (5%)	24	3

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	205	ILE
1	A	214	LEU
1	A	256	GLN
1	A	269	SER
1	A	296	LYS
1	A	298	THR
1	A	299	ASN
1	A	307	ASN
1	B	184[A]	THR
1	B	184[B]	THR
1	B	268	ARG
1	B	295	THR
1	B	299	ASN
1	B	301	GLU
1	B	307	ASN
1	C	197	ASP
1	C	214	LEU
1	C	254	LEU
1	C	270	ASP
1	C	273	ASN
1	C	296	LYS
1	C	299	ASN
1	C	300	LYS
1	C	307	ASN
1	C	348[A]	THR
1	C	348[B]	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	212	GLN
1	A	231	ASN
1	A	234	ASN
1	A	256	GLN
1	A	299	ASN
1	A	307	ASN
1	A	351	ASN
1	B	231	ASN
1	B	234	ASN
1	B	251	ASN
1	B	299	ASN
1	B	307	ASN
1	B	351	ASN
1	C	228	ASN
1	C	231	ASN
1	C	251	ASN
1	C	273	ASN
1	C	299	ASN
1	C	304	GLN
1	C	307	ASN
1	C	351	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	190/190 (100%)	0.50	17 (8%)	15	17	9, 25, 64, 108	9 (4%)
1	B	190/190 (100%)	0.38	11 (5%)	29	33	9, 23, 41, 72	11 (5%)
1	C	189/190 (99%)	0.77	24 (12%)	8	10	9, 27, 79, 136	9 (4%)
All	All	569/570 (99%)	0.55	52 (9%)	15	17	9, 25, 62, 136	29 (5%)

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	278	TYR	6.6
1	C	274	LEU	6.1
1	A	274	LEU	5.4
1	C	277	PRO	5.0
1	C	297	PRO	4.3
1	C	275	SER	4.1
1	C	295	THR	4.1
1	B	295	THR	3.9
1	B	296	LYS	3.6
1	C	179	LYS	3.6
1	C	276	GLN	3.4
1	A	297	PRO	3.4
1	C	298	THR	3.3
1	A	272	SER	3.3
1	C	279	LYS	3.3
1	A	275	SER	3.1
1	A	269	SER	3.1
1	C	296	LYS	3.0
1	C	180	LEU	3.0
1	A	295	THR	2.9
1	C	266	ASN	2.9
1	A	273	ASN	2.8
1	C	232	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	178	ASP	2.8
1	A	232[A]	THR	2.7
1	C	254	LEU	2.7
1	A	205	ILE	2.7
1	B	271	ASN	2.7
1	C	267	TYR	2.6
1	C	256[A]	GLN	2.6
1	A	296	LYS	2.6
1	B	297	PRO	2.6
1	B	302	ILE	2.6
1	C	269	SER	2.4
1	A	270	ASP	2.4
1	B	346	TYR	2.4
1	A	294	GLN	2.4
1	B	272	SER	2.3
1	B	301	GLU	2.3
1	A	271	ASN	2.3
1	C	273	ASN	2.3
1	A	298	THR	2.2
1	C	272	SER	2.2
1	C	367	GLU	2.1
1	B	300	LYS	2.1
1	B	256	GLN	2.1
1	B	299	ASN	2.1
1	C	217[A]	VAL	2.1
1	C	280	LYS	2.1
1	A	302	ILE	2.1
1	A	276	GLN	2.0
1	C	334	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.