



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:36 PM UTC

PDB ID : 1SUO / pdb_00001suo
Title : Structure of mammalian cytochrome P450 2B4 with bound 4-(4-chlorophenyl)imidazole
Authors : Scott, E.E.; White, M.A.; He, Y.A.; Johnson, E.F.; Stout, C.D.; Halpert, J.R.
Deposited on : 2004-03-26
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	16% 81% 15% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 2B4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	465	Total	C	N	O	S	0	0	0
			3734	2405	649	669	11			

There are 33 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	21	ALA	GLU	engineered mutation	UNP P00178
A	?	-	PHE	deletion	UNP P00178
A	?	-	SER	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	ALA	deletion	UNP P00178
A	?	-	PHE	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	ALA	deletion	UNP P00178
A	?	-	GLY	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	PHE	deletion	UNP P00178
A	?	-	ARG	deletion	UNP P00178
A	22	LYS	GLY	engineered mutation	UNP P00178
A	23	LYS	HIS	engineered mutation	UNP P00178
A	24	THR	PRO	engineered mutation	UNP P00178
A	25	SER	LYS	engineered mutation	UNP P00178
A	26	SER	ALA	engineered mutation	UNP P00178
A	27	LYS	HIS	engineered mutation	UNP P00178
A	29	LYS	ARG	engineered mutation	UNP P00178

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	221	SER	PRO	SEE REMARK 999	UNP P00178
A	226	TYR	HIS	engineered mutation	UNP P00178
A	492	HIS	-	expression tag	UNP P00178
A	493	HIS	-	expression tag	UNP P00178
A	494	HIS	-	expression tag	UNP P00178
A	495	HIS	-	expression tag	UNP P00178

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $\text{C}_{34}\text{H}_{32}\text{FeN}_4\text{O}_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is 4-(4-CHLOROPHENYL)IMIDAZOLE (CCD ID: CPZ) (formula: $C_9H_7ClN_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	0	0
			12	9	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	190	Total	O	0	0
			190	190		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytochrome P450 2B4



4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	233.39Å 233.39Å 56.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.00 – 1.90 49.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.00-1.90) 99.8 (49.00-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.79Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.215 , 0.239 (Not available) , 0.254	Depositor DCC
R_{free} test set	4155 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	43.2	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 43.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3979	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CPZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	4/3828 (0.1%)	1.13	21/5181 (0.4%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	351	ALA	CA-CB	8.97	1.67	1.53
1	A	314	MET	SD-CE	8.75	2.01	1.79
1	A	404	THR	CA-CB	6.98	1.58	1.52
1	A	356	ILE	CA-C	5.04	1.59	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	363	ILE	CA-C-N	10.70	130.37	119.24
1	A	363	ILE	C-N-CA	10.70	130.37	119.24
1	A	367	VAL	N-CA-C	-7.34	99.89	107.89
1	A	62	TYR	N-CA-C	6.85	121.81	113.17
1	A	102	ALA	N-CA-C	6.63	119.35	111.33
1	A	301	GLU	N-CA-C	6.61	118.37	111.03
1	A	285	HIS	N-CA-C	6.51	118.92	111.11
1	A	374	ASP	N-CA-C	-6.48	100.44	110.10
1	A	337	PRO	N-CA-C	6.24	117.25	110.58
1	A	114	ILE	CB-CA-C	-5.91	104.28	112.02
1	A	71	GLY	N-CA-C	-5.80	101.49	111.47
1	A	138	GLY	N-CA-C	-5.77	106.75	114.25
1	A	283	PHE	N-CA-C	5.54	118.21	109.39
1	A	361	ASP	N-CA-CB	-5.50	102.97	111.65
1	A	216	VAL	N-CA-C	-5.41	105.26	110.72
1	A	112	GLY	N-CA-C	-5.30	106.36	112.29
1	A	291	THR	N-CA-C	-5.26	105.22	111.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	289	ILE	CB-CA-C	-5.09	105.45	111.97
1	A	362	LEU	N-CA-C	5.03	118.67	112.54
1	A	209	ILE	CA-C-N	-5.02	114.68	122.67
1	A	209	ILE	C-N-CA	-5.02	114.68	122.67

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3734	0	3744	51	0
2	A	43	0	30	3	0
3	A	12	0	7	0	0
4	A	190	0	0	1	0
All	All	3979	0	3781	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:ARG:HH11	1:A:53:ARG:HA	1.38	0.87
1:A:95:PHE:O	1:A:369:HIS:HD2	1.69	0.76
1:A:53:ARG:HH11	1:A:53:ARG:CA	2.00	0.73
1:A:129:LEU:O	1:A:133:ARG:HB3	1.92	0.68
1:A:369:HIS:HE1	2:A:500:HEM:O2A	1.77	0.68
1:A:133:ARG:HG2	1:A:134:ASP:OD2	1.96	0.65
1:A:69:TYR:CE2	1:A:74:PRO:HB3	2.33	0.63
1:A:49:LYS:HE2	1:A:53:ARG:HG2	1.81	0.60
1:A:202:PHE:CE2	1:A:241:ILE:HD13	2.37	0.59
1:A:326:LYS:O	1:A:330:GLN:HG3	2.02	0.59
1:A:28:GLY:HA3	1:A:381:VAL:CG2	2.33	0.58
1:A:98:ARG:HG2	1:A:115:PHE:HA	1.86	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:MET:HG3	1:A:138:GLY:N	2.19	0.57
1:A:133:ARG:O	1:A:133:ARG:HG3	2.04	0.57
1:A:383:PRO:HG2	1:A:386:THR:OG1	2.05	0.57
1:A:140:ARG:HH12	1:A:148:GLU:HG2	1.69	0.56
1:A:53:ARG:HH11	1:A:53:ARG:CB	2.18	0.55
1:A:120:ARG:HA	1:A:282:GLU:HG3	1.89	0.54
1:A:168:THR:HA	1:A:308:ARG:HD2	1.90	0.54
1:A:53:ARG:NH1	1:A:53:ARG:HB2	2.24	0.53
1:A:135:PHE:CD2	1:A:139:LYS:HG3	2.45	0.52
1:A:133:ARG:HD2	4:A:1161:HOH:O	2.10	0.51
1:A:135:PHE:CE2	1:A:139:LYS:HG3	2.46	0.51
1:A:34:PRO:HD3	1:A:62:TYR:CE2	2.47	0.50
1:A:28:GLY:HA3	1:A:381:VAL:HG21	1.94	0.49
1:A:206:PHE:CE2	1:A:301:GLU:HB2	2.48	0.49
1:A:404:THR:HG23	1:A:404:THR:O	2.11	0.49
1:A:244:PHE:O	1:A:247:GLN:HG2	2.12	0.48
1:A:272:MET:HG2	1:A:283:PHE:O	2.13	0.48
1:A:49:LYS:CE	1:A:53:ARG:HG2	2.45	0.47
1:A:202:PHE:CZ	1:A:296:PHE:HB3	2.50	0.47
1:A:296:PHE:O	1:A:300:THR:HB	2.14	0.47
1:A:32:PRO:HD3	1:A:380:TYR:CZ	2.50	0.47
1:A:430:SER:CB	2:A:500:HEM:HBA1	2.46	0.46
1:A:435:ILE:O	1:A:436:CYS:C	2.59	0.46
1:A:86:GLU:O	1:A:90:ASP:HB2	2.16	0.45
1:A:430:SER:HB3	2:A:500:HEM:HBA1	2.00	0.44
1:A:31:PRO:HA	1:A:380:TYR:CE1	2.53	0.43
1:A:53:ARG:CB	1:A:53:ARG:NH1	2.80	0.43
1:A:95:PHE:O	1:A:369:HIS:CD2	2.59	0.42
1:A:47:ASP:HB2	1:A:57:ARG:HD3	2.00	0.42
1:A:206:PHE:CZ	1:A:301:GLU:HB2	2.55	0.42
1:A:105:ASP:N	1:A:106:PRO:CD	2.84	0.41
1:A:224:LEU:HA	1:A:227:PHE:HD2	1.86	0.41
1:A:165:LEU:C	1:A:165:LEU:HD12	2.46	0.41
1:A:28:GLY:O	1:A:29:LYS:HG2	2.20	0.41
1:A:32:PRO:HD3	1:A:380:TYR:CE1	2.56	0.41
1:A:309:TYR:CE2	1:A:313:LEU:HD11	2.57	0.40
1:A:135:PHE:O	1:A:136:GLY:O	2.40	0.40
1:A:360:GLY:O	1:A:361:ASP:C	2.64	0.40
1:A:403:GLU:C	1:A:404:THR:HG22	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/476 (97%)	446 (96%)	16 (4%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	136	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/421 (98%)	405 (98%)	6 (2%)	57	56

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	ARG
1	A	137	MET
1	A	186	LYS
1	A	300	THR
1	A	301	GLU
1	A	461	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	204	GLN
1	A	233	GLN
1	A	237	ASN
1	A	260	ASN
1	A	335	HIS
1	A	369	HIS
1	A	456	ASN
1	A	485	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	500	1,3	50,50,50	2.52	23 (46%)	67,82,82	1.31	7 (10%)
3	CPZ	A	600	2	12,13,13	2.06	3 (25%)	16,17,17	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1,3	-	3/14/54/54	-
3	CPZ	A	600	2	-	0/4/4/4	0/2/2/2

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	FE-ND	5.25	2.11	1.94
2	A	500	HEM	FE-NB	5.24	2.11	1.94
2	A	500	HEM	C1B-NB	-5.13	1.31	1.40
2	A	500	HEM	C4D-ND	-5.07	1.31	1.40
3	A	600	CPZ	C6-C4	-4.63	1.39	1.47
3	A	600	CPZ	C4-N3	-4.50	1.33	1.38
2	A	500	HEM	CMA-C3A	4.33	1.59	1.50
2	A	500	HEM	C1D-ND	-4.13	1.30	1.38
2	A	500	HEM	C1C-C2C	-3.97	1.37	1.45
2	A	500	HEM	CMC-C2C	3.91	1.58	1.50
2	A	500	HEM	C4B-NB	-3.79	1.31	1.38
2	A	500	HEM	CBB-CAB	3.26	1.46	1.30
2	A	500	HEM	CAC-C3C	-3.06	1.39	1.47
2	A	500	HEM	C1A-C2A	-2.95	1.38	1.44
2	A	500	HEM	CHD-C1D	-2.88	1.32	1.39
2	A	500	HEM	CMB-C2B	2.60	1.56	1.50
2	A	500	HEM	CBD-CGD	2.54	1.56	1.50
2	A	500	HEM	C4A-NA	-2.49	1.34	1.39
2	A	500	HEM	O2A-CGA	-2.49	1.22	1.30
2	A	500	HEM	FE-NC	2.46	2.03	1.95
2	A	500	HEM	CMD-C2D	2.46	1.55	1.50
2	A	500	HEM	C3B-C2B	2.29	1.41	1.37
3	A	600	CPZ	C2-N1	-2.27	1.33	1.37
2	A	500	HEM	C4A-C3A	-2.17	1.38	1.43
2	A	500	HEM	O2D-CGD	-2.17	1.23	1.30
2	A	500	HEM	C3C-C2C	2.08	1.41	1.37

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	HEM	C1B-NB-C4B	3.80	109.70	105.21
2	A	500	HEM	C4D-ND-C1D	3.42	109.26	105.21
2	A	500	HEM	C3B-C2B-C1B	-3.25	103.97	106.41
2	A	500	HEM	CHD-C4C-NC	-2.98	121.20	124.45
2	A	500	HEM	C4B-C3B-C2B	-2.58	104.91	107.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	HEM	C4C-CHD-C1D	2.07	130.42	126.02
2	A	500	HEM	CHC-C1C-NC	-2.03	122.24	124.45

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	500	HEM	CAA-CBA-CGA-O2A
2	A	500	HEM	CAA-CBA-CGA-O1A
2	A	500	HEM	CAD-CBD-CGD-O2D

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	465/476 (97%)	0.93	76 (16%) 4 4	26, 43, 76, 161	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	135	PHE	8.4
1	A	138	GLY	6.1
1	A	492	HIS	5.9
1	A	198	LEU	5.7
1	A	136	GLY	5.7
1	A	223	PHE	5.7
1	A	203	PHE	5.5
1	A	226	TYR	5.1
1	A	29	LYS	5.0
1	A	222	GLY	4.9
1	A	227	PHE	4.8
1	A	450	PHE	4.6
1	A	229	GLY	4.3
1	A	134	ASP	4.2
1	A	44	LEU	4.0
1	A	37	LEU	4.0
1	A	212	PHE	3.6
1	A	137	MET	3.5
1	A	380	TYR	3.5
1	A	28	GLY	3.5
1	A	30	LEU	3.4
1	A	194	VAL	3.3
1	A	230	THR	3.3
1	A	65	VAL	3.2
1	A	225	LYS	3.2
1	A	335	HIS	3.2
1	A	404	THR	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	379	GLY	3.1
1	A	286	GLN	3.0
1	A	228	PRO	3.0
1	A	191	LYS	3.0
1	A	60	GLU	3.0
1	A	139	LYS	2.9
1	A	253	ARG	2.9
1	A	49	LYS	2.8
1	A	66	PHE	2.7
1	A	35	SER	2.7
1	A	221	SER	2.7
1	A	297	PHE	2.7
1	A	72	SER	2.7
1	A	210	SER	2.7
1	A	48	ARG	2.6
1	A	202	PHE	2.6
1	A	224	LEU	2.6
1	A	131	THR	2.6
1	A	386	THR	2.6
1	A	388	VAL	2.5
1	A	220	PHE	2.5
1	A	140	ARG	2.5
1	A	259	SER	2.4
1	A	384	LYS	2.4
1	A	63	GLY	2.4
1	A	381	VAL	2.4
1	A	401	TYR	2.3
1	A	58	LEU	2.3
1	A	38	PRO	2.3
1	A	373	LYS	2.3
1	A	57	ARG	2.3
1	A	400	ARG	2.2
1	A	36	PRO	2.2
1	A	92	ALA	2.2
1	A	133	ARG	2.2
1	A	301	GLU	2.2
1	A	371	VAL	2.2
1	A	61	LYS	2.2
1	A	76	VAL	2.2
1	A	189	ASP	2.2
1	A	56	LEU	2.2
1	A	34	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	389	PHE	2.1
1	A	39	VAL	2.1
1	A	71	GLY	2.1
1	A	372	THR	2.1
1	A	148	GLU	2.1
1	A	132	MET	2.0
1	A	247	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CPZ	A	600	12/12	0.84	0.13	29,37,43,45	0
2	HEM	A	500	43/43	0.98	0.07	29,32,36,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.