



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 06:51 PM UTC

PDB ID : 6SUS / pdb_00006sus
Title : Crystal structure of RTX domain blocks IV and V of adenylate cyclase toxin from Bordetella pertussis
Authors : Motlova, L.; Barinka, C.; Bumba, L.
Deposited on : 2019-09-16
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	16% 76% 5% 19%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 2248 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Bifunctional hemolysin/adenylate cyclase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	18	0
			2008	1220	355	431	2			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	15	Total	Ca	0	0
			15	15		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		
3	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is NITRATE ION (CCD ID: NO3) (formula: NO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	N	O	0	0
			4	1	3		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	177	Total	O	0	0
			177	177		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional hemolysin/adenylate cyclase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	91.77Å 91.77Å 104.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.89 – 1.76 45.89 – 1.76	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.89-1.76) 99.9 (45.89-1.76)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.184 , 0.201 (Not available) , 0.203	Depositor DCC
R_{free} test set	2100 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	27.8	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2248	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, NO3, EDO, GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/2035	0.75	2/2752 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	1514	GLU	CA-C-N	5.54	125.21	119.56
1	A	1514	GLU	C-N-CA	5.54	125.21	119.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2008	0	1837	11	0
2	A	15	0	0	0	0
3	A	20	0	28	0	0
4	A	12	0	16	0	0
5	A	4	0	0	0	0
6	A	12	0	18	3	0
7	A	177	0	0	2	0
All	All	2248	0	1899	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1455[A]:LEU:HD12	1:A:1475[A]:LEU:CD2	2.07	0.84
1:A:1410:ILE:HB	1:A:1428[A]:LEU:HD23	1.78	0.66
6:A:1722:EDO:H22	7:A:1878:HOH:O	1.97	0.62
1:A:1455[A]:LEU:HD12	1:A:1475[A]:LEU:HD23	1.81	0.59
1:A:1410:ILE:HB	1:A:1428[A]:LEU:CD2	2.36	0.55
1:A:1455[A]:LEU:HD12	1:A:1475[A]:LEU:HD22	1.89	0.52
1:A:1613[A]:ILE:HG22	1:A:1615[A]:ALA:H	1.78	0.49
1:A:1410:ILE:HD12	1:A:1428[A]:LEU:HD21	1.98	0.46
6:A:1722:EDO:C2	7:A:1878:HOH:O	2.62	0.45
1:A:1648[A]:ASP:OD1	1:A:1651:HIS:ND1	2.51	0.43
1:A:1502:LEU:C	1:A:1502:LEU:HD12	2.43	0.43
1:A:1475[B]:LEU:HD12	1:A:1475[B]:LEU:N	2.34	0.43
1:A:1488:GLY:HA3	6:A:1722:EDO:H22	2.01	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/320 (85%)	261 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	198/220 (90%)	194 (98%)	4 (2%)	48 29

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1634	LEU
1	A	1663	VAL
1	A	1669	GLU
1	A	1673	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 15 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PGE	A	1716	-	9,9,9	0.39	0	8,8,8	0.34	0
4	GOL	A	1718	-	5,5,5	0.26	0	5,5,5	0.76	0
5	NO3	A	1719	-	1,3,3	0.27	0	0,3,3	-	-
6	EDO	A	1723	-	3,3,3	1.21	0	2,2,2	0.93	0
6	EDO	A	1722	-	3,3,3	0.45	0	2,2,2	0.65	0
6	EDO	A	1720	-	3,3,3	0.58	0	2,2,2	0.12	0
3	PGE	A	1721	-	9,9,9	0.38	0	8,8,8	0.48	0
4	GOL	A	1717	-	5,5,5	0.14	0	5,5,5	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	A	1716	-	-	1/7/7/7	-
4	GOL	A	1718	-	-	0/4/4/4	-
6	EDO	A	1723	-	-	1/1/1/1	-
6	EDO	A	1722	-	-	0/1/1/1	-
6	EDO	A	1720	-	-	1/1/1/1	-
3	PGE	A	1721	-	-	0/7/7/7	-
4	GOL	A	1717	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1723	EDO	O1-C1-C2-O2
3	A	1716	PGE	O1-C1-C2-O2
6	A	1720	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1722	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	258/320 (80%)	0.98	50 (19%) 3 3	12, 31, 61, 92	18 (6%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1497	ALA	8.6
1	A	1616	GLY	7.6
1	A	1513	ASP	6.7
1	A	1621	TRP	6.7
1	A	1671	LEU	6.7
1	A	1672	VAL	6.6
1	A	1617	ALA	6.5
1	A	1663	VAL	6.5
1	A	1675	MET	5.8
1	A	1666	ALA	5.5
1	A	1615[A]	ALA	5.3
1	A	1634	LEU	5.3
1	A	1668	ILE	4.4
1	A	1488	GLY	4.3
1	A	1635	GLY	4.3
1	A	1512	MET	4.3
1	A	1498	LYS	4.2
1	A	1673	GLU	4.1
1	A	1670	LYS	4.1
1	A	1665	GLN	4.0
1	A	1662	ALA	4.0
1	A	1660	ASN	3.9
1	A	1620	LEU	3.7
1	A	1598[A]	HIS	3.7
1	A	1499	GLY	3.7
1	A	1674	ALA	3.6
1	A	1605	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1659	ALA	3.3
1	A	1664	ASP	3.1
1	A	1410	ILE	3.1
1	A	1637	ASP	3.0
1	A	1618	ASP	3.0
1	A	1619	GLN	2.9
1	A	1636	THR	2.9
1	A	1661	GLN	2.9
1	A	1667	GLY	2.8
1	A	1422	ALA	2.8
1	A	1427	VAL	2.6
1	A	1632	ARG	2.6
1	A	1648[A]	ASP	2.5
1	A	1532	ARG	2.5
1	A	1420[A]	ARG	2.5
1	A	1633	ILE	2.4
1	A	1608	HIS	2.4
1	A	1625	GLN	2.2
1	A	1622	PHE	2.2
1	A	1669	GLU	2.2
1	A	1658	ALA	2.1
1	A	1594	VAL	2.1
1	A	1606	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	EDO	A	1720	4/4	0.74	0.23	48,51,51,58	0
6	EDO	A	1723	4/4	0.78	0.32	38,40,41,49	0
6	EDO	A	1722	4/4	0.83	0.19	44,52,55,60	0
3	PGE	A	1716	10/10	0.87	0.18	53,59,63,68	0
4	GOL	A	1718	6/6	0.91	0.12	37,40,45,45	0
3	PGE	A	1721	10/10	0.92	0.12	32,40,54,59	0
5	NO3	A	1719	4/4	0.94	0.09	47,49,52,56	0
4	GOL	A	1717	6/6	0.96	0.07	32,33,34,35	0
2	CA	A	1708	1/1	0.98	0.07	37,37,37,37	0
2	CA	A	1712	1/1	0.98	0.05	34,34,34,34	0
2	CA	A	1703	1/1	0.99	0.11	31,31,31,31	1
2	CA	A	1704	1/1	0.99	0.03	26,26,26,26	0
2	CA	A	1707	1/1	0.99	0.04	29,29,29,29	0
2	CA	A	1701	1/1	0.99	0.04	32,32,32,32	0
2	CA	A	1702	1/1	0.99	0.02	29,29,29,29	0
2	CA	A	1713	1/1	0.99	0.03	31,31,31,31	0
2	CA	A	1714	1/1	0.99	0.03	28,28,28,28	0
2	CA	A	1715	1/1	0.99	0.04	27,27,27,27	0
2	CA	A	1709	1/1	1.00	0.03	26,26,26,26	0
2	CA	A	1710	1/1	1.00	0.02	24,24,24,24	0
2	CA	A	1711	1/1	1.00	0.03	25,25,25,25	0
2	CA	A	1705	1/1	1.00	0.02	27,27,27,27	0
2	CA	A	1706	1/1	1.00	0.02	27,27,27,27	0

6.5 Other polymers

There are no such residues in this entry.