



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:37 AM UTC

PDB ID : 7SUV / pdb_00007suv
Title : APE1 exonuclease substrate complex with 8oxoG opposite A
Authors : Whitaker, A.W.; Freudenthal, B.D.
Deposited on : 2021-11-18
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	11	73% 55% 45%
2	D	10	90% 10%
3	E	21	33% 57% 43%
4	A	276	14% 92% 7%
4	B	276	9% 86% 14%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5553 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(P*TP*CP*GP*AP*CP*GP*GP*AP*TP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	11	Total	C	N	O	P	0	0	0
			225	106	41	67	11			

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*CP*TP*GP*AP*TP*GP*CP*GP*(80G))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	10	Total	C	N	O	P	0	0	0
			207	98	40	60	9			

- Molecule 3 is a DNA chain called DNA (5'-D(*GP*GP*AP*TP*CP*CP*GP*TP*CP*GP*AP*AP*CP*GP*CP*AP*TP*CP*AP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	21	Total	C	N	O	P	0	0	0
			427	203	82	122	20			

- Molecule 4 is a protein called DNA-(apurinic or apyrimidinic site) lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	276	Total	C	N	O	S	0	3	0
			2207	1410	383	406	8			
4	B	276	Total	C	N	O	S	0	4	0
			2224	1419	389	408	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	96	GLN	GLU	conflict	UNP P27695
A	138	ALA	CYS	conflict	UNP P27695
A	210	ASN	ASP	conflict	UNP P27695

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	96	GLN	GLU	conflict	UNP P27695
B	138	ALA	CYS	conflict	UNP P27695
B	210	ASN	ASP	conflict	UNP P27695


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	O	0	0
			1	1		
5	D	10	Total	O	0	0
			10	10		
5	E	12	Total	O	0	0
			12	12		
5	A	101	Total	O	0	0
			101	101		
5	B	139	Total	O	0	0
			139	139		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(P*TP*CP*GP*AP*CP*GP*GP*AP*TP*CP*C)-3')




- Molecule 2: DNA (5'-D(*GP*CP*TP*GP*AP*TP*GP*CP*GP*(8OG))-3')




- Molecule 3: DNA (5'-D(*GP*GP*AP*TP*CP*CP*GP*TP*CP*GP*AP*AP*CP*GP*CP*AP*TP*CP*AP*GP*C)-3')




- Molecule 4: DNA-(apurinic or apyrimidinic site) lyase



- Molecule 4: DNA-(apurinic or apyrimidinic site) lyase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.11Å 66.49Å 91.98Å 90.00° 110.69° 90.00°	Depositor
Resolution (Å)	24.96 – 1.99 24.96 – 1.99	Depositor EDS
% Data completeness (in resolution range)	89.1 (24.96-1.99) 73.4 (24.96-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 1.99Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.226 , 0.265 0.226 , 0.265	Depositor DCC
R_{free} test set	1992 reflections (3.60%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.514	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 28.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5553	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8OG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.51	1/251 (0.4%)	0.43	0/383
2	D	0.51	0/206	0.64	0/317
3	E	0.43	0/479	0.61	0/737
4	A	0.36	0/2271	0.56	0/3079
4	B	0.35	0/2285	0.56	1/3096 (0.0%)
All	All	0.38	1/5492 (0.0%)	0.56	1/7612 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	2	DT	OP3-P	6.43	1.61	1.48

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	111	LEU	N-CA-C	5.41	121.77	109.81

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	225	0	124	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	207	0	114	1	0
3	E	427	0	236	12	0
4	A	2207	0	2187	13	0
4	B	2224	0	2204	23	0
5	A	101	0	0	1	0
5	B	139	0	0	0	0
5	C	1	0	0	0	0
5	D	10	0	0	0	0
5	E	12	0	0	0	0
All	All	5553	0	4865	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:17:DT:H5'	4:B:127:GLY:HA3	1.72	0.71
1:C:6:DC:H42	3:E:6:DC:H42	1.40	0.70
1:C:3:DC:H4'	4:B:271:MET:HE2	1.74	0.68
4:A:245:GLN:HG3	4:B:293:PRO:HG2	1.76	0.66
4:A:101:GLU:HA	4:A:104:LEU:HD13	1.78	0.64
3:E:5:DC:H6	3:E:5:DC:H5''	1.64	0.62
4:A:177:ARG:NH1	5:A:401:HOH:O	2.26	0.62
4:A:61:THR:HB	4:A:318:LEU:H	1.66	0.60
3:E:5:DC:H2'	3:E:6:DC:C6	2.39	0.58
3:E:6:DC:H2'	3:E:7:DG:C8	2.40	0.56
1:C:6:DC:N4	3:E:6:DC:H42	2.01	0.56
3:E:13:DC:C2	4:B:270:MET:HE2	2.42	0.55
4:B:76:ILE:HD12	4:B:105:PRO:HG2	1.88	0.55
2:D:10:8OG:OP2	4:B:174:ASN:HB2	2.08	0.53
4:A:154:GLU:OE2	4:A:181:ARG:NH2	2.42	0.52
4:A:73:ARG:NH1	4:A:103:LYS:HD2	2.25	0.51
4:A:150:GLU:CD	4:A:187:ARG:HH21	2.18	0.51
4:B:174:ASN:OD1	4:B:176:GLY:N	2.39	0.50
4:A:104:LEU:HD21	4:A:119:TRP:CE2	2.47	0.49
4:B:268:THR:OG1	4:B:270:MET:HE3	2.13	0.49
3:E:3:DA:H2''	3:E:4:DT:H5'	1.95	0.48
4:B:110:GLU:O	4:B:112:PRO:HD3	2.14	0.47
4:A:231:GLY:H	4:A:236:GLU:CD	2.23	0.47
3:E:7:DG:H2''	3:E:8:DT:H5''	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:262:TYR:HA	4:B:264:TYR:CZ	2.49	0.47
3:E:4:DT:H2''	3:E:5:DC:H5''	1.97	0.46
4:B:150:GLU:OE1	4:B:187:ARG:NH2	2.49	0.46
4:A:122:PRO:HB3	4:A:154:GLU:HA	1.96	0.45
1:C:7:DG:C6	1:C:8:DG:C6	3.04	0.45
4:B:81:LEU:HD22	4:B:111:LEU:HD11	1.98	0.45
4:A:73:ARG:HH12	4:A:103:LYS:HD2	1.82	0.45
4:B:50:ASP:OD1	4:B:299:LYS:HE2	2.17	0.45
4:B:172:VAL:HG13	4:B:173:PRO:HD2	1.99	0.44
4:A:318:LEU:HD23	4:A:318:LEU:HA	1.78	0.43
4:B:114:LEU:HB3	4:B:134:LEU:HB3	2.00	0.43
4:B:161:GLU:OE2	4:B:203:LYS:NZ	2.49	0.42
4:B:72:LEU:HD21	4:B:108:LEU:HD11	2.02	0.42
4:B:265:THR:HG23	4:B:281:ARG:HG3	2.01	0.41
3:E:5:DC:H2'	3:E:6:DC:H6	1.84	0.41
4:B:254:ARG:HD3	4:B:254:ARG:HA	1.90	0.41
4:B:73:ARG:HG2	4:B:105:PRO:HG3	2.03	0.41
4:B:318:LEU:HA	4:B:318:LEU:HD23	1.82	0.40
3:E:11:DA:H2'	3:E:11:DA:OP2	2.21	0.40
4:A:224:LYS:HB2	4:A:224:LYS:HE2	1.82	0.40
4:B:196:LEU:HD22	4:B:205:LEU:HD21	2.03	0.40
4:B:209:GLY:O	4:B:283:ASP:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	A	277/276 (100%)	268 (97%)	9 (3%)	0	100	100
4	B	278/276 (101%)	269 (97%)	9 (3%)	0	100	100
All	All	555/552 (100%)	537 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	A	237/235 (101%)	232 (98%)	5 (2%)	47	52
4	B	238/235 (101%)	234 (98%)	4 (2%)	53	60
All	All	475/470 (101%)	466 (98%)	9 (2%)	50	56

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	A	61	THR
4	A	109	GLN
4	A	110	GLU
4	A	180	VAL
4	A	270	MET
4	B	68	ASN
4	B	109	GLN
4	B	181	ARG
4	B	213	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
4	A	68	ASN
4	A	96	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	8OG	D	10	2	22,25,26	2.26	6 (27%)	26,37,40	2.00	7 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8OG	D	10	2	-	1/7/21/22	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	10	8OG	C5-N7	5.78	1.47	1.37
2	D	10	8OG	C8-N9	4.79	1.49	1.40
2	D	10	8OG	O8-C8	-4.00	1.16	1.23
2	D	10	8OG	C6-N1	-3.09	1.33	1.38
2	D	10	8OG	C8-N7	-2.50	1.33	1.38
2	D	10	8OG	C5-C4	2.37	1.40	1.37

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	10	8OG	C2-N1-C6	-4.62	116.73	125.11
2	D	10	8OG	N9-C4-N3	4.59	131.86	126.13
2	D	10	8OG	C5-C6-N1	3.36	121.36	112.13
2	D	10	8OG	O6-C6-C5	-3.33	119.23	127.26
2	D	10	8OG	O8-C8-N9	2.87	129.90	126.00
2	D	10	8OG	O8-C8-N7	-2.73	121.35	126.47
2	D	10	8OG	C2-N3-C4	2.38	116.39	112.30

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	10	8OG	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	10	8OG	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	11/11 (100%)	2.49	8 (72%) 0 0	61, 92, 101, 101	0
2	D	9/10 (90%)	0.51	0 100 100	40, 43, 48, 50	0
3	E	21/21 (100%)	1.41	7 (33%) 1 1	40, 57, 91, 100	0
4	A	276/276 (100%)	1.20	38 (13%) 6 6	29, 40, 60, 76	3 (1%)
4	B	276/276 (100%)	1.10	26 (9%) 14 13	18, 38, 56, 73	4 (1%)
All	All	593/594 (99%)	1.17	79 (13%) 7 6	18, 40, 64, 101	7 (1%)

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	A	112	PRO	4.3
4	A	109	GLN	4.0
4	B	180	VAL	3.8
4	B	176	GLY	3.7
4	B	112	PRO	3.6
4	B	44	LEU	3.5
1	C	12	DC	3.3
4	B	111	LEU	3.3
1	C	6	DC	3.3
1	C	4	DG	3.2
3	E	1	DG	3.2
4	A	111	LEU	3.0
4	A	272	ASN	3.0
4	B	102	ASN	3.0
4	A	108	LEU	3.0
4	A	318	LEU	2.9
4	B	318	LEU	2.9
4	B	99	CYS	2.9
4	A	182	LEU	2.8
3	E	12	DA	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	B	221[A]	ARG	2.8
4	B	224	LYS	2.8
4	B	108	LEU	2.8
4	A	201	SER	2.7
4	A	140	LEU	2.7
4	B	43	ALA	2.7
4	A	177	ARG	2.7
1	C	5	DA	2.7
3	E	6	DC	2.7
4	A	270	MET	2.7
4	A	110	GLU	2.7
4	A	224	LYS	2.6
4	A	146	ILE	2.6
4	A	61	THR	2.6
4	A	128	TYR	2.5
1	C	8	DG	2.5
4	B	124	ASP	2.5
4	B	51[A]	GLN	2.5
4	A	164	SER	2.4
1	C	7	DG	2.4
4	B	225	GLY	2.4
4	B	45	TYR	2.4
4	B	184	TYR	2.4
4	B	110	GLU	2.4
4	A	148[A]	ASP	2.4
4	A	142	VAL	2.4
4	A	44	LEU	2.3
1	C	10	DT	2.3
3	E	5	DC	2.3
4	A	125	LYS	2.3
4	A	113	GLY	2.3
4	A	180	VAL	2.3
4	A	195	PHE	2.3
4	A	129[A]	SER	2.2
4	A	184	TYR	2.2
4	B	270	MET	2.2
4	A	149	GLU	2.2
4	A	245	GLN	2.2
4	A	147	GLY	2.2
4	B	175	ALA	2.2
4	B	171	TYR	2.2
4	B	269	TYR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	A	188	TRP	2.2
4	A	228	LYS	2.2
4	B	177	ARG	2.2
4	A	104	LEU	2.2
1	C	9	DA	2.2
3	E	10	DG	2.2
4	A	53	THR	2.1
4	B	72	LEU	2.1
4	B	187	ARG	2.1
4	A	130	GLY	2.1
3	E	7	DG	2.0
4	A	116	HIS	2.0
4	B	289	HIS	2.0
4	A	62	LEU	2.0
4	A	170	ALA	2.0
3	E	8	DT	2.0
4	A	247	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	8OG	D	10	23/24	0.78	0.26	41,57,68,72	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.