



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:59 AM UTC

PDB ID : 5SVD / pdb_00005svd
Title : Nop9, a new PUF-like protein, prevents premature pre-rRNA cleavage to correctly process mature 18S rRNA
Authors : Zhang, J.; Qiu, C.; Hall, T.
Deposited on : 2016-08-05
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	600	5% 80% 10% 10%
1	B	600	9% 79% 10% 11%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9186 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleolar protein 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	543	Total	C	N	O	S	0	1	0
			4443	2865	737	820	21			
1	B	534	Total	C	N	O	S	0	2	0
			4366	2812	725	808	21			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	430	THR	ALA	engineered mutation	UNP P47077
B	430	THR	ALA	engineered mutation	UNP P47077


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	290	Total	O	0	0
			290	290		
2	B	87	Total	O	0	0
			87	87		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nucleolar protein 9



• Molecule 1: Nucleolar protein 9



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.50Å 110.47Å 114.85Å 90.00° 92.22° 90.00°	Depositor
Resolution (Å)	32.02 – 2.10 32.02 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.8 (32.02-2.10) 96.0 (32.02-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.185 , 0.221 0.190 , 0.224	Depositor DCC
R_{free} test set	2023 reflections (2.11%)	wwPDB-VP
Wilson B-factor (Å ²)	40.1	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.001 for -h,l,k 0.011 for -h,-l,-k 0.024 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9186	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/4526	0.83	5/6104 (0.1%)
1	B	0.45	0/4449	0.76	1/5996 (0.0%)
All	All	0.54	0/8975	0.80	6/12100 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	279	SER	CA-C-N	8.07	136.22	121.70
1	A	279	SER	C-N-CA	8.07	136.22	121.70
1	A	160	GLU	N-CA-C	-6.75	105.01	113.18
1	A	285	GLN	N-CA-C	-6.22	101.07	110.28
1	B	276	GLY	N-CA-C	-5.49	106.02	111.56
1	A	185	MET	CG-SD-CE	-5.39	89.05	100.90

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	284	SER	Peptide
1	B	328	ALA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4443	0	4505	35	0
1	B	4366	0	4422	31	0
2	A	290	0	0	4	0
2	B	87	0	0	0	0
All	All	9186	0	8927	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:281:SER:HB2	1:A:282:ASP:HA	1.52	0.88
1:B:378:LYS:HE3	1:B:382:ARG:HH21	1.59	0.68
1:B:542:ILE:HG22	1:B:546:MET:HE2	1.79	0.64
1:B:153:SER:HB3	1:B:178:MET:HE2	1.84	0.59
1:A:483:GLU:OE1	1:A:486:ARG:NH1	2.33	0.57
1:A:50:GLN:HB2	1:A:135:GLY:HA3	1.86	0.57
1:A:375:ARG:NH1	2:A:705:HOH:O	2.29	0.55
1:A:620:ASP:OD2	2:A:701:HOH:O	2.18	0.55
1:B:139:HIS:HB3	1:B:142:ALA:HB3	1.89	0.54
1:A:279:SER:HB3	1:A:281:SER:O	2.08	0.53
1:B:199:HIS:HB3	1:B:202:ALA:HB3	1.90	0.53
1:B:328:ALA:HB1	1:B:329:ASP:CG	2.34	0.53
1:B:183:LEU:O	1:B:187:ASN:ND2	2.35	0.53
1:B:439:ARG:O	1:B:443:ILE:HG12	2.09	0.52
1:B:123:LYS:O	1:B:127:GLN:HG3	2.10	0.52
1:A:474:ASN:ND2	1:A:478:ASP:HB2	2.25	0.51
1:B:256:GLU:C	1:B:258:PHE:H	2.19	0.51
1:A:280:ARG:HG3	1:A:283:ILE:HG13	1.92	0.51
1:B:530:VAL:O	1:B:533:VAL:HG22	2.11	0.51
1:A:518:ILE:HD12	1:A:554:GLU:HG3	1.93	0.50
1:A:280:ARG:HD3	1:A:283:ILE:HD12	1.92	0.50
1:A:254:THR:O	1:A:259:LYS:NZ	2.45	0.50
1:B:439:ARG:HA	1:B:497:TYR:OH	2.10	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177:THR:HG23	1:B:180:ASN:H	1.76	0.49
1:A:194:LYS:HD2	1:A:272:GLY:HA3	1.93	0.48
1:A:500:LYS:NZ	2:A:718:HOH:O	2.46	0.48
1:A:119:GLU:OE2	1:A:159:ARG:NH1	2.46	0.48
1:A:284:SER:O	1:A:287:THR:HB	2.14	0.48
1:A:281:SER:CB	1:A:282:ASP:HA	2.30	0.47
1:A:364:VAL:HG12	1:A:397:LEU:HD21	1.96	0.47
1:B:261:GLU:O	1:B:265:ILE:HG12	2.16	0.46
1:A:542:ILE:HG22	1:A:546:MET:HE2	1.96	0.46
1:A:158:GLU:O	1:A:162:LEU:HG	2.16	0.46
1:B:291:PHE:CZ	1:B:304:ILE:HD12	2.50	0.46
1:B:455:SER:HB3	1:B:458:LYS:HB3	1.97	0.46
1:A:280:ARG:CG	1:A:281:SER:HA	2.46	0.45
1:B:523:HIS:O	1:B:527:SER:HB3	2.17	0.45
1:A:139:HIS:HB3	1:A:142:ALA:HB3	1.98	0.45
1:B:562:VAL:HG22	1:B:601:TYR:CE2	2.52	0.45
1:A:60:GLU:OE2	1:A:96:LYS:NZ	2.34	0.45
1:A:464[A]:CYS:SG	1:A:490:PHE:HD2	2.39	0.45
1:B:531:GLU:HG3	1:B:570:LYS:HD2	1.98	0.45
1:B:159:ARG:HA	1:B:160:GLU:HA	1.75	0.44
1:A:63:TYR:CE2	1:A:92:GLU:HG2	2.52	0.44
1:A:354:ASN:ND2	2:A:722:HOH:O	2.50	0.44
1:A:181:MET:HE3	1:A:181:MET:HB2	1.85	0.43
1:A:569:ASP:CG	1:A:608:ASN:HD22	2.26	0.43
1:A:178:MET:HE3	1:A:181:MET:HB3	2.01	0.43
1:B:464[B]:CYS:O	1:B:487:ARG:HD3	2.19	0.43
1:A:280:ARG:HG3	1:A:281:SER:HA	2.00	0.42
1:A:321:TRP:CZ3	1:A:359:ALA:HB2	2.54	0.42
1:A:154:ALA:O	1:A:157:VAL:HG22	2.18	0.42
1:B:153:SER:O	1:B:157:VAL:HG23	2.21	0.41
1:A:480:PRO:HB3	1:A:523:HIS:CE1	2.55	0.41
1:B:123:LYS:HD3	1:B:181:MET:HE1	2.03	0.41
1:B:323:LEU:HA	1:B:323:LEU:HD23	1.83	0.41
1:A:291:PHE:CZ	1:A:304:ILE:HD12	2.55	0.41
1:B:483:GLU:OE1	1:B:486:ARG:NH1	2.54	0.41
1:A:530:VAL:O	1:A:533:VAL:HG22	2.21	0.40
1:B:471:THR:O	1:B:474:ASN:ND2	2.47	0.40
1:B:80:GLU:OE1	1:B:80:GLU:N	2.50	0.40
1:B:436:GLY:O	1:B:439:ARG:HB2	2.21	0.40
1:A:197:MET:HE3	1:A:269:LEU:HD22	2.02	0.40
1:B:397:LEU:HD13	1:B:401:ASP:HB3	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:452:PRO:C	1:B:454:LYS:H	2.29	0.40
1:B:455:SER:C	1:B:457:ALA:H	2.28	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	538/600 (90%)	522 (97%)	16 (3%)	0	100	100
1	B	530/600 (88%)	513 (97%)	16 (3%)	1 (0%)	43	44
All	All	1068/1200 (89%)	1035 (97%)	32 (3%)	1 (0%)	48	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	278	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	495/554 (89%)	489 (99%)	6 (1%)	63	72
1	B	486/554 (88%)	482 (99%)	4 (1%)	73	81
All	All	981/1108 (88%)	971 (99%)	10 (1%)	68	76

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	LYS
1	A	194	LYS
1	A	339	LEU
1	A	376	ILE
1	A	415	MET
1	A	584	ARG
1	B	108	LYS
1	B	176	VAL
1	B	527	SER
1	B	549	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	144	HIS
1	B	127	GLN
1	B	180	ASN
1	B	305	GLN
1	B	354	ASN
1	B	503	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	543/600 (90%)	0.18	31 (5%)	29 31	20, 44, 91, 127	1 (0%)
1	B	534/600 (89%)	0.79	52 (9%)	13 14	26, 64, 101, 149	2 (0%)
All	All	1077/1200 (89%)	0.48	83 (7%)	19 21	20, 56, 95, 149	3 (0%)

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	456	ASP	8.0
1	B	277	ALA	6.5
1	A	280	ARG	5.6
1	B	478	ASP	5.2
1	A	282	ASP	4.9
1	A	279	SER	4.8
1	A	254	THR	4.5
1	A	175	TYR	4.2
1	B	329	ASP	4.2
1	B	479	TRP	4.1
1	A	281	SER	4.0
1	B	255	PRO	3.9
1	A	276	GLY	3.8
1	A	286	SER	3.8
1	A	256	GLU	3.8
1	B	328	ALA	3.7
1	A	421	MET	3.6
1	A	50	GLN	3.6
1	A	285	GLN	3.6
1	B	278	GLU	3.5
1	B	600	ILE	3.4
1	B	155	ALA	3.1
1	B	285	GLN	3.1
1	A	479	TRP	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	443	ILE	3.0
1	A	630	PHE	3.0
1	B	601	TYR	2.9
1	B	314	PHE	2.9
1	A	251	VAL	2.8
1	B	332	ASP	2.8
1	B	435	GLY	2.8
1	A	159	ARG	2.7
1	B	257	SER	2.7
1	B	89	ILE	2.7
1	A	160	GLU	2.7
1	A	255	PRO	2.7
1	A	252	TYR	2.7
1	B	497	TYR	2.7
1	B	95	GLY	2.7
1	B	258	PHE	2.6
1	B	177	THR	2.6
1	A	456	ASP	2.6
1	B	176	VAL	2.6
1	B	501	PHE	2.5
1	B	76	PHE	2.5
1	A	540	ASP	2.5
1	B	86	VAL	2.5
1	B	159	ARG	2.5
1	B	529	VAL	2.4
1	A	283	ILE	2.4
1	B	101	VAL	2.4
1	B	440	ASP	2.4
1	B	280	ARG	2.4
1	A	477	ASP	2.4
1	A	454	LYS	2.4
1	B	563	TYR	2.3
1	B	156	LEU	2.3
1	A	181	MET	2.3
1	A	453	GLU	2.3
1	B	105	ILE	2.3
1	B	480	PRO	2.3
1	A	277	ALA	2.3
1	B	313	ILE	2.2
1	B	495	ILE	2.2
1	B	276	GLY	2.2
1	B	602	GLY	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	455	SER	2.2
1	B	475	THR	2.2
1	B	124	ASP	2.2
1	B	504	ILE	2.2
1	B	78	ALA	2.2
1	B	630	PHE	2.1
1	B	464[A]	CYS	2.1
1	A	178	MET	2.1
1	B	275	ASN	2.1
1	A	163	THR	2.1
1	B	194	LYS	2.1
1	B	178	MET	2.1
1	B	453	GLU	2.1
1	A	51	MET	2.0
1	B	91	GLU	2.0
1	A	78	ALA	2.0
1	B	436	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.