



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 02:36 AM EDT

PDB ID : 8SV2 / pdb_00008sv2
Title : Pasteurella multocida alpha2,3/2,6 sialyltransferase D141N bound to CMP
Authors : Stubbs, H.E.; Iverson, T.M.
Deposited on : 2023-05-15
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	14% 79% 20%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3526 atoms, of which 122 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alpha-2,3/2,6-sialyltransferase/sialidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	387	3244	2022	110	521	584	7	0	1	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	26	MET	-	initiating methionine	UNP Q15KI8
A	142	ASN	ASP	conflict	UNP Q15KI8

- Molecule 2 is CYTIDINE-5'-MONOPHOSPHATE (CCD ID: C5P) (formula: $C_9H_{14}N_3O_8P$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	21	9	3	8	1	0	0

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0
			20	4	12	1	3	

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	224	Total	O	0	0
			224	224		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-2,3/2,6-sialyltransferase/sialidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.60Å 66.40Å 96.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.89 – 1.84 31.89 – 1.84	Depositor EDS
% Data completeness (in resolution range)	99.4 (31.89-1.84) 99.5 (31.89-1.84)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.95 (at 1.84Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.215 , 0.254 0.215 , 0.253	Depositor DCC
R_{free} test set	1869 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	26.7	Xtriage
Anisotropy	0.665	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.025 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3526	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: C5P, TRS, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	1/3209 (0.0%)	0.61	2/4359 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	235	THR	CA-C	-5.05	1.46	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	381	LYS	N-CA-C	-8.95	91.75	110.80
1	A	383	ALA	N-CA-C	-5.53	106.91	113.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3134	110	3050	94	1
2	A	21	0	12	0	0
3	A	8	12	12	3	0
4	A	17	0	23	5	0
5	A	224	0	0	16	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3404	122	3097	95	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:284:GLN:HG3	4:A:503:PGE:H2	1.40	1.03
1:A:26:MET:HB2	1:A:104:GLU:OE2	1.58	1.03
1:A:163:LYS:HE3	5:A:786:HOH:O	1.70	0.91
1:A:38:LEU:CD2	1:A:226:MET:HE1	2.10	0.81
1:A:333:PRO:O	5:A:602:HOH:O	1.96	0.81
1:A:143:GLY:HA2	1:A:201:LEU:HD11	1.63	0.79
1:A:228:TRP:CE3	1:A:342:MET:HG3	2.18	0.78
1:A:362:LEU:HG	1:A:363:PRO:HD2	1.64	0.78
1:A:360:PHE:HE1	1:A:388:TYR:CE1	2.03	0.75
1:A:350:LYS:NZ	5:A:603:HOH:O	2.17	0.74
1:A:74:GLN:NE2	5:A:605:HOH:O	2.21	0.72
1:A:170:SER:O	1:A:174:LEU:HG	1.90	0.71
1:A:284:GLN:OE1	4:A:503:PGE:H3	1.91	0.71
1:A:200:PHE:O	1:A:225:LYS:HE2	1.91	0.70
1:A:383:ALA:C	1:A:384:LEU:HD12	2.18	0.69
1:A:360:PHE:CE1	1:A:388:TYR:CE1	2.80	0.69
1:A:201:LEU:O	1:A:225:LYS:HE3	1.93	0.69
1:A:26:MET:HB2	1:A:104:GLU:CD	2.19	0.67
1:A:326:ALA:HB3	1:A:329:ILE:HD13	1.77	0.67
1:A:304:HIS:ND1	5:A:606:HOH:O	2.30	0.64
1:A:371:ILE:HG13	1:A:404:ILE:HB	1.80	0.64
1:A:207:GLU:HG2	1:A:216:LYS:HE2	1.79	0.63
1:A:378:VAL:HG13	1:A:380:SER:O	1.99	0.62
1:A:384:LEU:HA	5:A:649:HOH:O	1.99	0.62
1:A:284:GLN:CG	4:A:503:PGE:H2	2.24	0.60
1:A:211:PHE:CE2	1:A:212:LEU:HD13	2.37	0.59
1:A:392:MET:HE2	1:A:397:ILE:HG21	1.84	0.59
1:A:38:LEU:HD23	1:A:226:MET:HE1	1.84	0.58
1:A:336:ILE:N	5:A:602:HOH:O	2.37	0.56
1:A:197:LYS:HE2	5:A:756:HOH:O	2.04	0.56
1:A:378:VAL:HG23	1:A:386[A]:ASN:ND2	2.20	0.56
1:A:388:TYR:HB2	3:A:502:TRS:O2	2.05	0.56
1:A:382:ASP:C	1:A:384:LEU:H	2.16	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:TRP:CB	5:A:789:HOH:O	2.56	0.54
1:A:141:ASP:HB3	1:A:201:LEU:HD12	1.90	0.54
1:A:67:ILE:HD12	1:A:245:LEU:HD22	1.89	0.53
1:A:26:MET:CB	1:A:104:GLU:CG	2.86	0.53
1:A:306:LYS:HG2	1:A:308:TYR:CE1	2.43	0.53
1:A:382:ASP:C	1:A:384:LEU:N	2.63	0.52
1:A:280:TYR:OH	4:A:503:PGE:H4	2.10	0.52
1:A:143:GLY:HA2	1:A:201:LEU:CD1	2.37	0.51
1:A:26:MET:HB2	1:A:104:GLU:CG	2.40	0.51
1:A:275:THR:O	1:A:279:GLU:HG3	2.09	0.51
1:A:388:TYR:CB	3:A:502:TRS:O2	2.59	0.51
1:A:38:LEU:HB3	1:A:39:PRO:HD3	1.93	0.50
1:A:387:PRO:HA	1:A:390:LYS:HE2	1.93	0.50
1:A:380:SER:O	1:A:382:ASP:N	2.44	0.50
1:A:384:LEU:HD22	1:A:393:ARG:NH1	2.28	0.49
1:A:278:ARG:NH1	5:A:614:HOH:O	2.44	0.48
1:A:274:ASN:O	1:A:275:THR:C	2.57	0.48
1:A:38:LEU:HD21	1:A:226:MET:HE1	1.95	0.47
1:A:148:VAL:HG13	1:A:387:PRO:HB2	1.97	0.47
1:A:225:LYS:NZ	1:A:228:TRP:HE1	2.13	0.47
4:A:503:PGE:H12	5:A:632:HOH:O	2.13	0.47
1:A:26:MET:SD	1:A:104:GLU:HG2	2.55	0.47
1:A:154:GLU:HA	1:A:211:PHE:CD2	2.51	0.47
1:A:26:MET:HB3	1:A:104:GLU:HG3	1.97	0.46
1:A:378:VAL:CG2	1:A:386[A]:ASN:ND2	2.78	0.46
1:A:38:LEU:CG	1:A:226:MET:HE1	2.45	0.46
1:A:378:VAL:CG2	1:A:386[A]:ASN:HD22	2.29	0.46
1:A:392:MET:HE2	1:A:397:ILE:CG2	2.44	0.46
1:A:381:LYS:O	1:A:383:ALA:N	2.37	0.46
1:A:384:LEU:HB3	1:A:389:VAL:HG11	1.98	0.46
1:A:275:THR:O	1:A:276:ASP:C	2.57	0.46
1:A:38:LEU:HG	1:A:226:MET:HE1	1.98	0.45
1:A:26:MET:HB3	1:A:104:GLU:CG	2.47	0.45
1:A:141:ASP:CB	1:A:201:LEU:HD12	2.47	0.45
1:A:38:LEU:HD11	1:A:228:TRP:CH2	2.51	0.45
1:A:380:SER:C	1:A:382:ASP:N	2.75	0.45
1:A:82:GLU:OE2	5:A:604:HOH:O	2.21	0.45
1:A:392:MET:CE	1:A:397:ILE:HG21	2.47	0.45
1:A:397:ILE:HG22	1:A:398:ILE:HG23	1.98	0.45
1:A:386[A]:ASN:O	1:A:390:LYS:HG3	2.17	0.45
1:A:225:LYS:HZ1	1:A:228:TRP:HE1	1.65	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:362:LEU:HG	1:A:363:PRO:CD	2.40	0.44
1:A:228:TRP:CD2	1:A:342:MET:HG3	2.52	0.44
1:A:372:PHE:CD2	3:A:502:TRS:H11	2.53	0.44
1:A:407:ASP:OD1	1:A:407:ASP:N	2.51	0.43
1:A:326:ALA:CB	1:A:329:ILE:HD13	2.48	0.42
1:A:38:LEU:CD2	1:A:201:LEU:HD21	2.48	0.42
1:A:310:LYS:HE2	5:A:602:HOH:O	2.19	0.42
1:A:334:ALA:C	5:A:602:HOH:O	2.62	0.42
1:A:225:LYS:NZ	1:A:228:TRP:NE1	2.68	0.41
1:A:225:LYS:NZ	1:A:228:TRP:CD1	2.88	0.41
1:A:26:MET:CB	1:A:104:GLU:HG3	2.50	0.41
1:A:258:VAL:HG22	1:A:262:LYS:HE3	2.02	0.41
1:A:258:VAL:HG23	1:A:260:GLN:HB3	2.02	0.41
1:A:119:ARG:HB3	1:A:120:PRO:HD3	2.03	0.41
1:A:122:LEU:HD13	1:A:173:LEU:HD23	2.03	0.41
1:A:400:GLU:H	1:A:400:GLU:HG3	1.64	0.41
1:A:163:LYS:CE	5:A:786:HOH:O	2.47	0.41
1:A:252:VAL:O	1:A:255:SER:OG	2.26	0.40
1:A:174:LEU:HD12	5:A:721:HOH:O	2.21	0.40
1:A:332:ILE:O	1:A:333:PRO:C	2.64	0.40
1:A:130:ASP:OD1	1:A:130:ASP:N	2.55	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:LYS:NZ	1:A:328:ASN:OD1[4_555]	1.84	0.36
5:A:603:HOH:O	5:A:607:HOH:O[3_454]	2.10	0.10

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	386/388 (100%)	369 (96%)	16 (4%)	1 (0%)	36 25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	382	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	337/352 (96%)	336 (100%)	1 (0%)	86 81

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	289	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	74	GLN
1	A	128	HIS
1	A	135	GLN
1	A	136	GLN
1	A	331	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	C5P	A	501	-	22,22,22	0.95	1 (4%)	32,33,33	0.69	1 (3%)
4	PGE	A	503	-	9,9,9	0.35	0	8,8,8	0.40	0
3	TRS	A	502	-	7,7,7	0.62	0	9,9,9	0.47	0
4	PGE	A	504	-	6,6,9	0.39	0	5,5,8	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C5P	A	501	-	-	3/10/26/26	0/2/2/2
4	PGE	A	503	-	-	1/7/7/7	-
3	TRS	A	502	-	-	6/9/9/9	-
4	PGE	A	504	-	-	2/4/4/7	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	C5P	P-O1P	3.52	1.61	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	C5P	O2P-P-O1P	2.07	118.90	110.83

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	C5P	C5'-O5'-P-O3P
3	A	502	TRS	C1-C-C3-O3
3	A	502	TRS	C2-C-C3-O3
2	A	501	C5P	C5'-O5'-P-O1P
3	A	502	TRS	N-C-C3-O3
2	A	501	C5P	C5'-O5'-P-O2P
4	A	503	PGE	C4-C3-O2-C2
3	A	502	TRS	C2-C-C1-O1
4	A	504	PGE	O1-C1-C2-O2
4	A	504	PGE	C1-C2-O2-C3
3	A	502	TRS	C3-C-C1-O1
3	A	502	TRS	N-C-C1-O1

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	PGE	5	0
3	A	502	TRS	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/388 (99%)	0.77	54 (13%) 6 6	17, 32, 58, 89	1 (0%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	383	ALA	10.2
1	A	342	MET	4.9
1	A	384	LEU	4.7
1	A	277	VAL	4.7
1	A	126	PHE	3.2
1	A	377	GLN	3.1
1	A	379	LYS	3.1
1	A	258	VAL	3.1
1	A	201	LEU	3.1
1	A	341	LEU	3.1
1	A	388	TYR	3.0
1	A	397	ILE	2.8
1	A	346	LEU	2.8
1	A	343	MET	2.8
1	A	275	THR	2.7
1	A	362	LEU	2.7
1	A	387	PRO	2.7
1	A	220	ALA	2.7
1	A	281	TYR	2.7
1	A	276	ASP	2.7
1	A	378	VAL	2.7
1	A	228	TRP	2.6
1	A	338	PHE	2.6
1	A	280	TYR	2.6
1	A	399	ASP	2.6
1	A	391	VAL	2.6
1	A	381	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	273	GLY	2.5
1	A	360	PHE	2.5
1	A	301	ILE	2.5
1	A	274	ASN	2.4
1	A	174	LEU	2.4
1	A	26	MET	2.4
1	A	382	ASP	2.4
1	A	386[A]	ASN	2.3
1	A	389	VAL	2.3
1	A	259	GLN	2.3
1	A	361	SER	2.2
1	A	380	SER	2.2
1	A	376	LYS	2.2
1	A	217	GLU	2.2
1	A	256	LEU	2.2
1	A	260	GLN	2.2
1	A	304	HIS	2.2
1	A	400	GLU	2.2
1	A	269	THR	2.2
1	A	302	GLY	2.2
1	A	255	SER	2.1
1	A	272	GLU	2.1
1	A	212	LEU	2.1
1	A	375	ASN	2.1
1	A	385	ASN	2.1
1	A	295	ALA	2.1
1	A	261	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TRS	A	502	8/8	0.57	0.16	54,69,78,88	0
4	PGE	A	504	7/10	0.74	0.15	60,61,65,74	0
2	C5P	A	501	21/21	0.76	0.14	28,33,41,53	0
4	PGE	A	503	10/10	0.86	0.12	42,45,55,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around C5P A 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.