



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 12:35 PM UTC

PDB ID : 5SWH / pdb_00005swh
Title : c-Src V281C kinase domain in complex with Rao-IV-151
Authors : Merritt, E.A.; Dieter, E.M.
Deposited on : 2016-08-08
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	 17% 83% 7% 10%
1	B	286	 19% 85% 6% 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4243 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	1	0
			2052	1314	342	378	18			
1	B	261	Total	C	N	O	S	0	1	0
			2067	1325	345	380	17			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	expression tag	UNP P00523
A	249	HIS	-	expression tag	UNP P00523
A	250	MET	-	expression tag	UNP P00523
A	251	GLN	-	expression tag	UNP P00523
A	281	CYS	VAL	engineered mutation	UNP P00523
B	248	GLY	-	expression tag	UNP P00523
B	249	HIS	-	expression tag	UNP P00523
B	250	MET	-	expression tag	UNP P00523
B	251	GLN	-	expression tag	UNP P00523
B	281	CYS	VAL	engineered mutation	UNP P00523

- Molecule 2 is (2R)-3-[4-amino-5-(4-chlorophenyl)-7-(2-methoxyethyl)-7H-pyrrolo[2,3-d]pyrimidin-6-yl]-2-cyano-N-(propan-2-yl)propanamide (CCD ID: 71D) (formula: C₂₂H₂₅ClN₆O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 22	Cl 1	N 6	O 2	0	0
2	B	1	Total 31	C 22	Cl 1	N 6	O 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	34	Total O 34 34	0	0
3	B	28	Total O 28 28	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.82Å 64.14Å 74.91Å 77.94° 89.50° 89.83°	Depositor
Resolution (Å)	73.25 – 2.50 73.25 – 2.50	Depositor EDS
% Data completeness (in resolution range)	79.7 (73.25-2.50) 79.7 (73.25-2.50)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.241 , 0.287 0.245 , 0.290	Depositor DCC
R_{free} test set	1374 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.567	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4243	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 71D

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	1/2105 (0.0%)	0.95	2/2853 (0.1%)
1	B	0.81	0/2120	0.95	1/2875 (0.0%)
All	All	0.81	1/4225 (0.0%)	0.95	3/5728 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	309	GLN	N-CA	6.10	1.54	1.46

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	264	ARG	N-CA-C	7.25	120.60	111.69
1	B	309	GLN	N-CA-C	7.01	120.72	111.75
1	A	354	MET	N-CA-C	5.65	118.37	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2052	0	2017	12	1
1	B	2067	0	2033	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	31	0	0	0	0
2	B	31	0	0	1	0
3	A	34	0	0	1	0
3	B	28	0	0	0	0
All	All	4243	0	4050	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:314:MET:HE1	2:B:600:71D:CL	2.31	0.67
1:A:466:MET:HE2	1:A:471:VAL:HG22	1.84	0.59
1:B:466:MET:HE2	1:B:471:VAL:HG22	1.85	0.59
1:A:306:ALA:O	1:A:309:GLN:HB2	2.05	0.56
1:B:451:LEU:O	1:B:451:LEU:HD23	2.05	0.55
1:A:354:MET:HB2	3:A:721:HOH:O	2.09	0.52
1:B:518:ASP:O	1:B:519:TYR:C	2.58	0.46
1:B:323:VAL:HG21	1:B:393:LEU:HD12	1.98	0.46
1:A:323:VAL:HG21	1:A:393:LEU:HD12	1.98	0.45
1:B:354:MET:CA	1:B:354:MET:HE3	2.48	0.44
1:B:264:ARG:O	1:B:266:SER:N	2.44	0.43
1:A:354:MET:HE3	1:A:354:MET:CA	2.48	0.43
1:A:346:LEU:HD21	1:A:455:LEU:HD21	2.00	0.43
1:A:262:ILE:HB	1:A:263:PRO:HD2	2.01	0.42
1:B:346:LEU:HD21	1:B:455:LEU:HD21	2.01	0.42
1:B:354:MET:HE3	1:B:354:MET:HA	2.00	0.42
1:B:467:VAL:O	1:B:471:VAL:HG23	2.19	0.42
1:A:518:ASP:O	1:A:519:TYR:C	2.62	0.42
1:A:466:MET:CE	1:A:471:VAL:HA	2.50	0.41
1:A:467:VAL:O	1:A:471:VAL:HG23	2.19	0.41
1:B:262:ILE:HB	1:B:263:PRO:HD2	2.02	0.41
1:A:526:GLN:O	1:A:527:TYR:C	2.64	0.41
1:B:466:MET:CE	1:B:471:VAL:HA	2.52	0.40
1:A:354:MET:HE3	1:A:354:MET:HA	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:397:ASN:OD1	1:A:438:ARG:NE[1_455]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/286 (89%)	243 (95%)	11 (4%)	1 (0%)	30	49
1	B	258/286 (90%)	245 (95%)	13 (5%)	0	100	100
All	All	513/572 (90%)	488 (95%)	24 (5%)	1 (0%)	43	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	309	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/245 (89%)	216 (99%)	3 (1%)	59	81
1	B	219/245 (89%)	218 (100%)	1 (0%)	81	92
All	All	438/490 (89%)	434 (99%)	4 (1%)	70	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	309	GLN
1	A	451	LEU
1	A	505	GLU
1	B	505	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	526	GLN
1	B	287	ASN
1	B	526	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	71D	A	600	1	33,33,33	1.37	1 (3%)	37,46,46	0.78	1 (2%)
2	71D	B	600	1	33,33,33	0.93	1 (3%)	37,46,46	1.42	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	71D	A	600	1	-	9/20/22/22	0/3/3/3
2	71D	B	600	1	-	4/20/22/22	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	71D	C12-C11	7.53	1.53	1.49
2	B	600	71D	C12-C11	4.90	1.52	1.49

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	600	71D	C13-C12-C11	7.18	120.68	112.78
2	B	600	71D	C19-N5-C2	-3.44	118.99	125.00
2	A	600	71D	C19-N5-C2	-2.64	120.39	125.00

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	600	71D	C20-C19-N5-C2
2	B	600	71D	C20-C19-N5-C2
2	A	600	71D	N5-C19-C20-O1
2	A	600	71D	O-C15-N4-C16
2	A	600	71D	C13-C15-N4-C16
2	B	600	71D	N5-C19-C20-O1
2	A	600	71D	C17-C16-N4-C15
2	A	600	71D	C18-C16-N4-C15
2	B	600	71D	C19-C20-O1-C21
2	B	600	71D	C11-C12-C13-C14
2	A	600	71D	C14-C13-C15-O
2	A	600	71D	C4-C11-C12-C13
2	A	600	71D	C11-C12-C13-C14


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	600	71D	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	258/286 (90%)	1.01	48 (18%) 3 2	16, 45, 112, 165	1 (0%)
1	B	261/286 (91%)	0.98	53 (20%) 3 2	14, 41, 121, 166	1 (0%)
All	All	519/572 (90%)	1.00	101 (19%) 3 2	14, 42, 121, 166	2 (0%)

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	311	ALA	6.1
1	A	306	ALA	5.3
1	B	278	PHE	5.2
1	A	308	LEU	4.9
1	A	299	PRO	4.7
1	A	404	ASP	4.7
1	A	310	GLU	4.6
1	A	298	LYS	4.6
1	A	406	GLY	4.6
1	A	313	VAL	4.5
1	A	260	TRP	4.4
1	A	307	PHE	4.4
1	B	299	PRO	4.3
1	A	278	PHE	4.1
1	B	277	CYS	4.1
1	B	279	GLY	4.0
1	B	275	GLN	4.0
1	B	267	LEU	3.9
1	A	405	PHE	3.8
1	B	334	ILE	3.7
1	B	309	GLN	3.7
1	A	275	GLN	3.6
1	B	307	PHE	3.6
1	B	304	PRO	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	273	LEU	3.6
1	A	328	VAL	3.5
1	A	286	TRP	3.5
1	B	286	TRP	3.5
1	A	309	GLN	3.4
1	A	304	PRO	3.4
1	B	301	THR	3.3
1	B	328	VAL	3.3
1	B	308	LEU	3.3
1	B	300	GLY	3.2
1	B	318	ARG	3.1
1	A	314	MET	3.1
1	B	297	LEU	3.1
1	B	302	MET	3.1
1	A	300	GLY	3.1
1	B	403	ALA	3.1
1	A	312	GLN	3.0
1	A	403	ALA	3.0
1	B	264	ARG	3.0
1	B	317	LEU	3.0
1	A	334	ILE	2.9
1	A	276	GLY	2.9
1	A	332	GLU	2.9
1	A	277	CYS	2.9
1	A	337	VAL	2.9
1	A	336	ILE	2.8
1	A	472	LEU	2.8
1	A	259	ALA	2.8
1	B	336	ILE	2.8
1	B	335	TYR	2.7
1	B	262	ILE	2.7
1	A	325	LEU	2.7
1	B	271	VAL	2.7
1	B	268	ARG	2.7
1	B	324	GLN	2.7
1	A	317	LEU	2.7
1	B	276	GLY	2.6
1	B	333	PRO	2.6
1	B	404	ASP	2.6
1	B	305	GLU	2.6
1	A	315	LYS	2.6
1	B	306	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	405	PHE	2.6
1	B	256	ALA	2.6
1	A	343	LYS	2.6
1	A	302	MET	2.5
1	B	274	GLY	2.5
1	B	265	GLU	2.5
1	B	269	LEU	2.5
1	B	263	PRO	2.5
1	A	303	SER	2.4
1	A	262	ILE	2.4
1	A	323	VAL	2.4
1	A	301	THR	2.4
1	B	330	SER	2.3
1	A	279	GLY	2.3
1	A	355	GLY	2.3
1	B	526	GLN	2.3
1	B	354	MET	2.3
1	A	526	GLN	2.3
1	A	296	THR	2.2
1	A	273	LEU	2.2
1	A	297	LEU	2.1
1	B	469	ARG	2.1
1	B	311	ALA	2.1
1	B	255	LEU	2.1
1	B	290	THR	2.1
1	A	316	LYS	2.1
1	B	285	THR	2.1
1	B	288	GLY	2.1
1	A	469	ARG	2.1
1	A	382	TYR	2.1
1	B	472	LEU	2.1
1	B	331	GLU	2.1
1	B	322	LEU	2.0
1	B	332	GLU	2.0
1	B	282	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	71D	A	600	31/31	0.86	0.18	37,68,98,102	0
2	71D	B	600	31/31	0.86	0.16	51,69,99,103	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 71D A 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 71D B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.