



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:46 PM UTC

PDB ID : 6SW6 / pdb_00006sw6
Title : Crystal structure R264H mutant of the human S-adenosylmethionine synthetase 1
Authors : Panmanee, J.; Antonyuk, S.V.; Hasnain, S.S.
Deposited on : 2019-09-19
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	9% 91% 5%
1	B	395	9% 85% 6% 10%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-adenosylmethionine synthase isoform type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	0	5	0
			2947	1854	515	560	18			
1	B	357	Total	C	N	O	S	0	11	0
			2746	1734	475	520	17			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	264	HIS	ARG	engineered mutation	UNP Q00266
B	264	HIS	ARG	engineered mutation	UNP Q00266


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	O	0	0
			2	2		
2	B	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: S-adenosylmethionine synthase isoform type-1



- Molecule 1: S-adenosylmethionine synthase isoform type-1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	108.56Å 83.62Å 87.56Å 90.00° 107.99° 90.00°	Depositor
Resolution (Å)	51.87 – 2.85 51.87 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.5 (51.87-2.85) 99.5 (51.87-2.85)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.86Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.261 , 0.288 0.258 , 0.284	Depositor DCC
R_{free} test set	884 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	58.3	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 7.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5697	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/3005	0.80	0/4070
1	B	0.55	0/2798	0.78	0/3786
All	All	0.56	0/5803	0.79	0/7856

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2947	0	2879	7	0
1	B	2746	0	2659	7	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
All	All	5697	0	5538	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:ILE:HD11	1:B:266:ILE:HD11	1.71	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:25:VAL:HG12	1:B:181:LYS:HG3	1.75	0.67
1:B:285:LYS:HD3	1:B:289:LYS:HG2	1.85	0.58
1:B:361:VAL:HG23	1:B:366:LEU:HB2	1.88	0.55
1:A:25:VAL:HG12	1:A:181:LYS:HG2	1.91	0.53
1:B:18:PHE:C	1:B:19[A]:MET:HE2	2.35	0.51
1:A:334:THR:HG21	1:A:338:SER:HB3	1.96	0.48
1:A:308:ALA:HB2	1:A:393:LEU:HD12	1.96	0.47
1:B:308:ALA:HB2	1:B:393:LEU:HD12	1.96	0.47
1:A:69:GLY:HA2	1:B:259:ALA:HB2	1.98	0.45
1:A:354:ASP:HB3	1:A:359:VAL:HG11	2.02	0.41
1:A:180:SER:HB3	1:A:207:VAL:HG23	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/395 (96%)	366 (96%)	13 (3%)	1 (0%)	36	54
1	B	349/395 (88%)	341 (98%)	8 (2%)	0	100	100
All	All	729/790 (92%)	707 (97%)	21 (3%)	1 (0%)	48	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	114	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/332 (95%)	305 (97%)	10 (3%)	34	59
1	B	292/332 (88%)	282 (97%)	10 (3%)	32	58
All	All	607/664 (91%)	587 (97%)	20 (3%)	33	58

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	LYS
1	A	112	GLN
1	A	114	SER
1	A	127	GLU
1	A	128	GLU
1	A	137	LEU
1	A	213	ILE
1	A	222[B]	LEU
1	A	237	ASP
1	A	265	LYS
1	B	32	LYS
1	B	66	LEU
1	B	94[B]	ASP
1	B	99	PHE
1	B	137	LEU
1	B	213	ILE
1	B	222	LEU
1	B	265	LYS
1	B	288	THR
1	B	334[A]	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	GLN
1	B	36	GLN
1	B	80	GLN
1	B	135	GLN
1	B	208	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	382/395 (96%)	0.75	34 (8%) 15 12	35, 55, 91, 105	4 (1%)
1	B	357/395 (90%)	0.90	36 (10%) 12 10	20, 61, 85, 105	7 (1%)
All	All	739/790 (93%)	0.83	70 (9%) 14 11	20, 58, 89, 105	11 (1%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	257	GLY	6.7
1	B	308	ALA	5.3
1	B	356	ARG	5.3
1	B	273	GLY	5.2
1	A	276	ALA	5.1
1	A	310	LEU	5.1
1	B	355	LEU	5.0
1	A	98	GLY	4.9
1	B	351	LYS	4.6
1	A	275	GLY	4.5
1	A	277	HIS	4.4
1	B	16	GLY	4.2
1	A	298	ALA	4.0
1	B	387	TRP	3.9
1	B	349	VAL	3.8
1	B	274	TRP	3.7
1	A	200	ILE	3.4
1	B	296	TYR	3.4
1	B	275	GLY	3.4
1	B	136	GLY	3.3
1	B	137	LEU	3.3
1	A	196	ILE	3.2
1	B	305	LEU	3.2
1	B	104	CYS	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	118	ALA	3.1
1	A	183	GLN	3.1
1	B	267	ILE	3.1
1	A	194	ALA	2.9
1	A	198	VAL	2.8
1	A	108	VAL	2.8
1	A	273	GLY	2.8
1	B	314	VAL	2.7
1	A	316	VAL	2.7
1	B	114	SER	2.7
1	B	17	VAL	2.6
1	A	66	LEU	2.6
1	B	301	VAL	2.6
1	B	318	VAL	2.6
1	B	358	GLY	2.5
1	A	272	GLY	2.5
1	A	315	LEU	2.4
1	A	253	GLY	2.4
1	B	190	GLN	2.4
1	B	144	ASP	2.4
1	B	362	ARG	2.4
1	A	274	TRP	2.3
1	B	288	THR	2.3
1	A	251	VAL	2.3
1	B	278	GLY	2.3
1	A	355	LEU	2.3
1	A	87	ILE	2.3
1	A	202	THR	2.2
1	A	324	VAL	2.2
1	A	120	CYS	2.2
1	B	290	VAL	2.2
1	B	316	VAL	2.2
1	A	366	LEU	2.2
1	B	300	TRP	2.2
1	B	395	PHE	2.2
1	B	289	LYS	2.2
1	A	119	GLN	2.1
1	B	99	PHE	2.1
1	A	92	TYR	2.1
1	A	64	MET	2.1
1	A	201	HIS	2.1
1	B	388	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	95	SER	2.0
1	A	77	VAL	2.0
1	A	101	PHE	2.0
1	B	335	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.