



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 02:06 AM UTC

PDB ID : 9SWY / pdb_00009swy
Title : The structure of AfUAP1 in complex with GlcNAc-1P
Authors : Yan, K.; Raimi, O.; Chen, X.
Deposited on : 2025-10-07
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	484	
2	A	484	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7589 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-N-acetylglucosamine diphosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	476	Total	C	N	O	S	0	1	0
			3703	2365	631	694	13			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	26	GLY	-	expression tag	UNP A0A229XUD0


- Molecule 2 is a protein called UDP-N-acetylglucosamine diphosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	476	Total	C	N	O	S	0	2	0
			3650	2332	615	689	14			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	GLY	-	expression tag	UNP A0A229XUD0

- Molecule 3 is 2-acetamido-2-deoxy-1-O-phosphono-alpha-D-glucopyranose (CCD ID: GN1) (formula: C₈H₁₆NO₉P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			19	8	1	9	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	128	Total	O	0	0
			128	128		
5	A	88	Total	O	0	0
			88	88		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-N-acetylglucosamine diphosphorylase



- Molecule 2: UDP-N-acetylglucosamine diphosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.83Å 136.87Å 143.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.54 – 1.72 71.54 – 1.72	Depositor EDS
% Data completeness (in resolution range)	95.6 (71.54-1.72) 95.6 (71.54-1.72)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.72Å)	Xtriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.210 , 0.252 0.220 , 0.265	Depositor DCC
R_{free} test set	1984 reflections (1.72%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.390	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.002 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7589	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GN1, YCM, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.59	0/3775	0.96	1/5097 (0.0%)
2	A	0.59	0/3735	0.98	3/5058 (0.1%)
All	All	0.59	0/7510	0.97	4/10155 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	506	GLU	CB-CG-CD	6.55	123.74	112.60
2	A	279	ASP	CA-CB-CG	5.94	118.54	112.60
2	A	326	ASP	CA-CB-CG	5.15	117.75	112.60
2	A	64	HIS	CB-CA-C	5.03	119.14	110.79

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	178	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3703	0	3680	23	0
2	A	3650	0	3549	28	0
3	B	19	0	0	0	0
4	A	1	0	0	1	0
5	A	88	0	0	2	0
5	B	128	0	0	0	0
All	All	7589	0	7229	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:601:CL:CL	5:A:722:HOH:O	2.32	0.84
2:A:385:ILE:H	2:A:402:ASN:HD21	1.28	0.81
1:B:504:PHE:CZ	1:B:506:GLU:HG2	2.17	0.79
1:B:27:PRO:O	1:B:28:SER:HB3	1.83	0.77
2:A:471:THR:HG21	2:A:476:VAL:HG23	1.68	0.76
2:A:402:ASN:H	2:A:402:ASN:HD22	1.39	0.70
1:B:106:LEU:HD21	1:B:428:ARG:HG2	1.76	0.67
1:B:345:GLN:O	1:B:347:ASP:N	2.27	0.66
2:A:341:LYS:HE2	2:A:346:PRO:O	2.00	0.60
1:B:345:GLN:C	1:B:347:ASP:H	2.09	0.60
2:A:471:THR:C	2:A:472:GLU:HG3	2.26	0.60
1:B:344:LYS:O	1:B:345:GLN:C	2.47	0.57
1:B:189:TYR:CE1	1:B:218:ILE:HD13	2.39	0.57
2:A:385:ILE:H	2:A:402:ASN:ND2	1.99	0.57
1:B:141:ARG:O	1:B:141:ARG:HD2	2.05	0.57
1:B:344:LYS:HE2	1:B:344:LYS:C	2.31	0.56
1:B:142:LEU:HD21	1:B:436:LEU:HG	1.88	0.55
1:B:27:PRO:O	1:B:28:SER:CB	2.50	0.55
1:B:452:ARG:O	1:B:456:SER:HB3	2.09	0.53
1:B:140:THR:O	1:B:143:GLY:N	2.41	0.53
2:A:471:THR:C	2:A:472:GLU:CG	2.84	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:471:THR:CG2	2:A:476:VAL:HG23	2.42	0.49
2:A:325:PRO:HD3	2:A:419:LEU:HD13	1.94	0.48
2:A:402:ASN:HD22	2:A:402:ASN:N	2.10	0.47
2:A:179:ILE:HD13	5:A:739:HOH:O	2.15	0.46
2:A:30:GLU:OE1	2:A:34:GLN:NE2	2.48	0.46
1:B:504:PHE:CZ	1:B:506:GLU:CG	2.95	0.46
1:B:27:PRO:HD2	1:B:59:ARG:HH22	1.80	0.46
2:A:198:LYS:N	2:A:199:PRO:HD2	2.30	0.46
1:B:142:LEU:HD12	1:B:142:LEU:N	2.31	0.46
1:B:227:CYS:HB3	1:B:385:ILE:HD13	1.98	0.46
1:B:309:ARG:HH11	1:B:309:ARG:CG	2.30	0.45
2:A:297:LYS:HB2	2:A:299:VAL:HG13	1.97	0.45
2:A:138:GLN:NE2	2:A:145:SER:O	2.41	0.44
2:A:218:ILE:HD12	2:A:218:ILE:N	2.32	0.44
1:B:501:ALA:HA	1:B:502:PRO:C	2.43	0.44
2:A:319:LEU:HD11	2:A:328:VAL:CG2	2.48	0.43
1:B:286:ALA:O	1:B:287:ASP:C	2.62	0.43
1:B:500:LYS:O	1:B:503:ALA:HB2	2.19	0.43
2:A:125:ALA:HB1	2:A:180:SER:HB3	2.00	0.43
2:A:476:VAL:HG22	2:A:477:GLY:H	1.84	0.42
1:B:106:LEU:CD2	1:B:428:ARG:HG2	2.48	0.42
2:A:302:ALA:HB3	2:A:360:TYR:HB3	2.00	0.42
2:A:319:LEU:HD11	2:A:328:VAL:HG23	2.02	0.42
2:A:84:PRO:C	2:A:86:LYS:H	2.27	0.41
2:A:414:PHE:HB2	2:A:415:PRO:HD3	2.02	0.41
2:A:44:GLN:O	2:A:47:VAL:HG22	2.21	0.41
2:A:305:VAL:HG11	2:A:353:ALA:HB1	2.03	0.41
2:A:237:GLU:OE1	2:A:241:LYS:NZ	2.34	0.41
2:A:498:GLU:C	2:A:499:ILE:HD12	2.45	0.41
1:B:36:ARG:HA	1:B:48:PHE:CE1	2.56	0.40
2:A:458:GLY:HA3	2:A:478:VAL:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	470/484 (97%)	460 (98%)	7 (2%)	3 (1%)	21	9
2	A	470/484 (97%)	451 (96%)	17 (4%)	2 (0%)	30	16
All	All	940/968 (97%)	911 (97%)	24 (3%)	5 (0%)	24	11

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	28	SER
1	B	346	PRO
1	B	345	GLN
2	A	55	GLN
2	A	88	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	392/407 (96%)	377 (96%)	15 (4%)	29	8
2	A	380/408 (93%)	370 (97%)	10 (3%)	40	18
All	All	772/815 (95%)	747 (97%)	25 (3%)	35	11

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	99	ASP
1	B	111	LYS
1	B	122	LYS
1	B	140	THR
1	B	141	ARG
1	B	174	LEU
1	B	307	ARG
1	B	309	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	335	LYS
1	B	344	LYS
1	B	348	VAL
1	B	388	ILE
1	B	419	LEU
1	B	456	SER
1	B	506	GLU
2	A	30	GLU
2	A	51	VAL
2	A	62	LEU
2	A	122	LYS
2	A	134	MET
2	A	231	GLU
2	A	277[A]	CYS
2	A	277[B]	CYS
2	A	380	VAL
2	A	402	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	33	GLN
1	B	34	GLN
1	B	61	GLN
1	B	207	HIS
1	B	222	GLN
1	B	230	ASN
1	B	378	HIS
2	A	33	GLN
2	A	165	GLN
2	A	207	HIS
2	A	230	ASN
2	A	402	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	YCM	B	277	1	7,9,10	0.40	0	5,10,12	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	B	277	1	-	1/6/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	277	YCM	SG-CD-CE-OZ1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GN1	B	601	-	18,19,19	0.66	0	28,28,28	0.90	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GN1	B	601	-	-	0/10/31/31	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	GN1	C4-C3-C2	-2.26	107.11	110.40


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	B	475/484 (98%)	0.61	40 (8%)	17 22	21, 41, 74, 120	1 (0%)
2	A	476/484 (98%)	0.87	65 (13%)	6 10	22, 47, 78, 114	2 (0%)
All	All	951/968 (98%)	0.74	105 (11%)	10 15	21, 44, 75, 120	3 (0%)

All (105) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	345	GLN	7.1
1	B	87	ALA	5.7
1	B	476	VAL	5.3
2	A	396	PHE	5.0
2	A	476	VAL	5.0
1	B	343	PRO	4.9
2	A	85	PRO	4.8
2	A	179	ILE	4.3
2	A	333	ILE	4.3
1	B	346	PRO	4.2
2	A	343	PRO	4.1
1	B	395	PHE	4.1
2	A	395	PHE	4.0
1	B	471	THR	3.9
1	B	27	PRO	3.9
1	B	26	GLY	3.8
1	B	138	GLN	3.7
1	B	348	VAL	3.7
1	B	91	PRO	3.5
2	A	473	GLY	3.5
2	A	508	GLU	3.5
2	A	337	THR	3.4
2	A	27	PRO	3.4
1	B	137	GLY	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	392	THR	3.3
1	B	344	LYS	3.3
2	A	398	PRO	3.3
2	A	87	ALA	3.2
2	A	388	ILE	3.2
2	A	145	SER	3.2
1	B	140	THR	3.2
2	A	257	LEU	3.1
1	B	92	ALA	3.1
2	A	380	VAL	3.1
2	A	389	LYS	3.1
1	B	90	GLY	3.0
2	A	346	PRO	3.0
2	A	51	VAL	3.0
2	A	91	PRO	2.9
1	B	470	ILE	2.9
2	A	443	GLY	2.9
1	B	392	THR	2.8
1	B	139	GLY	2.8
2	A	142	LEU	2.8
2	A	88	ASP	2.8
2	A	90	GLY	2.8
2	A	86	LYS	2.8
2	A	386	PRO	2.7
2	A	338	ALA	2.7
2	A	146	ALA	2.7
2	A	56	THR	2.6
1	B	143	GLY	2.6
1	B	142	LEU	2.6
1	B	468	ILE	2.6
1	B	507	LYS	2.6
1	B	347	ASP	2.6
2	A	54	LEU	2.6
2	A	397	LYS	2.5
2	A	499	ILE	2.5
2	A	393	GLY	2.5
2	A	233	LYS	2.5
2	A	50	PHE	2.5
1	B	94	LEU	2.4
2	A	340	ALA	2.4
2	A	55	GLN	2.4
2	A	194	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	33	GLN	2.4
2	A	314	SER	2.4
1	B	337	THR	2.4
1	B	141	ARG	2.4
2	A	330	TYR	2.4
1	B	309	ARG	2.3
1	B	100	ILE	2.3
2	A	242	VAL	2.3
1	B	52	ASP	2.3
2	A	419	LEU	2.3
2	A	344	LYS	2.2
2	A	32	PHE	2.2
2	A	441	GLY	2.2
1	B	419	LEU	2.2
2	A	136	GLY	2.2
2	A	240	PHE	2.2
2	A	93	SER	2.2
1	B	340	ALA	2.2
2	A	140	THR	2.2
2	A	401	PRO	2.2
2	A	394	GLU	2.1
2	A	496	GLY	2.1
2	A	195	PRO	2.1
1	B	388	ILE	2.1
2	A	468	ILE	2.1
2	A	64	HIS	2.1
1	B	467	GLY	2.1
2	A	139	GLY	2.1
1	B	105	ILE	2.1
1	B	311	ALA	2.1
2	A	57	GLY	2.1
2	A	478	VAL	2.1
2	A	141	ARG	2.1
2	A	400	LYS	2.1
1	B	497	ARG	2.0
2	A	349	LEU	2.0
2	A	345	GLN	2.0
1	B	49	ALA	2.0
1	B	86	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	YCM	B	277	10/11	0.96	0.10	28,35,52,55	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	A	601	1/1	0.86	0.15	82,82,82,82	0
3	GN1	B	601	19/19	0.97	0.05	29,36,41,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GN1 B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.